



Annual Research & Review in Biology

23(4): 1-7, 2018; Article no.ARRB.38144
ISSN: 2347-565X, NLM ID: 101632869

Response of Broiler Chickens to *Carica papaya* and *Talinium triangulare* Leaf Meal under Normal and Subnormal Diets

B. E. Agboola¹, A. D. Ologhobo^{1*}, I. O. Adejumo^{2*} and G. O. Adeyemo¹

¹Animal Biochemistry and Nutrition Research Unit, Department of Animal Science, University of Ibadan, Ibadan, Nigeria.

²Animal Nutrition, Biotechnology and Food Safety Laboratory, Department of Animal Science, Federal University Gashua, Nigeria.

Authors' contributions

This work was carried out in collaboration among all authors. Author ADO designed the study. Authors ADO and GOA supervised the study. Author BEA carried out the feeding trial. Authors BEA and IOA carried out the statistical analyses. Author IOA handled literature and wrote the first draft. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/ARRB/2018/38144

Editor(s):

(1) George Perry, Dean and Professor of Biology, University of Texas at San Antonio, USA.

Reviewers:

(1) Hana Zakaria, The University of Jordan, School of Agriculture, Jordan.

(2) Mahmoud I. El-Kelawy, Assiut University, Egypt.

(3) Juan Carlos Troiano, University of Buenos Aires, Argentina.

Complete Peer review History: <http://www.sciencedomain.org/review-history/23019>

Original Research Article

Received 14th September 2017
Accepted 14th December 2017
Published 5th February 2018

ABSTRACT

The use of locally available and cheap feed ingredients has received particular attention as a viable alternative to the use of conventional feedstuffs in developing countries. Vegetable-based feeds are rich sources of essential plant amino acids, vitamins, minerals, and antioxidants. Further to the rich contents mentioned, it has been established that green vegetable leaves are cheap and abundant sources of protein because of their ability to synthesize amino acids from a wide range of available primary materials. The aim of this study was to investigate the response of broiler chickens to *Carica papaya* and *Talinium triangulare* under normal and subnormal diets. In this study, the effects of *C. papaya* leaf meal and *T. triangulare* chopped leaves were assessed on growth performance and haematological parameters of broiler chickens. Two hundred and seventy 1-d old Arbor acres broiler

*Corresponding author: E-mail: tonyologhoobo@yahoo.com, smogisaac@gmail.com;

chicks were used for the study. Each treatment had 5 replicates with 6 birds per replicate. Normal diets and subnormal diets were compounded with varying amounts of ingredients. The test ingredients were supplied at the rate of 10% of the main diets. The design of the experiment was a completely randomized design. All the treatments with subnormal diets obtained higher feed intake than treatments with normal diets. Birds on subnormal diet mixed with *C. papaya* leaf meal (24.03%) obtained the highest packed cell volume. The haemoglobin concentrations followed a similar pattern. Normal diets compared with control for feed conversion ratio.

Keywords: Broiler chickens; *Carica papaya*; haematology; leaf meal; *Talinum triangulare*; performance.

1. INTRODUCTION

Feeding in poultry production is based on the science of nutrition in which nutritional requirements are mostly known. High cost of feeds and scarcity of essential raw materials have necessitated poultry farmers to look out for systems which could help to identify feed ingredients of lower cost and sound biological values that can partly supplement the conventional protein and energy sources in broiler diets.

The use of locally available and cheap feed ingredients has received particular attention as a viable alternative to the use of conventional feedstuffs [1-5]. Vegetable-based feeds are rich sources of essential plant amino acids, vitamins, minerals, and antioxidants [6]. Further to the rich contents mentioned, it has been established that green vegetable leaves are the cheapest and most abundant sources of protein because of their ability to synthesize amino acids from a wide range of available primary materials [7].

Talinum triangulare is an agricultural product whose nutritional importance on livestock has been ignored probably because little is known about its potential in livestock production. *Talinum triangulare* contributes significantly to the nutritional security in some rural communities in Africa. *Carica papaya* is popular in the tropics and sub-tropics because of its easy cultivation, rapid growth, quick economic returns and easy adaptation to diverse soils and climates [8,9]. Exposing meat type broilers to a partial scavenging lifestyle may enhance their nutritional requirements. Hence, this study evaluated the performance characteristics and haematological response of broiler chickens scavenging on *C. papaya* leaf meal and *Talinum triangulare* chopped leaves under normal and subnormal diets.

2. MATERIALS AND METHODS

2.1 Experimental Site, Source, and Processing of Test Ingredients

The study was carried out at the Teaching and Research Farm, University of Ibadan, Nigeria. Fresh pawpaw leaves (*Carica papaya*) and water leaves (*Talinum triangulare*) were obtained from the farm premises and identified at the Department of Botany, University of Ibadan. Pawpaw leaves were separated from the stalk and sundried for 5 days on a clean concrete floor. By the 5th day, they had become crispy at constant weight. The leaves were crumbled by hand and were stored in an air-tight container. Water leaves were supplied fresh on daily basis in a chopped form. The proximate composition of the test ingredients was carried out in triplicates according to the method of AOAC [10].

2.2 Management of Experimental Animals

The poultry house was cleaned, washed and disinfected using Povidine Iodophore™ (Animal Care Services Konsult Nig. Ltd) and allowed without stocking for 2 weeks. Two hundred and seventy 1-d old Arbor acres broiler chicks were purchased from a commercial hatchery in Ibadan. After brooding, the birds were randomly allotted to 9 treatments in a complete randomized design in a deep litter system. Each treatment had 5 replicates with 6 birds per replicate. Two main rations were compounded with varying amount of ingredients. One was tagged 'Normal diet' while the other was tagged 'Subnormal diet' (Table 1). The normal diet had 21% crude protein while the subnormal diet had a very low crude protein (15.57%) and metabolizable energy lower than the recommended (Table 1). The test ingredients were supplied at the rate of 10% of the main diets. The treatment diets were as follows:

- T₁= Control diet (Normal diet – leaf meal);
- T₂= Normal diet mixed with *C. papaya* leaf meal
- T₃= Subnormal diet mixed with *C. papaya* leaf meal
- T₄= Normal diet separate, *C. papaya* leaf meal separate
- T₅= Subnormal diet separate, *C. papaya* leaf meal separate
- T₆= Normal diet mixed with chopped *T. triangulare*
- T₇= Subnormal diet mixed with chopped *T. triangulare*
- T₈= Normal diet separate, chopped *T. triangulare* separate
- T₉= Subnormal diet separate, *T. triangulare* separate

The experimental diets (Table 1) were fed for 28 days and the birds had access to the feed and cool drinking water *ad libitum*. All routine vaccinations and bio-security measures were carried out as recommended. Litter materials were regularly replaced.

2.3 Data Collection

Feed intake was obtained daily by subtracting left-over feed from feed served. Weekly body weight gain was measured using weekly weigh-back mechanism, by subtracting the present week's weight from that of the previous week. Feed conversion ratio (FCR) was obtained by

dividing the average total feed intake by the average total body weight gain.

2.4 Haematological Studies

Blood samples were collected from the experimental animals through making incision on the jugular veins and evacuating the blood into heparinised bottles. The haematological parameters were analysed for at the Physiology Laboratory, Department of Animal Science, University of Ibadan.

2.5 Statistical Analysis

The data obtained were subjected to one-way analysis of variance (ANOVA) and the means were compared using the Duncan's Multiple range test as outlined by Steel and Torrie [11].

3. RESULTS

The proximate composition of the experimental diets, experimental leaves, mixed feeds and performance characteristics of broiler chickens fed *T. triangulare* and *C. papaya* leaf meal under normal and subnormal diets are presented in Tables 2, 3, 4 and 5 respectively. No significant differences were observed across the treatments for initial body weights and feed intake. The highest average total body weight gain (2901.05g) was obtained by animals on T₈, with

Table 1. Gross composition of finisher experimental diets experimental diets (normal and subnormal)

Ingredients (%)	Normal	Subnormal
Maize	55.00	50.00
Soya meal	25.00	6.70
Fish meal	2.00	---
Groundnut cake	10.00	5.00
Wheat offal	4.70	35.00
Oyster shell	1.50	1.50
Bone meal	1.00	1.00
Methionine	0.15	0.15
Lysine	0.10	0.10
Premix	0.25	0.25
Salt	0.30	0.30
Total	100.00	100.00
Calculated nutrients		
Crude protein	21.90	15.57
Metabolizable Energy (Kcal/Kg)	2894.50	2672.40

Vitamins and Mineral premix for Broilers and Chicks (2.5kg/ton) Vitamin A12,500.00iu. Vitamin D3 2,5000.00iu. Vitamin E= 35,000iu Vitamin K=200g, Thiamine B1=200g, Ribloflavin B2= 5.00g, Niacin B3 = 40.00g, D-Calpan B5 11.00g, Pyridoxine B6=400g, Biotin=0.10g, Folic acid=1.50g Vitamin B12=0.012g, Manganese= 70.00g, Zinc=50.00g, Copper = 6.00g: Iron= 40.00g, Iodine = 1.00g: Cobalt = 11.25g. Selenium = 0.15g: Choline chloride = 500.00g

normal feed and *T. triangulare* leaf separated. All the treatments with subnormal diet except T₇ (i.e. T₃, T₅, T₉) had lower body weight gain ($P = 0.05$). Birds fed the control diet with no experimental leaf had a final body weight gain (2830.05g) which was significantly ($P = 0.05$) lower than those of T₄, T₆ and T₈ (2851.03g, 2880.02g and 2901.05g respectively). All the treatments with subnormal diets obtained higher feed intake than the treatments with normal diets. However, birds in T₇ (subnormal feed mixed with *T. triangulare* leaf) obtained the highest average total feed intake (6251g). Also, there was no significant difference in the feed efficiency of all the birds fed subnormal diets (T₃ (0.43), T₅ (0.42), T₇ (0.44), T₉ (0.42)).

Table 2. Proximate compositions of experimental diets

Nutrients (%)	Normal	Subnormal
Dry matter	92.00	90.00
Crude protein	20.63	15.50
Crude fibre	3.62	6.05
Ether extract	4.90	3.80
Ash	4.80	5.00
Metabolizable energy (Kcal/Kg)	2850.00	2610.00

The haematological parameters of broiler chickens fed pawpaw leaf meal and water leaves are shown in Table 6. Birds on T₃ (24.03%) obtained the highest PCV content, followed by those on T₂ (21.93%) while T₆ (17.03%) and T₈ (17.00%) obtained the least values. The haemoglobin concentrations were significantly higher for birds on T₃ than those of the control group, while those on T₂, T₄, T₅, T₇ and T₉ compared with the control group. For white blood cell counts (X10³/mm³) birds on T₇ (31.20) and T₉ (31.12) obtained the highest values followed by those on control diet (27.85). Birds on T₅ (4.6410⁶/mm³) obtained the highest mean value of red blood cell counts followed by those on T₃ (4.0510⁶/mm³) while those on T₈ (1.6110⁶/mm³) obtained the least values.

4. DISCUSSION

The crude protein of water leaf obtained in this study was close to reported by Oguntona [12] but lower than the value reported by Mohammed and Mann [13] for fresh water leaf. The value obtained for crude fibre content of water leaf in the present study was similar to the value obtained by Mohammed and Mann [13]. The value reported in the present study is lower than

those of *C. olerius*, *A. cruentus* and *C. argenta* as reported by Chionyedua et al. [14]. The fibre content of water leaf and other vegetables could be a beneficial factor in aiding effective digestion process. The mean value obtained for ether extract by water leaf is similar to the value reported by Ogbonnaya and Chinedum [15]. Crude protein, ether extract, crude fibre and ash content of the pawpaw leaf are similar for the same leaf as reported in the previous finding Unigwe [16].

Table 3. Proximate composition of experimental leaves

Nutrients (%)	Water leaf (wet)	Pawpaw leaf (sundried)
Crude protein	2.11	23.90
Ether extract	1.50	0.38
Crude fibre	1.03	10.50
Ash	1.38	7.61
Dry matter	9.80	91.60

4.1 Performance Characteristics

The average weight of 3.0Kg at the end of the experiment justifies the recommendation of 1.60 to 2.00Kg body weight at 9 weeks old by Oluoyemi and Robert [17] and 1.80Kg to 2.0Kg/bird at 9 weeks projected by Olomu and Offiong [18]. Generally, birds on normal diets had higher live weight and body weight gain than those on sub normal diet because the crude protein, ether extract, metabolizable energy and other nutrients in the normal diets fell within the recommended amount for broiler production and performed better than that of subnormal diet. The highest feed consumption was obtained in the treatments with subnormal diets. The feed consumption pattern is in agreement with the report of Leeson [19] that birds eat more feed as the nutrient level of a diet is reduced. The lowest FCR obtained for T₆ and T₈ is likely due to contribution of the vegetable to proper digestion process. Vegetables act as buffering agents for acidic substances produced during the digestion process [20]. This is likely responsible for the highest weight gain. Generally, birds fed with normal diet with water leaf had better weight gain than birds fed with normal diet with pawpaw leaf. Water leaf extract has appetizing and digestion stimulating properties as earlier observed by Langhout [21]. Nonethelss, Nworgu et al.[22] earlier reported that dietary inclusion of water leaf meal (sundried) in broiler starter diets depressed feed intake.

Table 4. Proximate composition of the mixed diets

Nutrients (%)	Normal diet + water leaf	Subnormal diet + water leaf	Normal diet + pawpaw leaf	Subnormal diet + pawpaw leaf
Dry matter	93.00	90.00	92.00	90.40
Crude fibre	4.52	5.80	4.85	5.95
Ether extract	5.40	3.60	4.60	3.20
Ash	5.00	6.05	7.63	6.46
Crude protein	20.81	16.31	20.51	16.28
Metabolizable energy (Kcal/Kg)	2815.00	2623.00	2802.00	2615.00

Table 5. Performance characteristics of the experimental birds at different treatments

Parameters	T ₁	T ₂	T ₃	T ₄	T ₅	T ₆	T ₇	T ₈	T ₉	SEM
Av. Initial body wt. (g)	320.55	320.01	320.00	319.87	320.04	320.08	319.95	319.00	310.00	5.33
Av. final body wt. (g)	3150.60 ^d	3110.00 ^e	2980.02 ^g	3170.90 ^c	2950.10 ^h	3200.10 ^b	3040.00 ^f	3220.05 ^a	2940.02 ⁱ	42.71
Av. total body wt. gain (g)	2830.05 ^d	2789.99 ^e	2660.02 ^g	2851.03 ^c	2630.06 ^h	2880.02 ^b	2720.05 ^f	2901.05 ^a	2630.02 ^h	51.89
Av. Total feed intake (g)	5860.01 ^e	5790.05 ^h	6210.10 ^d	5808.08 ^g	6219.00 ^c	5830.02 ^f	6251.00 ^a	5860.04 ^e	6250.00 ^b	55.52
Feed conversion ratio	2.07 ^{bc}	2.08 ^{bc}	2.34 ^{ab}	2.04 ^c	2.37 ^a	2.02 ^c	2.30 ^{abc}	2.02 ^c	2.38 ^a	0.42
Feed efficiency	0.48 ^{abc}	0.48 ^{abc}	0.43 ^{cd}	0.49 ^{ab}	0.42 ^d	0.50 ^a	0.44 ^{bcd}	0.50 ^a	0.42 ^d	0.09

^{a, b, c}Means with different superscripts on the same row differ significantly ($p = 0.05$), T₁= Control diet (Normal diet with no leaf included); T₂= Normal diet mixed with pawpaw leaf; T₃= Subnormal diet mixed pawpaw leaf; T₄= Normal diet separate, pawpaw leaf separate; T₅= Subnormal diet separate, pawpaw leaf separate; T₆= Normal diet mixed with water leaf; T₇= Subnormal diet mixed with water leaf; T₈= Normal diet separate, water leaf separate; T₉= Subnormal diet separate, water leaf separate

Table 6. Effect of waterleaf and pawpaw leaf on haematology of experimental broilers

Parameters	T ₁	T ₂	T ₃	T ₄	T ₅	T ₆	T ₇	T ₈	T ₉	SEM
PCV (%)	20.01 ^d	21.93 ^b	24.03 ^a	19.90 ^d	21.02 ^c	17.03 ^f	17.95 ^e	17.00 ^f	18.08 ^e	3.78
Hb (g/100ml)	6.70 ^{bc}	7.30 ^b	7.98 ^a	6.73 ^{bc}	7.08 ^b	6.00 ^d	6.17 ^{cd}	6.01 ^d	6.12 ^{cd}	1.28
RBC (10 ⁶ /mm ³)	2.47 ^d	3.43 ^c	4.05 ^b	2.47 ^d	4.64 ^a	1.62 ^g	1.85 ^e	1.61 ^g	1.78 ^f	0.51
WBC (X10 ³ /mm ³)	27.85 ^b	25.70 ^c	18.59 ^e	27.88 ^b	16.85 ^f	22.03 ^d	31.20 ^a	22.03 ^d	31.12 ^a	4.7
MCV(fl)	81.00 ^d	63.8 ^e	59.30 ^f	80.80 ^d	45.30 ^g	105.20 ^a	97.20 ^c	105.90 ^a	101.70 ^b	15.82
MCHC (%)	33.48 ^{cd}	32.37 ^e	33.22 ^d	33.83 ^b	33.68 ^{bc}	35.25 ^a	33.89 ^b	35.34 ^a	33.85 ^b	6.52
MCH (Pg/cell)	27.18 ^d	20.68 ^e	19.70 ^f	27.23 ^d	15.24 ^g	37.00 ^a	33.52 ^c	37.41 ^a	34.43 ^b	5.39

^{a, b, c}Means with different superscripts on the same row differ significantly ($p = 0.05$), PCV= Packed cell volume; Hb= Haemoglobin; RBC= Red blood cell; WBC= White blood cell, T₁= Control diet (Normal diet with no leaf included); T₂= Normal diet mixed with pawpaw leaf; T₃= Subnormal diet mixed pawpaw leaf; T₄= Normal diet separate, pawpaw leaf separate; T₅= Subnormal diet separate, pawpaw leaf separate; T₆= Normal diet mixed with water leaf; T₇= Subnormal diet mixed with water leaf; T₈= Normal diet separate, water leaf separate; T₉= Subnormal diet separate, water leaf separate

4.2 Haematological Indices

Blood represents a means of assessing clinical and nutritional health status of animals in feeding trial and the haematological parameters and most commonly used haematological indices in nutritional studies include PCV, RBC, HBC, MCHC, MCV and clotting time [23,24]. Highest values of packed cell volume, red blood cell, and haemoglobin were recorded in all the pawpaw leaf-included treatments and the findings are similar to the report by Nodu et al. [25]. This is likely attributed to the chemical and nutritional components found in pawpaw leaves. Pawpaw leaf is reported to be rich in minerals which are useful in the coagulation of blood, proper functioning of the heart and nervous system, useful as both antiseptic and a tonic or blood purifier, as well as the normal contraction of muscles [26]. The pawpaw leaf had better impact on the haematological parameters of the experimental birds than water leaf. The contribution of pawpaw leaf to the pigmentation of the blood was noticeable through recorded highest RBC, Hb and PCV in the pawpaw leaf-supplemented treatments. *C. papaya* has been reported to increase in platelet counts in the patients of dengue [27,28].

5. CONCLUSION

This study concludes that the contribution of pawpaw leaf to the pigmentation of the blood was noticeable through recorded highest RBC, Hb and PCV in the pawpaw leaf-supplemented treatments. Sundried pawpaw leaf meal may however contain some secondary metabolites which raised the white blood cell counts of the experimental animals, which may be investigated through further studies.

ETHICAL APPROVAL

As per international standard or university standard written ethical approval has been collected and preserved by the author(s).

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Nwakpu PE, Omeke SI, Alaku SO. The response of weaner pigs to diets containing fish meal and blood meal as separate sources of animal protein. *Tropical Journal of Animal Science*. 2000;3:45–48.
2. Odunsi AA. Assessment of *Lablab purpureus* leaf meal as a feed ingredient and Yolk colouring agent in the diet of layers. *International Journal of Poultry Sci*. 2003;2:71-74.
3. Adejumo IO, Ologhobo AD. Haematological response of broiler finishers fed differently processed taro cocoyam (*Colocasia esculenta* [(L.) Schott]). *Agricultura Tropica Et Subtropica*. 2012;45(3):112-116.
4. Ologhobo AD, Adejumo IO. Effects of differently processed taro cocoyam (*Colocasia esculenta* [(L.) Schott]) on growth performance and carcass characteristics of broiler finishers. *International Journal of Agri Science*. 2011;1(4):244-248.
5. Ekenyem BU, Madubuike FN. An assessment of *Ipomoea asarifolia* leaf meal as feed ingredient in broiler chick production. *Pakistan Journal of Nutrition*. 2006;5(1):46-50.
6. Omenka RO, Anyasor GN. Vegetable-based feed formulation on poultry meat quality. *Afr J Food Agric Nutr Dev*. 2010;10(1):2001-2011.
7. Fasuyi AO. Nutritional potential of some tropical vegetable leaf meals: Chemical characterisation and functional properties. *Afr J Biotechnol*. 2006;5(1):49-53.
8. Harkness RW. Papaya growing in Florida. *Bulletin of Florida Agric Exp Sta*. 1967;7: 5100.
9. Campbell CW. Papaya-tropical fruits and nuts, In: F. W. Martin (Ed.), *Handbook of Tropical Food Crops*, CRC Press Inc., Boca Raton FL.; 1984.
10. Association of Official Analytical Chemists (AOAC). Washington DC. 15th Edition; 1990.
11. Steel RGD, Torrie JH. Principles and procedures of statistics, Second Edition, New York: McGraw-Hill Book Co.; 1980.
12. Oguntona T. In: Osagie AU, Eka OU (Eds). *Nutritional quality of plant food*. Nigeria Post Harvest Research Unit. 2008;133.
13. Mohammed G, Mann A. Evaluation of the nutritional values of dry season Fadama vegetables in Bida, Nigeria. *African Journal of Food Science*. 2012;6(11):302-307.
14. Chionyedua TO, Anuoluwa MO, Adedoja DW. The proximate and mineral

- composition of three leafy vegetables commonly consumed in Lagos, Nig. Afr J Biotechnol. 2009;3(6):102-107.
15. Ogbonnaya EC, Chinedum EK. Bioactive constituents and in vitro antioxidant capacity of water Leaf (*Talinum triangulare*) as affected by domestic cooking, European Journal of Medicinal Plants. 2013;3(4):540-551.
 16. Unigwe CR, Okorafor UP, Ogbu UM, Nwufoh OC. The nutritive profile of sun-dried paw-paw (*Carica papaya*) leaf meal and its effect on the growth performance of broiler chickens. International Journal of Pure and Applied Sciences and Technology. 2014;20(2):72-78.
 17. Oluyemi JA, Roberts FA. Poultry production in warm wet climates. Macmillan, London. 1979;198.
 18. Olomu JM, Offiong SA. Effects of protein, energy level, and time change from starter to finisher ration on the performance of broiler chicks in the tropics. British Poultry Science. 1978;59:825-835.
 19. Steeve Leeson. Is feed efficiency still a useful measure of broiler performance? A publication of Department of Animal and Poultry Science, University of Guelph; 2000.
 20. Fayemi PO. Nigerian Vegetables, 1st Edition, Heinemann Educational Books Nigeria. 1999;1-8.
 21. Langhout P. New additives for broiler chickens. World Poultry Science. 2000;16(1):22-27.
 22. Nworgu FC, Alikwe PCN, Egbunike GN, Ohimain EI. Performance and nutrient utilization of broiler chickens fed water leaf meal supplement. International Journal of Farming and Allied Science. 2015;3(8):876-883.
 23. Olorede BR, Longe OG. Effects of replacing palm kernel cake with shear butter cake on quality characteristics, haematology and serum chemistry of laying hens. Nigerian Journal of Animal Production. 2000;27:19-23.
 24. Adeyemi OA, Fasina OE, Balogun MO. Utilization of full fat *Jatropha* seeds in broiler diet: Effect on hematological parameters and blood chemistry. Proceedings of the 25th conference of Nigerian Society for Animal Production held at Michael Okpara University of Agriculture, Umudike, Nigeria. 2000;108-109.
 25. Nodu MB, Okpeku M, Abezi AE. Heamatological Characteristics of Young Rabbits fed a mixture of *Alchornea cordifolia*/Pawpaw leaf meal. Journal of Agriculture and Veterinary Science (IOSR-JAVS). 2014;7(1):92-96.
 26. ATTA KB. The power of Garlic. Cardiovascular Disease Prevention Association, Buea, Cameroon. 1999, P72.
 27. Yunita F, Hanani E, Kristianto J. The effect of *Carica papaya* L. leaves extract capsules on platelets count and hematocrit level in dengue fever patient. Int J Med Aromat Plants. 2012;2:573-8.
 28. Jaykaran Charan, Deepak Saxena, Jagdish Prasad Goyal, Sandul Yasobant. Efficacy and safety of *Carica papaya* leaf extract in the dengue: A systematic review and meta-analysis. Int J Appl Basic Med Res. 2016; 6(4):249-254.

© 2018 Agboola et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<http://www.sciencedomain.org/review-history/23019>