



Productivity and Economic Feasibility of Pigeonpea Base Companion Cropping under Additive Series Planting System

Shailendra Pratap Singh ^{a++*}, C. L. Maurya ^{b#},
Naushad Khan ^{c#}, Bal Veer Singh ^{c++},
Sarvesh Kumar ^{a#} and Kaushal Kumar ^{a#}

^a Department of Soil Conservation and Water Management, C.S. Azad University of Agriculture and Technology, Kanpur (U.P.), India.

^b Department of Seed Science and Technology, C.S. Azad University of Agriculture and Technology, Kanpur (U.P.), India.

^c Department of Agronomy, C.S. Azad University of Agriculture and Technology, Kanpur (U.P.), India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JABB/2024/v27i5787

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/114014>

Original Research Article

Received: 03/02/2024

Accepted: 05/04/2024

Published: 06/04/2024

ABSTRACT

Over two consecutive Kharif seasons in 2016-2017 and 2017-18, a study was conducted at the Soil Conservation and Water Management Farm of C.S. Azad University of Agriculture and Technology in Kanpur. The study aimed to determine the yields and economic benefits (in terms of net profit) of

⁺⁺ Teaching Associate;

[#] Professor;

*Corresponding author: E-mail: shailendrapratapsingh01989@gmail.com;

the primary crops: pigeonpea, black gram, and sesame. The experimental field's soil was classified as sandy loam with a pH of 7.8, containing 0.32% organic carbon, 17.3 kg/ha of available P₂O₅, and 181.2 kg/ha of available K₂O, indicating a medium fertility level. The experiment included five different cropping treatments: sole pigeonpea, sole black gram, sole sesame, a combination of pigeonpea and black gram (1+1) in an additive series, and a combination of pigeonpea and sesame (1+1) in an additive series. The combined cultivation of pigeon pea and black gram yielded the highest productivity at 23.53 quintals per hectare (q/ha), outperforming the pigeonpea and sesame combination at 20.30 q/ha and sole pigeonpea at 17.34 q/ha. The analysis of aggregated data revealed that the highest cultivation cost was Rs. 57,876.00 per hectare for pigeonpea and black gram, followed by pigeonpea and sesame at Rs. 56,191 per hectare. The greatest gross returns were observed with the pigeonpea and black gram combination at Rs. 134,024 per hectare, with a net return of Rs. 76,148 and a benefit-cost ratio (BCR) of 1:2.32. This was followed by the pigeonpea and sesame combination, whereas sole pigeonpea cultivation resulted in a net return of Rs. 50,387 and a BCR of 1:2.01.

Keywords: BCR; companion cropping; net income; pigeonpea base; pooled data.

1. INTRODUCTION

Pigeonpea (*Cajanus cajan* L.) is an important grain legume crop of the semi-arid tropics cropping, about the area and production with 35% and 29% world, respectively available in this tract. Further, pigeonpea occupied second position after chickpea and contributed 4.2 mt. from an area of 4.43 mt. with average productivity of 960 kg/ha. In Uttar Pradesh, pigeonpea covered 2.85 lakh hectares and total production is 2.72 lakh tones during 2020 [1]. Pigeonpea grown as single crop display in efficient utilization of resources especially the space because of its slow initial growth rate, therefore, cultivation of pigeonpea as a sole crop is reported less profitable due to higher duration and wider spacing [2]. To make the cultivation of pigeonpea more viable, it is necessary to utilize the inter row space through companion cropping. Intercropping with short duration pulse like black gram, green gram, pearl millet like finger millet, foxtail millet in pigeonpea may enhance total productivity and may also provide early cash flow. Companion cropping also suppress the weed growth in between two rows of pigeonpea. Efficient of resource utilization can be increased by companion cropping and it also help in harnessing benefits of positive interaction in crop association. Growing a intercrop which produces the maximum cover, reduces soil loss, black gram is important cover crop for rainy season. The crop gives early and dense ground cover, which generally co-occurrence with peak rate of runoff.

In areas where the annual rainfall is in the range of 600-850 mm companion cropping is being recommended and practiced. The different

intercropping systems including sesame or oilseeds are followed in different parts of country. Sesame and pulse crop like black gram which are drought resistant and have a capacity to low level management conditions may be ideal combination for sustainable cropping system (Yadav et al., 2013). Companion cropping is practiced as an insurance of crop failure under rainfed condition. The main objective is higher productivity per unit area in addition to stability in production. Companion cropping system utilizes resource efficiency and their productivity is increased.

The economic resources of rainfed area are the national treasure and need proper planning to make best use of them. Therefore, sustainable management practices are urgently needed all over the world to preserve the production potential of agricultural land. Efficient management and maintenance of soil health is the being to accomplish sustainable high productivity, food security and environment safety [3].

Therefore, productivity and profitability increases through companion cropping system is the subject matter of this manuscript.

2. MATERIALS AND METHODS

An experiment was conducted under rainfed condition during two consecutive Kharif seasons of 2016-17 and 2017-18 at Soil Conservation and Water Management Farm, C.S. Azad University of Agriculture and Technology, Kanpur. The five treatments were tested i.e., pigeonpea sole planted at 60 cm apart, black gram sole planted at 30 cm apart, sesame sole planted at 30 cm

apart, pigeonpea + black gram (1+1) planted under additive series and pigeonpea + sesame (1+1) planted under additive series. The main crop was pigeonpea, while black gram and sesame were planted in the inter spaces of two rows of pigeonpea. The soil of experimental site was a typical eroded Gangetic alluvial representing Kanpur Type-1. The soil was sandy loam, having pH 7.7, organic carbon 0.33%, available N 172.00 kg/ha, available P₂O₅ 17.50 kg/ha and available K₂O 183.00 kg/ha, therefore fertility status of plant nutrients was medium. The pH was determined by Electrometric glass electrode method as discussed by Piper [4]. The organic matter was analyzed by Walkley and Black's rapid titration method [5]. The available P₂O₅ and available K₂O were analyzed by Olsen's method and Flame photometric method, respectively, [6]. The available N was analyzed by Kjeldahl's method as suggested by Subhiah and Asija [7]. The cultivar Amar (KA-32-1) of pigeonpea, cv. Shekhar-2 (KU-300) of black gram and cv. Shekhar (SH-446) of sesame were shown under intercropping. The sole crops of pigeonpea and black gram were fertilized with 20 kg N + 40 kg P₂O₅/ha, while sole and intercropped sesame were fertilized with 40 kg N + 20 kg P₂O₅ + 20 kg K₂O/ha. The recommended methods of fertilizer application were applied for obtaining better response. The recommended conservation agronomical practices were followed for raising of experimental crops during two experimental seasons. The harvesting of crops was made at complete maturity stage. The treatments were replicated thrice in a split plot design. The obtained data analyzed by standard method as suggested by Cochran and Cox [8]. The economics computed and pooled to draw valid

conclusion from the study, is the subject matter of this study.

3. RESULTS AND DISCUSSION

The aggregated data on seed yield for both the main crops and intercrops, along with the economic analysis, are shown in Table-1 and will be discussed below under relevant sections.

3.1 Total Productivity

Purusal of data available in Table 1 make it clear that the highest total productivity was weighed under pigeonpea + black gram intercropping system by 23.53 q/ha, followed by inter cropping of pigeonpea + sesame 20.30 q/ha. The order of performance of alone cropping system was pigeonpea alone (17.34 q/ha) > black gram alone (9.44 q/ha) and > sesame alone (6.32 q/ha).

3.2 Economic Study

The maximum cost of cultivation was observed under pigeonpea + black gram (1+1) additive intercropping by Rs. 57876/ha. It might be attributed to total population adjustment under both component crops of the system and their total input requirements. The highest gross return (Rs. 134024/ha), net return (Rs. 76148/ha) and BCR (1:2.32) were observed under pigeonpea + black gram intercropping followed by pigeonpea + sesame (1+1) additive intercropping. It may due to higher total productivity of the planting system. These results are similar, with the findings of Reddy et al. [9] Dudhade et al. [10] Sharma et al. [11] and Kumawat et al. [12].

Table 1. Yield and economic outcomes of pigeonpea and intercropping systems across various treatments (aggregated over two years)

S.N.	Cropping system	Yield (q/ha)			Cost of cultivation (Rs./ha)	Gross return (Rs./ha)	Net return (Rs./ha)	BCR
		Main crop	Inter crop	Total productivity				
1.	Pigeonpea alone	17.34	-	17.34	49861	100248	50387	2.01
2.	Black gram alone	09.44	-	09.44	32966	51025	18059	1.54
3.	Sesame alone	06.32	-	06.32	31561	34056	2495	1.07
4.	Pigeonpea + black gram (1+1) a.s.	19.34	4.19	23.53	57876	134024	76148	2.32
5.	Pigeonpea + sesame (1+1) a.s.	16.61	3.69	20.30	56191	116053	54862	2.07

4. CONCLUSION AND RECOMMENDATION

Given that the intercropping of pigeonpea with black gram yielded a net return of Rs. 76,148 per hectare and pigeonpea with sesame produced a net profit of Rs. 54,862 per hectare, therefore, farm families residing in the vicinity of rainfed area may be advocated for adoption of intercropping of pigeonpea + black gram and pigeonpea + sesame for higher total productivity and profitability and harvest the fruits of newly generated technology.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Anonymous. Kharif keeSaghanPadhatiyan. Publication, Directorate of Agriculture, U.P. Lucknow. 2021;229.
2. Sekhon FS, Singh T, Singh S. Growth penology and yield of pigeonpea (*Cajanus cajan*) as affected by intercropping systems and application of nutrients level to intercrop. *Indian Journal of Agricultural Science*. 2018;88(3):509.
3. Kumar SNA, Gowda J, Venkate, Porama VR, Ramkrishna and Satish A. Qualifying and mapping of soil nutrients in Bantanahalli micro watershed using GIS and GPS. *Journal of Soil and Water Conservation*. 2017;16(1):18-24.
4. Piper CS. *Soil and Plant Analysis*. Hans Publication, Bombay; 1966.
5. Wakley AJ, Black IA. Estimation of organic carbon by Chromic acid titration method. *Soil Science*. 1934;37:29-38.
6. Muhr GR, Datta NP, Sankarasubramoney H, Diver RF, Leley VK, Donadhue RL. *Soil Testing in India*, U.S. Agency for International Development Mission to India, New Delhi; 1963.
7. Subhiah BV, Asija CL. A rapid procedure for estimation of available nitrogen in soil. *Current Science*. 1956;25:259-260.
8. Cochran WG, Cox GM. *Experimental Design*. Asia Publication House, New Delhi; 1963.
9. Reddy MM, Padmaja B, Rao LJ. Performance of pigeonpea cultivars in intercropping system under rainfed conditions in vertisols. *Indian Journal of Dryland Agricultural Research and Development*. 2007;22(2):208-212.
10. Dudhade DD, Deshmukh GP, Harer PN, Patil JV. Studies on intercropping of pulse crops with pigeonpea under rainfed condition. *Legume Research*. 2009;32(3): 215-217.
11. Sharma A, Rathod PS, Dharmraj PS, Mohan Charan. Response of pigeonpea to bio-fertilizers in pigeonpa based intercropping systems under rainfed conditions. *Karnataka Journal of Agricultural Sciences*. 2012;25(3):322-325.
12. Kumawat N, Singh RP, Kumar R, Kumari, Anupma and Kumar P. Response of intercropping and integrated nutrition on production potential and profitability on rainfed pigeonpea. *Journal of Agricultural Science (Toronto)*. 2013;4(7): 154-162.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/114014>