

Journal of Experimental Agriculture International

Volume 46, Issue 7, Page 881-893, 2024; Article no.JEAI.118375 ISSN: 2457-0591 (Past name: American Journal of Experimental Agriculture, Past ISSN: 2231-0606)

Minor Millets: Miracle Grain of South India

Sharmila Bharathi, C^{a++*} and A. Elango ^{b#}

 ^a ICAR – Krishi Vigyan Kendra, Tamil Nadu Veterinary and Animal Sciences University, Kallakurichi District, Tamil Nadu, India.
^b Veterinary College and Research Institute, Tamil Nadu Veterinary and Animal Sciences University, Salem District, Tamil Nadu, India.

Authors' contributions

This work was carried out in collaboration between both authors. Authors SBC and AE has initiated the idea of the article. Author CSB all sections of the article except sections on Minor millets as a source of fodder, minor millets-therapeutic potential and processing of minor millets. Author AE authored Minor millets as a source of fodder and processing of Minor millets. Both authors read and approved the final manuscript.

Article Information

DOI: https://doi.org/10.9734/jeai/2024/v46i72642

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: https://www.sdiarticle5.com/review-history/118375

> Received: 23/04/2024 Accepted: 28/06/2024 Published: 02/07/2024

Review Article

ABSTRACT

Minor millets are important traditional food crops for dry land farmers in India. India has the third largest area under minor millets cultivation in the world. Minor millets such as finger millet, foxtail millet, proso millet, kodo millet, little millet and barnyard millet are considered as a climate-smart crop. Since these crops are widely divergent for thermo and photoperiods and hence these are known for their climatic resilience and are relatively less prone to biotic stress factors. Added these crops are rich in protein, fat, crude fiber, iron and other minerals and vitamins with low glycemic index in comparison to cereals like wheat and rice. Hence they are labelled as nutri- cereals and act as the source of food, feed and fodder. Minor millets have curative and nutrient - rich properties

Cite as: C, Sharmila Bharathi, and A. Elango. 2024. "Minor Millets: Miracle Grain of South India". Journal of Experimental Agriculture International 46 (7):881-93. https://doi.org/10.9734/jeai/2024/v46i72642.

⁺⁺ Professor (Horticulture);

[#] Dean;

^{*}Corresponding author: E-mail: csbkvk79@gmail.com;

in the form of antioxidants which prevent weakening of human health such as lowering blood pressure, risk of heart disease, obesity, prevention of cancer, cardiovascular diseases and diabetes etc. Owing to its health benefits, the minor millets output has increased recently due to the availability of high-yielding varieties, pest and disease - resistant varieties, and improved cultivation techniques, rising by 1.05 percent annually. This review aims to document the production status, demand, varietal diversity, current consumption pattern, processing, nutrient composition and therapeutic potential of minor millets for further research on various aspects.

Keywords: Minor millets; nutrients; phytochemicals; antioxidant.

1. INTRODUCTION

Millets are a group of cereal type of food grain crops that are tiny - seeded, acclimatized to cultivation over a wide range of tropical and subtropical climatic conditions and can be grown with very low critical inputs. They are the sustenance food of the millions inhabiting the arid and semiarid tropics of the world. Millets are spread in most of the Asian and African countries and parts of Europe. According to FAO, the world production of millets is 89.17 million metric tons from an area of 74 million ha in 2020. "Sorghum. pearl millet, and maize are some of the major millets grown in huge quantities in India compared to minor or small millets, in spite of the fact that minor millets have higher nutritional value" [1-4]. "In India, minor millets, which consist of six species, are cultivated over around 2 million hectares, mainly in semi-arid, hilly, and mountainous regions" [5]. The six minor millet species grown are finger millet, little millet, italian or foxtail millet, barnvard millet, proso millet and kodo millet. Out of these, finger millet occupies about 60% of the area and contributes 70% to the overall minor millet production.

Minor millets are nutritious, climate - resilient, hardy and dry land crops, hence called as Nutricereals, contribute to a great extent to food and providing alimentary security. Recently, minor millets have gained the attention of the masses due to their non-gluten tendency. Nutrition wise, minor millets are rich in minerals, polyphenols, antioxidants, and fibers that are important for healthy body functioning. Minor millets are a habitual food source for the people of Manipur, Meghalava, and Nagaland, located in northeastern India. Although these grains were once referred to as coarse cereals, their nutrient richness has led the Government of India to officially categorize them as Nutri cereals.

2. ORIGIN AND DISTRIBUTION OF MINOR MILLETS

Millets were the first crops to be habituated by mankind in Asia and Africa which later distributed

across the globe as critical food sources to the evolving civilizations. Minor millets are small coarse grains belonging to the group of forage grass called millet [6] belongings to the family Poaceae; most of the genera belongs to the subfamily Panicoideae, which can grow in exorbitant ecological conditions [7]. "The word millet coined from the French word "Mille" which means "thousand", meaning that a handful of millet can hold up to a thousand grains" [8]. The minor millets include foxtail millet, little millet, kodo millet, proso millet and barnyard millet (Fig. 1). "There are four other types of minor millets namely brown top millet, fonio, tef, and job's tear millet (Coix laeryma), which are of minor importance in India because they are grown mostly in Africa" [9].

"Finger millet [Eleusine coracana (L.) Gaertn.] is an allotetraploid developed from its wild africana. progenitor. E. Domestication of cultivated finger millet started around 5,000 years ago in Western Uganda and Ethiopian highlands and the crop has spread to Western Ghats of India around 3.000 BC" [10]. "Foxtail millet [Setaria italica (L.) P. Beauv.] is a member of the subfamily Panicoideae and the tribe Paniceae. The green foxtail (S. viridis) is a wild progenitor of cultivated foxtail millet. Foxtail millet is grown since >10,500 years before in China" [11]. "Proso millet or common millet (Panicum miliaceum L.) is the true millet of history, and it is one of the oldest human foods and believed to the first cultivated cereal be grain. Recent studies suggested that Proso millet probably was domesticated in China and Europe" [12].

"Little millet (*Panicum sumatrense* Roth. ex. Roem. & Schult.) was habituated in India particularly in the Eastern Ghats of India, where it forms an important part of tribal agriculture" [13]. "In barnyard millet, two species namely *Echinochloa crus-galli* and *E. colona* are cultivated as cereals. *E. crus-galli* is native to temperate Eurasia and was spread in Japan some 4000 years ago, while *E. colona* is widely distributed in the tropics and subtropics of the Old World, and was domesticated in India" [14]. "Kodo millet belongs to the genus Paspalum, a multiple genus comprising about 400 species, most of which are native to the tropical and subtropical regions of the Americas, and the main center of origin and diversity of the genus is considered to be South American tropics and subtropics" [15]. "Kodo millet, also known as cow grass, rice grass, ditch millet, Native Paspalum, or Indian Crown Grass and it is estimated to have been domesticated in India 3000 years ago. It is cultivated by tribal people in limited areas throughout India, from Kerala and Tamil Nadu in the south, to Rajasthan, Uttar Pradesh and West Bengal in the North" [16].

"Teff (*Eragrostis tef*) is the most important minor millet in Ethiopia, especially in the poorly drained, heavy soils that predominate in the Central Plateau. However, the crop has not become important outside Ethiopia. Fonio (*Digitaria exilis*), also known as hungry rice, is grown as a minor millet crop throughout the savanna zone of West Africa. In parts of Guinea and Nigeria, it is the staple crop and is considered to be the oldest West African cereal and its cultivation is thought to date back to 5000 B.C" [17].




Fig. 1. Minor millet production status in south India - Area (A) in lakh ha, Production (P) in lakh tons and Yield (Y) in kg/ha



a. Finger millet



b. Proso millet



c. Foxtail millet



e. Barnyard millet



d. Kodo millet



f. Little millet

Fig. 2. Minor millets of south India

3. SCENARIO OF MINOR MILLETS IN INDIA

"India occupies the third largest area under minor millets cultivation in the world. Over the last eleven years, the area under minor millet has decreased considerably from 2011-12 to 2021-22. The area of minor millet shrunk from 7.99 lakh hectares (2011-2012) to 4.23 lakh hectares (2021-2022). Likewise, production under minor millet also reduced from 4.52 lakh tons to 3.75 lakh tons for the same period. However, the marginal gain in the productivity of minor millets from 565 q/ha to 885 q/ha was observed, but it was very minimal as compared to other crops like cereals, pulses, groundnut and cotton in India" [18]. "The declining trend in area and production was also reported in recently published studies" [19]. "The major reason for the reduction of area and production was the tradeoff between rice and wheat with minor millet" [20].

"In India, the major states involved in minor millet cultivation are Madhya Pradesh (32.4%,89000 hectares) followed by Chhattisgarh (19.5%, 52000 hectares), Uttarakhand (8.0%, 47000 hectares,), Maharashtra (7.8%), Gujarat (5.3%), and Tamil Nadu (3.9%) for the past seven years from 2015 – 2016 to 2021- 2022" [21,22].

In south India, the growth rate of area under minor millet cultivation (from 0.51 to 0.14 lakh ha in Andhra Pradesh, from 0.28 to 0.27 in Karnataka, from 0.31 to 0.23 lakh ha in Tamil Nadu) has declined over time (from 2015-16 to 2021-22) which was shown in Fig 2. In Karnataka, the increased trend of production was noticed during 2017-18 (0.26 lakh tons), then decreased during 2018-19 (0.17 lakh tons) and again increased (0.37) during 2019-20. The decreased production trend was observed in Andhra Pradesh for the past seven years (from 0.49 to 0.11 lakh tons), whereas in Tamil Nadu sustained growth of production and productivity was recorded during the same years.

Among the South Indian states, Karnataka state is well known for cultivating minor millets, with finger millet (Karnataka awarded with Geographical Indication (GI) tag for finger millet by the central government) serving as an important meal in the southern portion of the state [23] and its accounting 65% of area and production followed by Uttarakhand and Tamil But Tamil Nadu has the highest Nadu. productivity (3246 kg/ha) of finger millet followed by Karnataka (1726 kg/ha) which is above the national average yield (1697 kg/ha). The biggest small millets producing state is Karnataka (56% of total production) followed distantly by Tamil Nadu (14.0%) and Uttarakhand (9.3%). Rest of the states together contribute about one-fifth of the total production. Nearly 68% of area under small millets is occupied by finger millet, followed by little and kodo millets (about 10%) and rest by barnyard, foxtail and proso millets.

4. MINOR MILLETS - CLIMATE RESILIENT CROP

"Minor millets can be grown in India from June to November, and loamy, well-drained soil works best for growth. "These crops are widely differing for thermo and photoperiods and hence are known for their climatic resilience and are relatively less prone to biotic stress factors" [24]. "It needs very meagre inputs for its growth and development. Millets may be cultivated with very low rainfall since they are crops fed by the rain. Major millets require 450 mm of rainfall whereas minor millets only need 350 mm" [25,26]. Different minor millets can thrives well to the climatic conditions prevailing in a topographical location. For example, proso millet can finishes its life cycle in 60-80 days, while foxtail millet can mature in 70-90 days. Little millet and barnvard millet generally completes in 75-100 days, whereas, kodo millet and finger millets took slightly longer duration of 95-130 days. Because of earliness compared to other crops, these are highly acceptable for contingency crop planning and cropping system and have the unexpected feature of faster recovery and growth after stress alleviation, especially in finger millet and kodo millet.

5. VARIETAL WEALTH OF MINOR MILLETS IN INDIA

"Indian Institute of Millets Research and AICRP Project on Millets under Indian Council of Agricultural Research, has achieved in the development of more than 90 varieties for various agro- climatic regions so far. Till 2022, 302 varieties in different minor millets - 142 in finger millet, 41 in kodo, 38 in foxtail, 32 in little, 27 in proso millet and 22 in barnyard millet were developed and released for cultivation in different parts of the country" [27].

A number of varieties with high yield - potential have been released for different states. The list of improved and popular varieties recommended for different states of South India is given below (Table 1).

"Apart from the above varieties, nineteen different landraces from five millet crop species can be found in the Kolli Hills of Tamil Nadu. These include finger millet (Sundangi kelvaragu, Perungelvaragu, Karunguliyan kelvaagu, Arisi kelvaragu, Sattai kelvaragu, Kara kelvaragu), Italian millet (Perunthinai, Koranthinai, Senthinai, Mookanthinai. Palanthinai). little millet (Vellaperum samai. Katta vetti samai. Karumsamai, Malliasamai. Thirigulasamai, proso Sadansamai), common millet or (panivaragu) and kodo millet (varagu). The later two are cultivated only to a very limited extent" [28].

5.1 Bio Fortified Varieties in Minor Millets

Bio fortification in minor millets is a feasible strategy to control the micronutrient malnutrition among the rural poor. A few bio fortified cultivars with higher iron and zinc content along with high grain yield were released for farmers (Table 2) [29].

6. MINOR MILLETS AS SOURCE OF FODDER

"Minor millets are also used as fodder crops due to their higher biomass/green fodder yield. Kodo millet, little millet and proso millet are the most preferred animal feeds due to their higher palatability and crude protein content" [30]. "The minor millet grains can also become good alternatives for other cereal grains in the preparation of concentrate mixture of livestock and poultry, thereby reducing the cost of feed as well. Stover from millets is a highly recognized fodder, especially if cut and dried immediately after the heads have been harvested for grain" [31]. The nutritive value in terms of nitrogen and soluble nutrients is better in minor millet stover as compared to slender straws of paddy and wheat. Improved varieties and hybrids of the minor millets yield higher, both grain and stover (Table 3).

Minor millets	Karnataka	Tamil Nadu	Andhra Pradesh
Finger millet	GPU 28, GPU-45, GPU-48, PR 202, MR 1, MR 6, Indaf 7, ML-365, GPU 67, GPU 66, KMR 204, KMR 301, KMR 340, DHFM-78-3	GPU 28, CO 13, TNAU 946 (CO 14), CO 9, CO 12, CO 15	VR 847, PR 202, VR 708, VR 762, VR 900, VR 936, PPR-2700
Foxtail millet	SiA 3088, SiA 3156, SiA 3085, Lepakshi, SiA 326, Narasimharaya, Krishnadevaraya, PS-4	SiA 326, HMT 100-1 and PS 4, Narasimharaya, SiA 3088, SiA 3156, SiA 3085, DHFt-109-3, PS-4	TNAU 43, TNAU-186, TNAU 196, CO 1, CO 2, CO 4, CO 5, K2, K3, SiA 3088, SiA 3156, SiA 3085, PS-4
Kodo millet	GPUK 3, RBK 155, RK 390- 25, TNAU-86	KMV 20 (Bamban), CO 3, TNAU 86, GPUK 3, RK 390-25	-
Little mille	OLM 203, JK 8, BL-6, DHLM-36-3	Paiyur 2, TNAU 63 and CO 3,C0-4,K1, OLM - 203, OLM -20, BL-6, DHLM-36-3, DHLM-14-1	OLM 203, JK 8, BL-6, DHLM-36-3
Barnyard millet	VL 172, RAU 11, VL 181, DHBM-93-3, DHB-93-2	CO 1, CO 2, VL 181, VL 29, DHBM-93-3	-
Proso millet	GPUP 8, GPUP 21, TNAU 145, TNAU-151, TNAU-164, TNAU-202, TNPm-230, DHP-2769	Co-5, TNAU 151, TNAU 164, TNAU 145, TNAU 202, Co 4, K2, Co 3, Co 2, GPUP 21, GPUP 8 , TNPm-230	Sagar, Nagarjuna, Co 4, Co 3, TNAU- 151, TNAU-164, TNAU-202, TNPm- 230

Table 1. Improved and popular varieties of South India

Table 2. List of released bio fortified minor millet varieties in India

Crop	Bio fortified varieties	Characteristics	
Finger millet	VR 929 (Vegavathi)	Rich in iron (131.8 ppm). Grain yield:3110kg/ha	
Finger millet	ger millet CFMV1 (Indravati) Rich in calcium (428 mg/100g), iron (58.0 zinc (44.0 ppm).Grain yield: 2950kg/ha		
	CFMV 2 (Gira)	Rich in calcium (454 mg/100g), iron (39.0 ppm) and zinc (25.0 ppm). Grain yield:3200kg/ha	
Little millet	CLMV1(Jaicar Sama-1)	Rich in iron (59.0 ppm) and zinc (35.0 ppm).Grain yield:1580kg/ha	

Minor millets yields 3-8 tons of fodder depending on cultural as well as climatic conditions. Finger millet stover is considered a good feed in India and Nepal. Finger millet produces 2-3 t of dry fodder from a one hectare area under rain fed conditions, whereas under irrigated condition, the fodder yield reaches up to 6-10 t/ha. Green fodder from kodo millet is readily eaten by dairy animals and is highly digestible, when it will be harvested before flowerina. However. the stover obtained fed from the rain crop is relished more by livestock animals compared to an irrigated crop because stover of irrigated crop is

tough and fibrous and hence it becomes less palatable.

"Finger millet stover can be used to feed crossbred dairy cows along with concentrate feed. Such feeding method can increases the milk yield of 8 to 9 litres with improved body weight gain of 200–300 g/day. In addition to that it can also be used for growing heifers when supplemented with wheat bran (25%) or groundnut cake. Dry matter intake of stover is about 3.5 kg/day and the daily weight gain is 310–350 g. Further, it can also be made into silage with suitable supplements" [32].

Crop	Variety / Hybrid	Grain yield (q/ha)	Fodder yield (q/ha)
Finger millet	ML-365	50-55	65
	Vegavathi (VR 929)	36	77
	CFMV 1 (Indravathi)	31.1	84.4
	CFMV 2	29.5	86.1
	CFMV 3	32	87
Barnyard millet	Pratap Sanwa 1	15-17	50-55
Kodo millet	CKMV 1 (ATL 2)	28	70
Proso millet	GPUP 21	15.7	42
	Pratap Cheena	15-17	48-50
	TNAU 202	19	37
Foxtail millet	Pratap Kangni-1	16-18	46-50
	CO-7	18.6	51
Little millet	Sabara	12-21	52
	CO-4	15.7	58
	DHLM -14-1	16	61
	CLMV 1	16	55

Table 3. Stover yield of recent cultivars of minor millet in India

7. MINOR MILLETS: A POWER HOUSE OF NUTRIENTS

Minor millets are important due to their several nutritional and other useful characteristics. These crops are rich in protein, fat, crude fiber, iron and other minerals and vitamins in comparison to fine cereals like wheat and rice.

The protein content in minor millets like foxtail millet (11.2-15.0), little millet (10.0-15.0) and proso millet (10.0-13.0) is on par with wheat (11.8) and much higher than rice (6.8). Though the finger millet contains lesser protein (7.0-10.0), but it is rich in mineral nutrients and calcium in comparison to wheat and rice. All the millets contain more fiber than fine cereals. Particularly, the minor millets namely barnyard millet (9.5-14.0), kodo millet (5.0-9.0) little millet (4.0-8.0) and foxtail millet (4.5-7.0) are the richest source of fiber in comparison to wheat (1.0) and rice (1.0). The fat content of the minor millets varies from 1 to 7g, which is lowest in finger (1.3-1.8), proso (1-3.5) and kodo millet (1.4 -3.6) and highest in foxtail, and little millets (4 -7). Therefore, millets are now being tagged as "Miracle grains and nutria-cereals"

The minor millets are rich sources of Vitamin E and B-complex vitamins (except Vitamin B 12). Total niacin content present is 10.88 mg. Matured grains of millets have shown low levels of Vitamin C. "Finger millet contains over >10 fold higher calcium (240-410 mg per 100g), little and barnyard millets are rich in iron (13 to 20 mg per 100g) compared to other major cereals" [33].

7.1 Minor millet as Miracle grain -Nutritional composition and its uses

7.1.1 Finger millet

"The grains are an abundant source of natural calcium (240- 410 mg) which helps for bone strengthening and helps in reducing the risk of bone fractures. On daily consumption of whole grain of finger millet and its products can protect against the risk of cardiovascular diseases, Type II diabetes and gastro intestinal cancers and other health issues" [34]. "It is generally milled with the testa which is rich in dietary fiber and micronutrients to prepare flour" [35]. "The dietary fiber. minerals, phenolics and vitamins accumulated in the outer laver of the seed coat form the part of the food and offer their nutritional and health benefits" [36]. "In addition, it helps to increase the hemoglobin level and also helps to fight malnutrition and degenerative diseases" [37]. "Fibers of finger millet provides fullness feeling thus controlling the excessive food consumption" [12].

7.1.2 Barnyard millet

"Barnyard millet is a good source of protein, which is highly digestible and is an excellent source of dietary fiber with good amount of soluble and insoluble nutrients" [38]. "It is known for its low glycemic index, high phosphorous (280 mg) and magnesium content. Among minor millets, barnyard millet has the highest dietary fiber (9.5-14.0g) and niacin content" [39]. "The carbohydrate content of barnyard millet (55.0 – 65.0 g) is low and slowly digestible, which makes the barnyard millet a nature's gift for the modern mankind who is engaged in sedentary activities" [40]. Barnyard millet is most effective in reducing blood glucose and lipid levels.

7.1.3 Little millet

"Little millet is known for its magnesium (120-133mg), phosphorous (251-260mg), and protein content (10.0-15.0g). The unique feature of little millet is that it is rich in Poly Unsaturated Fatty Acids (PUFA) and flavonoids" [41]. "In addition to that, it is a rich source of sulphur-containing amino acids (cysteine and methionine) and lysine, which is lacking in most cereals" [42], "It is generally considered to induce a lower glycemic response due to the presence of abundant dietary fiber, resistant starch, and slowly digestible starch" [43]. It is wonderful millet which is suitable for people of all age groups. It helps to prevent constipation and heals all the problems related to stomach. It improves the semen counts of men and also helps for women with irregular periods problems. Its high fiber helps to reduce the fat depositions in the body. Little millet is rich in cholesterol, when consumed increases good cholesterol in the body, suitable for growing kids strengthens the body. lts complex and carbohydrate digests slowly which is very helpful for diabetic patients [44] and especially good for people who have low body mass [45].

7.1.4 Proso millet

"Proso millet has a high protein profile (10.0-13.0 g) following foxtail millet and also has a higher nutritional content when compared with major cereals as it contains a higher concentration of minerals and dietary fibre. Furthermore, it is a rich source of vitamins and minerals such as iron (4.0-5.2mg), calcium (20-23mg), potassium (250-320mg), phosphorus (230-281mg), zinc (1.4-2.4mg), magnesium (117-153mg), vitamin Bcomplex, niacin (4.5mg), and folic acid. Proso millet contains essential amino acids like lysine, isoleucine, and methionine leucine in comparatively higher quantities unlike other crops". [46]. "However, proso millet has an almost 51% higher essential amino acid index than wheat" [47]. "In grains, the quality of protein decreases after drying but the amount of protein increases" [48]. "Traditionally, it is used as recuperative food, especially post - pregnancy or illness" [39].

7.1.5 Kodo millet

Kodo millet is highly drought resistant crop among the millets. The grain of kodo millet is

coarsest of all food grains. It is a nutritious grain and a good alternative to rice and wheat. The grain is covered with a tough seed coat that must be removed before cooking.

"Kodo millet has the highest phosphorous content (215-310mg) and radical scavenging activity owing to its high phenol content. Consuming kodo millet reduces the risk of cardiovascular dysfunction" [49]. "The protein, fiber and mineral content present in this grain are much higher than the major cereals like rice. The major protein fraction in kodo millet is gluten" [50]. "It is a traditional food which helps to use in weight loss. It is easily digestible and is rich in phytochemicals and antioxidants which help in preventing different lifestyle related diseases. Kodo millet helps in reducing the joints and knee pain and helps in regularizing the menstruation in woman" [40].

"Apart from being a rich source of nutrients, kodo millet contains high amounts of polyphenols, tannins, phosphorus and phytic acids. The antioxidant activity of kodo millet decreases when the whole grain is dehulled and cooked" [51]. Kodo millet contains high amounts of vitamins and minerals, especially B-complex vitamins, B6, niacin and folic acid, Fe, Ca, Mg, K, and Zn. Kodo millet is very easy to digest and thus can be beneficial for infant and geriatric product formulation.

7.1.6 Foxtail millet

"Foxtail is also named as Italian millet and German millet. It is a good source of beta– carotene, which is the precursor of Vitamin and also it helps in slow and steady release of glucose without affecting the metabolism of the body. When people consume foxtail millet, the prevalence of diabetes is reduced and it is also known as healthy heart food due to its good source of magnesium (100-130mg)" [52].

Foxtail millet has a greater nutritional value when compared to major cereals such as wheat and rice due to its copious dietary fiber content, resistant starch, vitamins, minerals, and essential amino acids, except for lysine and methionine, but it is richer than most cereals. Foxtail millet contains highest protein (11.2-15.0g) among the minor millets. Added higher amount of stearic and linoleic acids present in foxtail millet helps in maintaining a good lipid profile.

7.2 Phytochemicals in Minor Millets

Phyto - chemicals are the chemical compounds that occur naturally in plants. Much of the health beneficial effects of minor millets have been to the presence attributed of abundant phytochemicals, including phenolic compounds. phytosterols etc. Phenolic compounds, viz. phenolic acids, flavonoids etc., which are present in millets impart bioactivities including antioxidant, anti-microbial, anti-diabetic, and antiproperties. hypertensive The 3-deoxy anthocyanin which is found in millets is known to exhibit several bioactivities including the ability to protect from certain types of cancers: they act as antioxidants and also play many roles in the body immune system. Lignans are a type of phytonutrient found in millet that is beneficial to the human body, which protest against hormonedependent malignancies like breast cancer and reduce the risk of heart disease.

7.3 Anti-nutrients: Challenges and Solutions

"Millets are nutritionally rich and even superior to major cereals in terms of energy value, proteins, fat and minerals. However, due to the presence of anti-nutrients like phytate, polyphenols, oxalates and tannins, mineral bioavailability is affected. These anti-nutrients form complexes with dietary minerals, such as calcium, zinc and iron leading to a marked reduction in its bioavailability and make them biologically unavailable to human beings" [53]. However, the negative impact of these anti nutrients can be taken care by using common household food processing techniques like decortications, milling, soaking, malting, germination, fermentation, popping and cooking etc. These methods reduce the content of phytates, phenol, tannins and trypsin inhibitor activity, improve the digestibility of millets apart from enhancing bio-availability of minerals.

8. SHELF LIFE OF MINOR MILLETS

The shelf- life of minor millet grain is normally above 6–8 months for fair and average quality grain, with 10–12% moisture. The shelf-life deterioration of processed products especially flour is a big challenge. Because minor millets are small grains, the separation of the oil rich germ from the endosperm might pose issues when millet flour is stored. Microorganisms also play a vital role in determining the shelf life of the product.

8.1 Processing of Minor Millets

The harvesting process followed for minor millets is mostly carried in a traditional manner wherein a lot of refractions such as immature grains, chaffs, mud particles, stones, admixed grains as well as obnoxious material, dust, etc. will be mixed. For this, de-stoners, graders and aspirator systems suitable for millets are available and millet processors are using them effectively. Minor millets have good grain qualities suitable for processing. Being a staple food and consumed at household levels, processing must be considered at both traditional as well as industrial levels, involving small, medium and large-scale entrepreneurs.

8.2 Primary Processing

The primary processing (wetting, dehulling and milling) of minor millets is an important step in converting the grain into edible form and thereby enhancing their quality as well as consumer acceptability. Among the minor millets, finger millet is naked grains as almost all the glumes get detached from grains during harvesting.

On the other hand, the processing of little, proso, kodo, barnyard, browntop and foxtail millets is complicated as they have an inedible husk that needs to be removed, followed by the debranning to a desirable extent through primary processing. It was noticed that barnyard, little, brown top and kodo millets needs multiple passages whereas foxtail and proso could be dehusked in a single stage.

8.3 Secondary Processing

"Secondary processing such as fermentation, malting, roasting, extrusion, flaking, popping etc., and product development involves the conversion of the primary processed raw material into different Ready to-Eat and Ready-to-Cook minor millet products. Although the dehulled and de-branned minor millets are largely used for cooking and consumption similar to milled rice, they are pulverized into flour and suji and for use as roti and other foods similar to rice/wheat flour and semolina" [54].

During processing, some nutrient losses will also be occur. To overcome this problem, fortified products such as cookies, vermicelli, pasta, khichadi mix and bread etc., are also developed by adding natural nutrient-rich ingredients like garden cress, spinach (rich in iron), gingelly seed (rich in zinc) etc., to enhance us enhance the iron and zinc proportions in to them. Various processing methods such as germination or thermal processing, malting, soaking and fermentation can minimize the nutritional and increase the loss also well chemical physiological as as accessibility of micronutrients in the body and also decrease the anti-nutrients like phytates.

9. ECONOMICS AND MARKET TREND

"The cultivation of millets is highly beneficial to small-scale farmers as they are relatively easy to grow and require low inputs, which results in lower costs and higher profitability. The total cost of cultivation of foxtail millet worked out to be Rs.20,055/- per hectare. Farmers got about 12.05 quintals of main product and 4.22 tonnes of byproduct. The gross return from main product worked out to Rs.2619/- per quintal. Thus, gross returns realized from both main product and byproduct were Rs.32171/- per hectare. The total cost of cultivation of little millet worked out to be Rs.22278/per hectare. Farmers got about 10.65 guintals of main product and 2.44 quintals of byproduct. Thus, gross returns realized from both main product and byproduct were Rs.33784/- per hectare. The total cost of cultivation of finger millet worked out to be Rs.38128/- per hectare. Farmers could receive 14.05 guintals of main product and 7.50 guintals of byproduct. In turn, gross returns obtained from both main product and byproduct were Rs.51603.50/- per hectare, respectively. The net return realized from millet cultivation after deductina cost of cultivation came to foxtail millet was Rs.12116/- per hectare, little Rs.11506/per hectare, millet was kodo millet was Rs.5391/- per hectare, barnyard millet was Rs.9728/- per hectare, brown top millet was Rs.7352/- per hectare and in case of finger millet was Rs.13475.50/- per hectare" [55].

Minor millets is grown mostly as food crop. Farmers sell millets as a grain at the local markets to both traders as well as consumer. Now Consumers are aware that, minor millets are significant to their diet and recent fast food habits. "Further procurement of millets through government agency at MSP, establishment of advanced processing units, storage units and training on value added products of millets etc. will enhance the economic status of farmers" [56].

10. CONCLUSION

Over the last few years, there has been increasing recognition of their nutrient composition and benefits as healthy food. In addition, minor millets output has increased recently due to the availability of high-yielding varieties, pest and disease - resistant varieties, and improved cultivation techniques, rising by percent annually. Therefore, dietary 1.05 modification by increasing the consumption of a wide variety of fruits, vegetables and minor millet grains daily is a practical strategy for consumers to optimize their health and reduce the risk of chronic diseases. Because of their health benefits, these grains do need a great promotion to reach the heights of the major cereals in terms of their production and utilization.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declared that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

ACKNOWLEDGEMENTS

The authors wish to thank the anonymous reviewer(s) for their suggestions

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Tiwari H, Naresh RK, Kumar L, Kataria SK, Tewari S, Saini A, Yadav RK, Asati R. Millets for food and nutritional security for small and marginal farmers of North West India in the context of climate change: A review. Int. J. Plant Soil Sci. 2022 Dec. 9 [cited 2024 Jun. 4];34(23):1694-705. Available:https://journalijpss.com/index.php /JJPSS/article/view/2594
- Iqbal S, Bhat MA, Sheikh TA, Baba ZA. Deciphering the importance of underutilized millet crop towards sustainable feed and food sources. Curr. J. Appl. Sci. Technol. 2023 Dec. 30 [cited 2024 Jun. 4];42(48):165-7. Available:https://journalcjast.com/index.ph p/CJAST/article/view/4351

- Balasubramanian S, Viswanathan R. Influence of moisture content on physical properties of minor millets. Journal of food science and technology. 2010 Jun;47:279-84.
- Kaur B, Singh A, Suri S, Usman M, Dutta D. Minor millets: A review on nutritional composition, starch extraction/modification, product formulation, and health benefits. Journal of the Science of Food and Agriculture. 2023 Aug 15;103(10):4742-54.
- Prashanthi A, Geetha R. Millet status in India - production and consumption. Just Agriculture multidisciplinary e – Newsletter. 2023;3(5):244-250.
- Dayakar R, Gill B, Thakur MS, Manjula S, Ghoora D, Kowsalya S, Pant KK, Tonapi VA. Millet international recipes: A Culinary Journey of Tradition and Innovation; 2022.
- 7. Bhag M, Padulosi S and Bala Ravi S. A book on Minor Millets in South Asia; 2010.
- Gowri UM, Shivakumar KM. Millet scenario in India. Economic Affairs. 2020;65(3):360-370.
- 9. ASSOCHAM. Millets status way forward; 2021.
- 10. Indian Institute of Millet Research. Nutritional and Health Benefits of Nutri Cereals; 2018.
- Anbukkani P, Balaji SJ and Nithyashree ML. Production and consumption of minor millets in India - A structural break analysis. Ann. Agric. Res. New Series. 2017;38(4):1-8.
- Sangappa DR, Srinivasa Babu K. A study on area-production-productivity of minor millets in India. Biological Forum – An International Journal. 2023;15(1):275-280.
- 13. Ashoka P, Gangaiah B, Sunitha NH. Millets -foods of twenty-first century. International Journal of Current Microbiology and Applied Sciences. 2020;9(12):2404- 2410.
- 14. Weber S. Out of Africa: The Initial Impact of Millets in South Asia. Current Anthropology. 1998;39(2):267- 274.
- 15. Seetharam A, Riley K, Harinaryana G. Small millets in global agriculture. Oxford & IBH, New Delhi. 1989;9-12.
- Baljeet K, Ajay S, Shweta S, Mohammad U, Debashish D. Minor millets: A review on nutritional composition, starch extraction/modification, product formulation, and health benefits. J. of the Science of Food and Agriculture. 2023. DOI: 10.1002/jsfa.12493

- Hilu KW, de Wet JMJ, Harlan JR. Archeobotany and the origin of finger millet. Am. J. Bot. 1979;66:330-333.
- Yang X, Wan Z, Perry L, Lu H, Wang Q, Zhao C, Li J, Xie F, Yu J, Cui T, Wang T, Li M, Ge Q. Early millet use in northern China. Proc. Natl. Acad. Sci. 2012;109:3726–3730.
- Harlan JR. Crops and man. American Society of Agronomy and Crop Science. 1975;295.
- 20. De wet JMJ, Prasad R, Brink D. Systematics and domestication of *Panicum sumatrense* (Gramineae). Journal d'agriculture traditionnelle et de botanique appliquée. 1983,30:159-168.
- 21. De wet JMJ, Prasada Rao, Mengesha MH, Brink De. Domestication of *Sawa millet* (*Echinochloa colona*). Econ. Bot. 1983;37:283-291.
- 22. Chase A. The North American species of paspalum. Contributions from the United States National Herbarium. 1929;28:1-310.
- 23. De wet JMJ, Prasada R, Mengesha MH, Brink D. Diversity in Kodo milet (*Paspalum scrobiculatum*). Econ. Bot. 1983;37:159-163.
- Jones MK. Between fertile crescents: Minor grain crops and agricultural origins In: MK Jones (ed) Traces of Ancestry: Studies in Honour of Colin Renfrew, Oxbow Books, Cambridge. 2004;127-135.
- 25. Goron TL, Raizada M. Genetic diversity and genomic resources available for the small millet crops to accelerate a New Green Revolution. Front. Plant Sci. 2015;6:157-158.
- 26. Sukanya T, Prabhakar S, Krishne Gowda KT, Swarna R, Hariprasanna K, Tonapi VA. Good agronomic practices for higher yield in small millets, ICAR - All India Coordinated Research Project on Small Millets, ICAR-Indian Institute of Millets Research, Hyderabad; 2022.
- 27. Upadhyaya HD, Vetriventhan M Dwivedi SL, Pattanashetti SK, Singh SK. Proso, barnyard, little and kodo millets. In: Mohar Singh and HD Upadhyaya (eds) Genetic and Genomic Resources for Grain Cereals Improvement. 2015;321-343.
- 28. Hariprasanna K. High yielding varieties for enhancing the production of small millets in India. Indian farming. 2023;73(1):42-46.
- 29. Hiroyuki T, Latha N. Minor millets in Tamil Nadu, India: Local market participation, onfarm diversity and farmer welfare; 2015.

Available:https: // www.researchgate.net /publication /259425992

- Raju RC, Manjeet S, Monu K, Harsh C, Mukesh P, Babulal D. Biofortification in India: Present status and future prospects. The Pharma Innovation Journal. 2022;SP-11(1):782-786.
- Vetriventhan M, Azevedo VC, Upadhyaya HD, Nirmalakumari A, Kane PJ, Anitha S. Genetic and genomic resources and breeding for accelerating improvement of small millets: Current status and future interventions. Nucleus. 2020;63(3):217– 239.
- Seetharam AK, Riley W, Harinarayana G. Small millets in global agriculture proceedings of the first international small millets workshop Bangalore, India; 1986.
- Venkatesh B, Avinash S, Balakrishna D, Sooganna. Millet crops as source of fodder. Indian Farming. 2023;73(01):19-21.
- Saleh AS, Zhang Q, Chen J, Shen Q. Millet grains: Nutritional quality, processing, and potential health benefits. Comprehensive reviews in food science and food safety. 2013;12(3):281-295.
- 35. McKeown NM, Meigs JB, Liu S, Wilson PW, Jacques PF. Whole- grain intake is favorably associated with metabolic risk factors for type 2 diabetes and cardiovascular disease in the Framingham Offspring Study, A. J. of Clinical Nutrition. 2002;76(2):390–398.
- Devi PB, Vijayabharathi RS, Sathyabama NG, Malleshi VB and Priyadarisini. Health benefits of finger millet (*Eleusine coracana* L.) polyphenols and dietary fiber: a review. J. of Food Science Technology. 2014;10(10):26-30.
- Antony U, Moses LG and Chandra TS. Inhibition of Salmonella typhimurium and Escherichia coli by fermented flour of finger millet (*Eleusine coracana*), World. J of Microbiology and Biotechnology. 1998;14(6):883–886.
- 38. Reddy OSK. Smart millet and human health; 2017.
- Hadimani NA, Malleshi NG. Studies on milling, physic- chemical properties, nutrient composition and dietary fiber content of millets. J. of Food Science and Technology. 1993;30: 17-20.
- 40. Pramitha L, Subhalakshmi J, Jeeva G, Surendar A. Small millets exploring the hidden potential; 2021.
- 41. Veena B, Chimmad BV, Naik RK, Shanthakumar G. Physico chemical and

nutritional studies in barnyard millet. Karnataka J. of Agricultural Sciences. 2005;18:101-105.

- 42. Indirani K, Devasena M. Review on nutritional profiles and health benefits of little millets –India. Int. J. Res. Eng. Sci. 2021;9(7):21-29.
- 43. Deshpande SS, Mohapatra D, Tripathi MK, Sadvatha RH. Kodo millet-nutritional value and utilization in Indian foods, Journal of Grain Processing and Storage. 2015;2:16-23.
- 44. Chethan S, Malleshi NG. Finger millet polyphenols: Optimization of extraction and the effect of pH on their stability. Food chemistry. 2007;105(2):862-870.
- 45. Gayathri A. Little millet nutrition, health benefits and facts you must know; 2015.
- 46. Nutritive value of Indian Foods- Millets in your meals, NIN; 2007.
- Kalinova J, Moudry J. Content and quality of protein in Proso millet (*Panicum miliaceum* L.) varieties. Plant Foods and Human Nutrition. 2006;61:45–9.
- 48. Jana kalinova. Nutritionally important components of Proso millets (*Panicum miliceum* L.) food. 2007;1(1):91-100.
- 49. Deepak Taggelli RG, Thakur V. Minor millets-their potential health benefits and medicinal properties: A review. Int. J. Pure Appl. Sci. 2018;6(1):1677.
- 50. Sudharshana L, Monteiro PV, Ramachandra G. Studies on the proteins of kodomillet (*Paspalum scrobiculatum*). J. Science of Food and Agriculture. 1988;42(4):315-323.
- 51. Chandrasekara A, Naczk M, Shahidi, F. Effect of processing on the antioxidant activity of millet grains. Food Chemistry. 2012;133(1):1-9.
- Murugan R, Nirmalkumari A. Genetic divergence in foxtail millet (Setaria italica (L.) Beauv). Indian Journal of Genetics. 2006;66(4):339-340.
- 53. Malik M, Singh U, Dahiya S. Nutrient composition of pearl millet as influenced by genotypes and cooking methods. Journal of Food Science and Technology. 2002;39(5):463-468.
- 54. Arora P, Sehgal S, Kawatra A. Content and HCI extractability of minerals as affected by acid treatment of pearl millet. Food Chemistry. 2003;80(1): 141–144.
- 55. Arjuman B, Ganapathy MS, Govinda V Siddayya G, Girish MR, Shamshad B, Mohan Kumar

TL. Value chain analysis of minor millets for improving economic status of farmers in Tumakuru District of Karnataka. Mysore J. Agric. Sci. 2022;56 (4):51-60.

56. Uma Gowri M, Prabhu R. Millet production and its scope for revival in India with special reference to Tamil Nadu. International Journal of Farm Sciences. 2017;7(2):88-93.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history: The peer review history for this paper can be accessed here: https://www.sdiarticle5.com/review-history/118375