



Soil Fertilization and Texture on Boron Accumulation and Wood Volume in *Corymbia citriodora* (Hook) K.D. Hill & L.A.S Johnson

**Antônio Leis Pinheiro¹, Karine Fernandes Caiafa¹, Daniel Teixeira Pinheiro^{2*},
Tássia Fernanda Santos Neri Soares², Rodrigo Vieira Leite¹,
Patrícia Ramalho de Barros³ and Matheus Ferreira França Teixeira²**

¹*Departamento de Engenharia Florestal, Universidade Federal de Viçosa, Av. PH. Rolfs s/n, 36570000, Viçosa, MG, Brazil.*

²*Departamento de Fitotecnia, Universidade Federal de Viçosa, Av. PH. Rolfs s/n, 36570000, Viçosa, MG, Brazil.*

³*Departamento de Solos e Nutrição de Plantas, Universidade Federal de Viçosa, Av. PH. Rolfs s/n, 36570000, Viçosa, MG, Brazil.*

Authors' contributions

This work was carried out in collaboration among all authors. Author ALP designed the study, performed the statistical analysis, wrote the protocol and wrote the first draft of the manuscript. Authors KFC, DTP and PRB worked in the final draft of the manuscript. Authors TFSNS, RVL and MFFT managed the literature review and worked translating Portuguese to English. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JEAI/2019/v37i130255

Editor(s):

(1) Dr. Peter A. Roussos, Assistant Professor, Lab. Pomology, Agricultural University of Athens, Greece.

(2) Dr. Lixiang Cao, Professor, Department of Biotechnology, Sun Yat-sen University, China.

Reviewers:

(1) Dale Loussaert, USA.

(2) Lucas Aparecido Manzani Lisboa, São Paulo State University (Unesp), College of Agricultural and Technological Sciences, Dracena, São Paulo, Brazil.

(3) Dr. Alok Nahata, Ying Zhi Agricultural and Industries Sdn Bhd, Malaysia.
Complete Peer review History: <http://www.sdiarticle3.com/review-history/48777>

Received 12 March 2019

Accepted 28 May 2019

Published 03 June 2019

Original Research Article

ABSTRACT

This study aimed to evaluate the effects of soil fertilization and texture on leaf Boron (B) accumulation and its relation with wood volume of *Corymbia citriodora* Hill & Johnson. The experiment was set in randomized block with four replications, four B fertilization levels (0, 1.1, 2.2

*Corresponding author: E-mail: pinheiroagroufv@gmail.com;

and 4.4 g.plant⁻¹) in two soils types (sandy and clayey). To determine leaf B content, 25 leaves were collected from the median portion of four trees for each treatment. Leaves were dried on oven and B content was determined by the Azomethine-H method using extract obtained by dry digestion. The diameter at breast height and the total height of 25 trees were collected in all treatments and wood volume was calculated. Data were submitted to analysis of variance and the means adjusted to regression equations. The regression coefficients were evaluated by t-test at 1 and 5% probability. It was verified that clay soil produces more wood, compared to sandy soil. There was a gain increase in foliar B as B doses increased in both soil types. Leaf B affected *Corymbia citriodora* productivity only in clay soil.

Keywords: *Eucalyptus*; nutrition; fertility; productivity.

1. INTRODUCTION

The increase in wood production, especially after the evolution of clonal propagation and improvements on genetic techniques, has led to clones that are more demanding in a nutritional view. Micronutrient deficiency symptoms have been more commonly observed, mainly boron (B), which, according to Malavolta and Kliemann [1], is the micronutrient that presents more limiting levels in forest soils. B deficiency occurs frequently in eucalyptus and has been related to many other factors such as nutrients accumulation, soil type and others [2,3,4].

The genera *Eucalyptus* and *Corymbia* are the main forest essences cultivated in Brazil today. Among the species, *Corymbia citriodora* KD Hill & LAS Johnson stands out for having easy adaptation, rapid growth, good quality wood and uses of essential oils production [5]. The planting of this species has grown greatly, especially in areas with water and nutritional restrictions, as occurs in the Northern region of Minas Gerais, Brazil. In these areas, the use of drought-tolerant genetic materials, associated with B fertilization, has been shown to be effective in mitigating water deficit effects [6].

Among the main B functions are cell wall formation and components synthesis, such as pectin, cellulose and lignin [7]. Thus, species with a cell wall rich in pectin, such as dicotyledons, usually have high B requirements, compared to other species [8].

Barreto et al. [9] found that clones of *Eucalyptus grandis* x *Eucalyptus urophylla* responded to B fertilization in both shoot growth and biomass production, with gains in height and biomass, 240 days after transplanting in pots, 35-54%, and 21-64%, respectively. Silva et al. [10] reported that B foliar application improved nutrition of this

nutrient levels in eucalyptus clones leaves, even under water stress conditions.

The B absorption occurs mainly by mass flow, which is directly affected by soil texture and its water retention capacity [11]. Mattiello et al. [12] studying B transport in the soil and its absorption by eucalyptus, verified that maximum production of dry shoot matter was obtained in the B doses corresponding to 0.96 and 1.82 mg.dm⁻³ in -10 and -40 kPa water potentials, respectively.

Nutritional factors have been important for reducing production of planted forests in Brazil since they limit plant growth and consequently reduce productivity [12]. Therefore, researches that seek to maximize forest productivity through more adequate fertilization have fundamental importance.

Considering these factors, this study aims to evaluate the effects of soil fertilization and texture on B foliar accumulation and its relation with wood volume in *Corymbia citriodora*, considering the importance of this species for the economy and the high cost-benefit ratio for implementing adequate nutrition programs in forest species.

2. MATERIALS AND METHODS

2.1 Location and Soil Characterization

The experiment was set in João Pinheiro (sandy soil) and Três Marias (clayey soil) cities, located in Minas Gerais, Brazil. The climate is classified as dry-subhumid tropical. João Pinheiro city has an average annual temperature of 22.6°C and rainfall of 1406 mm. Três Marias has an average annual temperature of 23.5°C and rainfall of 1214 mm. Both cities have long dry periods, mainly between May and September. The experimental area João Pinheiro city is located at 17°00' south latitude; 45° 50' west longitude and

altitude of 500-550 m. The experimental area in Três Marias city is located at 18°08' south latitude; and 45°12' west longitude and 700 m altitude.

The main physical and chemical soils characteristics of experimental areas are described in Table 1.

2.2 Plant Cultivation and Treatments

The *C. citriodora* seedlings were planted at 3x2 m spacing, with a useful area per plant of 6 m² and a population density of 1,666 plants/ha. Before the planting, liming (Ca:Mg ratio of 4:1) was applied to soil aiming the percent base saturation of 80%. At the planting, fertilization (N, P,K) was applied in grooves and performed according to the soil analyze and type [13]. The fertilization of K and N were divided into two stages, at planting and 6 months later. For both experimental areas, three months after planting, B fertilization was carried out with the application of 0; 1.1; 2.2; and 4.4 g.plant⁻¹ in the soil. At 15 months, additional fertilization (levels 1, 2 and 4) with 3.6 g.plant⁻¹ of borax (Na₂B₄O₇.10H₂O) in both soil types. The management consisted of normal cultural practices with periodic weeding during the entire experiment.

2.3 Evaluations

All evaluations were performed with plants at seven years. The diameter at 1.30 m height (DBH) of 25 plants in all treatments was measured with a tape measure. The trees' heights were obtained using a hypsometer and the volume in m³/ha of both species was also calculated.

To determine leaf B content, 25 leaves removal from the median portion of four trees for each treatment were collected. After that, they were dried in a forced circulation air oven at 70°C until weight stabilized. The samples were ground and the B content was obtained by the Azomethine-H method, using extract obtained by dry digestion [14].

2.4 Experimental Design and Statistical Analysis

The experiment was arranged in a randomized complete block design (RCB) in a subdivided plots scheme, with 4 fertilization levels, 2 types of soils, 1 eucalyptus species and 4 replications. The analysis of variance joint was performed according to Banzatto and Kronka [15]. Data of

wood diameter and leaf B were submitted to analysis of variance and regression. The regression coefficients were evaluated by the t-test at 1 and 5% probability.

3. RESULTS AND DISCUSSION

The analysis of variance showed no significant interaction between soil and boron for wood volume. However, for leaf Boron, this interaction was significant. These results show dependence or independence of these factors for different analysis (Table 2).

The wood production of *C. citriodora* after different B doses application had a higher efficiency in the clayey soil, compared to sandy soil by the t-test, at 5% probability (Table 3). The clayey soil has great natural fertility (Ca²⁺, Mg²⁺, K⁺ and B_{daq}) and its granulometry associated with a higher organic matter content (Table 1) allows a better water retention, which may explain its higher performance when compared to sandy soil.

The studied area had a low rainfall distribution at the beginning of the year, with an intense water deficit between May and August. Thus, a greater water capacity retention had fundamental importance to guarantee gains in productivity. Ramos et al. [16] describes soil moisture as an important parameter to B uses in the soil by *C. citriodora*. It's also important to emphasize that water deficit interferes in B uptake in eucalyptus, causing shoot dieback, especially in young plants. In this context, Dias et al. [17] observed a relevant occurrence of this disease in treatments without addition of B and in the more restrictive water regime. In addition, according to these authors, the fertilization with 0.55 g boron plant⁻¹ in soil was sufficient to avoid shoot dieback.

Organic matter is the biggest source of B in tropical soils [18]. As shown in Table 3, the sandy soil has organic matter content almost three times lower, compared to the clayey soil. The same was observed to B content, thus, there is a correlation between organic matter and B content in the soil.

In general, wood volume (m³.ha⁻¹) increased as B doses increased. According to the regression equation, the B dose required to reach the highest wood volume (161.01 m³.ha⁻¹) would be 6.49 g.plant⁻¹ (Fig. 1).

The results show that the fertilization using B is fundamental to reach higher yields in the forest plantations. B plays a key role in stem growth

and is directly related to volume increment, mainly acting on cell walls formation in the wood [18,19]. Similar results were obtained by Sgarbi et al. [20,21], evaluating nutritional status and soil fertility in *Eucalyptus grandis* plantations in two different areas. In these studies, it was observed that B is one of the nutrients that most limit this species growth in these studied areas. Furthermore, a positive relation of this element and *Eucalyptus* productivity was observed.

It is observed that in the very clayey soil, the maximum level of leaf B is 108.37 mg.kg⁻¹, which by the curve would be equivalent to an application of 3.35 g.plant⁻¹ of B (Fig. 2). In the soil with a sandy-loam texture, the maximum point of leaf B is 256.31 mg.kg⁻¹, which is reached when 8.05 g of boron.plant⁻¹ is applied.

In environments with low water availability, B is the micronutrient that more affects tree growth. Moreover, this micronutrient is associated with a series of metabolic reactions such as sugar transport, cell wall synthesis, respiration and others. Thus, the B deficiency may lead to

disorders in these processes, further aggravating problems related to water deficit [22].

Tirloni et al. [23] observed that fertilization using B influenced *C. citriodora* growth, evaluating height and diameter only when was performed at the beginning of the rainy season and when plants were already at 29 months old. In younger plants, there was no increase in height or diameter with B application, independent the dry or rainy seasons. According to Paula [24], no differences related to eucalyptus wood volume was verified when plants were fertilized with different B doses and evaluated at 20 and 24 months old. In the present work, the observed influence of B can be related a many factors, such as genotype and climate.

In the sandy soil, it is expected that B availability, absorption and translocation are lower mainly because its absorption by mass flow and due to lower water retention capacity [25,26]. However, in this study was verified that the leaf B concentration was higher in grown trees in sandy soil than in clayey soil (Fig. 2).

Table 1. Chemical soil characteristics for each experimental area

Area	pH ⁽¹⁾ (H ₂ O)	Al ³⁺ (²)	Ca ²⁺ (³)	Mg ²⁺ (³)	P(³)	K(³)	B _{dag}	OM ⁽⁵⁾	Sand	Silt (%)	Clay
Sandy	4.48	0.8	0.43	0.1	3	58	0.3	0.85	79	4	17
Clayley	4.7	1.27	0.55	0.39	3	109	0.6	2.74	11	20	69

⁽¹⁾ Ratio 1:2.5; ⁽²⁾ KCl Extractor 1 mol.L⁻¹; ⁽³⁾ Mehlich⁻¹ Extractor; ⁽⁴⁾ Hot water Extractor (65 °C) e ⁽⁵⁾ Organic Matter: Walkley & Black Method

Table 2. Analysis of variance of wood volume and leaf Boron by *C. citriodora* under crescent boron doses (g/plant) on two soils with different textures

Source of variation	df	Mean square	
		Wood volume	Leaf boron
Soil	1	15044.81 **	75763.23**
Boron	3	2818.86 **	66371.17**
Block	3	785.92**	148.93**
S x B	3	231.33 ^{ns}	1850.84**
Error	31	514.05	136.11
CV (%)		14.64	11.70

Note. ns, **: non significant and significant by F test (P < 0.01)

Table 3. Wood volume means (m³.ha⁻¹) produced by *C. citriodora* under crescent boron doses (g/plant) on two soils with different textures

Boron (mg.plant ⁻¹)	Clayey (Três Marias)	Sandy (João Pinheiro)	Means
0	133.69	134.19	133.94
1.1	154.36	126.86	140.61
2.2	157.7	142.36	150.03
4.4	165.53	150.53	158.03
Means	152.86 a	138.53 b	-

Note. a > b by test F (P < 0,05)




Fig. 1. Volume of wood produced by *C. citriodora* in response to application of different B doses in the soil (g.plant⁻¹)
ns: non significant, bars: standard deviation




Fig. 2. Leaf boron level of *C. citriodora* under different boron fertilizing doses on two different soil types
*** : significant (P < 0.01), Bars: standard deviation*




Fig. 3. Relation between wood volume increment (m³.ha⁻¹) and leaf boron (mg.kg⁻¹) of *C. citriodora* cultivated in two different soil types

The clayey soil has a greater water retention capacity, which would facilitate the absorption of B. However, due to a greater presence of iron and aluminum oxides, the availability of these nutrients is compromised once adsorption of these elements occurs. According to Barros and Novais [27], B occurs in the soil in the form of boric acid or borate and can be adsorbed to organic and inorganic fractions. Ferreira [28] describes that different minerals types of the soil's clay fraction play a fundamental role in controlling the available B content. The adsorption of B is mainly due to iron and aluminum oxides, since these oxides influence more than any other silicate clay type.

Tirioni et al. [23] reports that in very weathered soils, such as those observed in the cerrado, eucalyptus usually presents symptoms of B deficiency. These symptoms appear mainly when the rainfall regime is characterized by prolonged periods of water deficit [17]. Mattiello et al. [12] concluded that the water deficit elevates the external requirements of B and promotes a higher nutrient concentration in the plant tissue.

In sandy soils B may be leached during the rainy season, on the other hand, in clayey soils, B may be adsorbed in the organic and inorganic fraction, being the soil type directly related to nutrient deficiency occurrence [29]. Ramos et al. [16] observed that B distribution in plant tissues of *C. citriodora* was directly influenced by the amount of nutrient available, in other words, by nutritional status. Ferreto et al. [3] evaluated the relation between B and liming in Eucalyptus cultivated in sandy soils and observed that limed soils showed higher B availability at the same time that plants showed a higher B concentration in shoots and roots.

It is important to note that other important factors must be considered. In general, B translocation and redistribution in plant tissues are related to its low mobility in the phloem and are also strongly influenced by the plant transpiration flow, which depends not only on soil texture, but also on relative humidity, temperature and light intensity [30,31,32]. Being a dry sub-humid tropical climate in the experimental areas, the plants underwent different dry cycles, which may have influenced the translocation of B.

The leaf B content and the wood volume increment ($m^3 \cdot ha^{-1}$) of *C. citriodora* for both soil types are shown in Fig. 3.

For the sandy-loam soil, there was no significant difference in wood volume increment ($m^3 \cdot ha^{-1}$) as increasing leaf B dose, suggesting that in this soil the B absorbed was not directed to physiological processes related to the increase of the wood volume, although there is great absorption of this nutrient. On the other side, in clayey soil, there was an increase in wood volume as increasing B absorption. Biomass production and nutrient content present in the eucalyptus are positively related to each other. In addition, biomass production depends on water availability and the plant's ability to absorb, distribute and use this nutrient [30,31,32,33].

4. CONCLUSION

The clayey soil has been shown to be more efficient in the production of wood than the sandy soil because it has a higher natural fertility and a higher content of organic matter, which allows a greater retention of water. Under the studied conditions, $4.4 \text{ g} \cdot \text{plant}^{-1}$ of B allows higher leaf accumulation and volume of wood in *Corymbia citriodora*.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Malavolta E, Kliemann HS. Desordens nutricionais no Cerrado. 1985;136. Portuguese.
2. Silveira RLVA, Moreira A, Takashi EN, Sgarbi F, Branco EF. Deficiency symptoms of macronutrients and of boron in hybrids clones of eucalyptus grandis with eucalyptus urophylla. Cerne. 2015;8(2): 107-116.
3. Ferreto DOC, Rodrigues JDP, Ibarra MA, De Oliveira CS, Vieira RCB, Weber MA, Vieira FCB. Boron fertilization and liming for Eucalyptus urograndis cropped on sandy arenosol of Brazilian pampa. Journal of plant nutrition. 2016;39(3),399-409.
4. Albaugh TJ, Rubilar RA, Maier CA, Acuña EA, Cook RL. Biomass and nutrient mass of Acacia dealbata and eucalyptus globulus bioenergy plantations. Biomass and Bioenergy. 2017;97:162-171.
5. Morais E, Zanotto ACS, Freitas MLM, Moraes MLT, Sebbenn AM. Variação

- genética, interação genótipo solo e ganhos na seleção em teste de progênes de *Corymbia citriodora* Hook em Luiz Antonio, São Paulo. Scientia Forestalis. 2010;38(85):11-18. Portuguese.
6. Barros Filho NF. Discriminação isotópica do ¹³C e nutrição com cálcio e boro em clones de eucalipto submetidos ao déficit hídrico. 2014. Universidade Federal de Viçosa. 2014;59. Portuguese.
 7. Marschner H. Mineral nutrition of higher plants; 2012.
 8. Hu H, Brown PH, Labavitch JM. Species variability in boron requirements is correlated with cell wall pectin. Journal Experimental Botany. 1996;(47):227-232.
 9. Barretto VCM, Valeri SV, Silveira RLVA, Takahashi EO. Eficiência de uso de boro no crescimento de clones de eucalipto em vasos. Scientia Forestalis. 2007;76:21-33. Portuguese.
 10. Silva PHM, Campoe OC, Vieira IG, Paula RC. Aplicação foliar de boro em eucalipto sob estresse hídrico. Scientia Forestalis. 2015;43(106):395-405. Portuguese.
 11. Gonçalves JLM, Valeri SV. Micronutrientes para culturas: Eucalipto e pinus. In: Ferreira, M. E. et al. (Ed.). Micronutrientes e elementos tóxicos na agricultura. Jaboticabal. 2001;393-423. Portuguese.
 12. Mattiello EM, Ruiz HA, Silva IR, Guerra PC, Andrade VM. Características fisiológicas e crescimento de clones de eucalipto em resposta ao boro. Revista Árvore. 2009;33(5):821-830. Portuguese.
 13. Andrade JD, Cantarella H, Quaggio JA. Análise química para avaliação da fertilidade de solos tropicais. Instituto Agrônomo. 2001;47. Portuguese.
 14. Tedesco MJ, Gianello C, Bissani CA, Bohnen H, Volkweiss SJ. Análise de solo, plantas e outros materiais. 2. Ed. Boletim Técnico de Solos. 1995;174. Portuguese.
 15. Banzatto D, Kronka SN, Experimentação Agrícola, Funep Unesp, Jaboticabal SP. 1989;245. Portuguese.
 16. Ramos SJ, Faquin V, Ferreira RMA, Araújo JL, Carvalho JGD. Crescimento e teores de boro em plantas de eucalipto (*Eucalyptus citriodora*) cultivadas em dois latossolos sob influência de doses de boro e disponibilidade de água. Revista Árvore. 2009;33(1):57-65. Portuguese.
 17. Dias LPR, Gatiboni LC, Miquelutti DJ, Brunetto G, Campos DJP. Boron in preventing of shoot dieback on *Eucalyptus benthamii* grown in soil with different water regimes. Scientia Agrarian. 2017;18(3):37-43.
 18. Gupta UC. Factors affecting boron uptake by plants. In: Gupta U.C. (Ed.). Boron and its Role in Crop Production. 1993:87-104.
 19. Dechen AR, Haag HP, Carmello QAC. Funções dos micronutrientes nas plantas. In: Ferreira M.E, Cruz MCP. (Ed.). Micronutrientes Na Agricultura. 1991;65-98. Portuguese.
 20. Sgarbi F, Silveira RLA, Higashi EN, Gonçalves NA. Avaliação do estado nutricional e da fertilidade do solo em plantios de *Eucalyptus grandis* e *Eucalyptus saligna* na região de Capão Bonito/SP (compact disc). In: Reunião Brasileira de Fertilidade do Solo e Nutrição de Plantas, Santa Maria; 2000. Portuguese.
 21. Sgarbi F, Silveira RLA, Higashi EN, Neto CBL. Avaliação do estado nutricional e da fertilidade do solo em plantios de *Eucalyptus grandis* na região de Lençóis Paulista/SP (compact disc). In: Reunião Brasileira de Fertilidade do Solo e Nutrição de Plantas, Santa Maria. 2000. Portuguese.
 22. Nunes FN. Crescimento e expressão gênica em clones de eucalipto influenciados pelo boro e déficit hídrico, Universidade Federal de Viçosa. 2010;65. Portuguese.
 23. Tirloni C, Daniel O, Vitorino ACT, Novelino JO, Carducci CE, Heid DM. Crescimento de *Corymbia citriodora* sob aplicação de boro nas épocas secas e chuvosas no Mato Grosso do Sul, Brasil. Silva Lusitana. 2011;19(2):185-194. Portuguese.
 24. Paula TA. Doses, fontes e formas de aplicação de boro em floresta de eucalipto. Universidade de São Paulo, Piracicaba. 2009;67. Portuguese.
 25. Reichardt K. Processos de transferência no sistema solo-planta-atmosfera, Fundação Cargill. 1985;466. Portuguese.
 26. Sakya AT, Dell B, Huang L. Boron requirements for eucalyptus globulus seedlings. Plant and Soil. 2002;246(1):87-95.
 27. Barros NF, Novais RF. Eucalyptus nutrition and fertilizer regimes in Brazil. In: Attiwill PM, Ada MA. (Ed.). Nutrition of Eucalyptus. Collin-wood: CSIRO. 1996; 335-355.
 28. Ferreira GB. Interferências da matéria orgânica e ferro na dosagem de boro com

- azometina-H e comparação de extratores para boro disponível no solo. Universidade Federal de Viçosa. 1998;75. Portuguese.
29. Shorrocks VM. The occurrence and correction of boron deficiency. *Plant and Soil*. 1997;193(1-2):121-148.
30. Brown PH, Shelp BJ. Boron mobility in plants. *Plant Soil*. 1997;193(1):85-101. 1997.
31. Hu H, Brown PH. Absorption of boron by plants roots. In: Dell B, Brown, PH, Bell RW. (Ed.). *Boron in Soils and Plants: Reviews*, Dordrecht. Kluwer Academic. 1997;193(1-2):49-58.
32. Furlani AMC. Nutrição mineral. In: Kerbauy, G. B. (Org.) *Fisiologia Vegetal*. 2004;452. Portuguese.
33. Santana RC, Barros NF, Novais RF, Leite HG, Comerford NB. Alocação de nutrientes em plantios de eucalipto no Brasil. *Revista Brasileira de Ciências do Solo*. 2008;32:2723-2733. Portuguese.

© 2019 Pinheiro et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<http://www.sdiarticle3.com/review-history/48777>