



## Further Scalable Test Functions for Multidimensional Continuous Optimization

George Anescu<sup>1\*</sup>

<sup>1</sup>Faculty of Power Plant Engineering, Polytechnic University of Bucharest, Romania.

### Author's contribution

The sole author designed, analysed, interpreted and prepared the manuscript.

### Article Information

DOI: 10.9734/JAMCS/2019/v34i430221

*Editor(s):*

(1) Prof. Radko Mesiar, DSc., Department of Mathematics, Faculty of Civil Engineering, Slovak University of Technology Bratislava, Radlinskho 11, 813 68 Bratislava, Slovakia.

*Reviewers:*

(1) K. Lenin, Jawaharlal Nehru Technological University Hyderabad, India.

(2) M. Bhanu Sridhar, G. V. P. College of Engineering for Women, India.

Complete Peer review History: <http://www.sdiarticle4.com/review-history/52793>

*Received: 29 September 2019*

*Accepted: 06 December 2019*

*Published: 14 December 2019*

**Original Research Article**

## Abstract

Multidimensional scalable test functions are very important in testing the capabilities of new optimization methods, especially in evaluating their response to the increase of the search space dimension. As a continuation of a previous published paper, new sets of test functions for continuous optimization are proposed, both unconstrained (or only box constrained, 7 new test functions) and constrained (10 new test functions).

*Keywords: Optimization; Continuous Global Optimization Problem (CGOP); Unconstrained Optimization; Constrained Optimization.*

**2010 Mathematics Subject Classification:** 90C06, 90C25, 90C26, 90C30.

## 1 Introduction

The real world optimization problems that emerge from various scientific and engineering fields are characterized by complexity, non-linearity and increased numbers of decision variables and constraints. In order to be able to handle such difficult problems the researchers in the optimization

\*Corresponding author: E-mail: [george.anescu@gmail.com](mailto:george.anescu@gmail.com);

field are continually proposing new improved optimization algorithms. Due to the intrinsic mathematical difficulty of the global optimization problem, in the last decades there is a trend in researching new nature inspired optimization algorithms, capable to provide acceptable solutions in convenient computing time, even though the global solution is not guaranteed. Such nature inspired algorithms, also named meta-heuristic, or population based algorithms, have some advantages over the traditional gradient based algorithms: they are able to handle more general classes of optimization problems, are derivative free (can be successfully applied when the derivatives are not available or do not exist) and can be easily parallelized on modern multiprocessor computers. Before the newly proposed optimization methods are applied to real world optimization problems, their properties are extensively evaluated by using known test functions from standard literature. In most of the cases the global solutions of the test functions are theoretically known, but sometimes only the best experimentally found solutions are available (for the so called open problems) and any improvement to the best known solutions provided by the tested optimization algorithm is considered as a competitive advantage. One important property that modern optimization methods need (especially in modern Big Data applications) is scalability, i.e. the ability to respond well when the dimension of the search space increases. In order to appropriately evaluate the scalability property of the optimization methods there is a need of multidimensional scalable test functions. Many sets of optimization test functions (benchmarks) are already known from the literature (see [1], [2], [3], [4], [5], etc.), but there is still a need for multidimensional scalable test functions. The present paper is a continuation of a previously published paper, [6], both papers having the purpose to supplement the known collections of optimization test functions with some new proposals of multidimensional scalable problems, especially deceptive problems (for which the size of the basin of attraction of the global solution is small compared to the sizes of the basins of attractions of some local solutions), which can prove useful in further testing and comparing the capabilities of the numerous modern optimization methods. The indexing of the proposed test functions continues from [6] with  $f_{14}$  for unconstrained optimization test functions and  $f_{30}$  for constrained optimization test functions. Initially published as a working paper on the Research Gate website, the paper already accumulated 4 citations [7], [8], [9], [10] which is a further proof of its usefulness.

The rest of the paper is organized as follows: Section 2 presents the general model of Continuous Global Optimization Problem (*CGOP*); Section 3 presents the additional proposed unconstrained (or box constrained) optimization test functions; Section 4 presents the additional proposed constrained optimization test functions; and finally, Section 5 summarizes and draws some conclusions.

## 2 Continuous Global Optimization Problem (*CGOP*)

The Continuous Global Optimization Problem (*CGOP*) is generally formulated as [11]:

$$\text{minimize} \quad f(\mathbf{x}) \quad (2.1)$$

$$\text{subject to} \quad \mathbf{x} \in D$$

with

$$D = \{\mathbf{x} : \mathbf{l} \leq \mathbf{x} \leq \mathbf{u}; \text{ and } g_i(\mathbf{x}) \leq 0, \quad i = 1, \dots, G; \\ \text{and } h_j(\mathbf{x}) = 0, \quad j = 1, \dots, H\} \quad (2.2)$$

where  $\mathbf{x} \in \mathbb{R}^n$  is a real  $n$ -dimensional vector of decision variables ( $\mathbf{x} = (x_1, x_2, \dots, x_n)$ ),  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  is the continuous objective function,  $D \subset \mathbb{R}^n$  is the non-empty set of feasible decisions (a proper subset of  $\mathbb{R}^n$ ),  $\mathbf{l}$  and  $\mathbf{u}$  are explicit, finite (component-wise) lower and upper bounds on  $\mathbf{x}$ ,  $g_i : \mathbb{R}^n \rightarrow \mathbb{R}$ ,  $i = 1, \dots, G$  is a finite collection of continuous inequality constraint functions, and

$h_j : \mathbb{R}^n \rightarrow \mathbb{R}$ ,  $j = 1, \dots, H$  is a finite collection of continuous equality constraint functions. In the black box approach of the *CGOP* problem, which is specific for the derivative free meta-heuristic population based optimization methods, no other additional suppositions are made and it is assumed that no additional knowledge about the collections of real continuous functions can be obtained, i.e. for any point  $\mathbf{x}$  in the boxed domain  $\{\mathbf{x} : \mathbf{l} \leq \mathbf{x} \leq \mathbf{u}\}$  it is assumed the ability to calculate the values of the functions  $f(\mathbf{x})$ ,  $g_i(\mathbf{x})$ ,  $i = 1, \dots, G$ ,  $h_j(\mathbf{x})$ ,  $j = 1, \dots, H$ , but nothing more. However, in the gradient based optimization methods it is assumed that the methods have also access to the derivatives of the mentioned set of functions (if they exist).

The general mathematical model presented in this section will be applied to the formal presentations of all the proposed optimization test functions in the next sections of the paper. All the proposed functions are multidimensional and scalable to the dimension of the search space,  $n$ . All other properties of the functions (such as, unimodality or multimodality) are described when known. The known global solutions (theoretically provable), or the best known global solutions (for open problems) are also specified. All the numerical results presented were obtained by applying metaheuristic optimization methods (see the methods presented in [12], [13] and [14]).

### 3 Unconstrained Optimization Test Functions

In this section 7 ( $f_{14} - f_{20}$ ) additional new test functions are proposed to the set of unconstrained (or box constrained) optimization test functions.




Fig. 1. Difficult optimization problem (2-dimensional case)




Fig. 2. Example of non-quasi-convex unimodal function [15]

- $f_{14}$  (*Perturbed Sphere Function*) - multimodal, *global minimum* value (theoretical)  $f_{14}^* = -1$  at  $\mathbf{x}^* = (1, 1, \dots, 1)$ . This is another very difficult test function, if approached as a black box model, due to the small dimension of the attraction basin of the global minimum, which is contained in a larger attraction basin of the local minima at  $\mathbf{x}^* = (0, 0, \dots, 0)$ . Usually the optimization methods are trapped by the local minimum with the larger attraction basin. A graphical representation of this function is given in Fig. 1 for the 2-dimensional case:

$$f_{14}(\mathbf{x}) = -(n+1)e^{-10\sqrt{n}\left[\sum_{j=1}^n (x_j - 1)^2\right]^{1/2}} + \sum_{j=1}^n x_j^2 \quad (3.1)$$

$$n \geq 1, \quad -2 \leq x_j \leq 2, \quad j = 1, \dots, n$$

- $f_{15}$  - multimodal, *global minimum* value (theoretical)  $f_{15}^* = 0$  at  $\mathbf{x}^* = (1, 1, \dots, 1)$ :

$$f_{15}(\mathbf{x}) = \sum_{j=1}^n \left(2x_{j-1} - x_j - \frac{1}{x_j x_{j+1}}\right)^2 \quad (3.2)$$

$$n \geq 3, \quad x_{n+1} = x_1, \quad x_0 = x_{n-1}, \quad 10^{-6} \leq x_j \leq 2, \quad j = 1, \dots, n$$




Fig. 3. Example of quasi-convex unimodal function with almost flat region [15]

- $f_{16}$  - multimodal, *global minimum* value (theoretical)  $f_{16}^* = n$  at  $\mathbf{x}^* = (2, 2, \dots, 2)$ . The function has many local minima which have the potential of trapping the optimization methods, notably the one located close to  $(3.2, 3.2, \dots, 3.2)$ :

$$f_{16}(\mathbf{x}) = \sum_{j=1}^n \log_{x_j} (|x_{j-1}x_{j+1} - 5.2x_j + 6.4| + x_j) \quad (3.3)$$

$n \geq 3, x_{n+1} = x_1, x_0 = x_{n-1}, 1.000001 \leq x_j \leq 3.3, j = 1, \dots, n$

- $f_{17}$  - unimodal, *global maximum* value (theoretical)  $f_{17}^* = 4$  at  $\mathbf{x}^* = (2, 2, \dots, 2)$ :

$$f_{17}(\mathbf{x}) = \min\{3x_1 + x_2 - x_3^2, 3x_2 + x_3 - x_4^2, \dots, 3x_n + x_1 - x_2^2\} \quad (3.4)$$

$n \geq 3, 0 \leq x_j \leq 4, j = 1, \dots, n$

- $f_{18}$  - unimodal, *global maximum* value (theoretical)  $f_{18}^* = 1$  at  $\mathbf{x}^* = (1, 1, \dots, 1)$ :

$$f_{18}(\mathbf{x}) = \min\{3x_1x_2 - 2x_3^3, 3x_2x_3 - 2x_4^3, \dots, 3x_nx_1 - 2x_2^3\} \quad (3.5)$$

$n \geq 3, 0 \leq x_j \leq 2, j = 1, \dots, n$

- $f_{19}$  - Not Quasi-Convex Function, unimodal, *global maximum* value (theoretical)  $f_{20}^* = 0$  at  $\mathbf{x}^* = (0, 0, \dots, 0)$ :

$$f_{19}(\mathbf{x}) = \sum_{j=1}^n x_j^4 + 16 \sum_{j=1}^n x_j^2 x_{j+1}^2, \quad (3.6)$$

$$-2 \leq x_j \leq 2, \quad j = 1, \dots, n, \quad x_{n+1} = x_1,$$

This is an example of unimodal function that is not quasi-convex [15], the 2-dimensional case being presented in Fig. 2.

- $f_{20}$  - Plateau Function, unimodal, *global minimum* value (theoretical)  $f_{20}^* = -1$  at  $\mathbf{x}^* = (1, 1, \dots, 1)$ :

$$f_{20}(\mathbf{x}) = -(n+1)e^{-10\sqrt{n} \left[ \sum_{j=1}^n (x_j - 1)^2 \right]^{1/2}} + \max \left\{ \sum_{j=1}^n x_j^2, n + \frac{0.01}{n} - 0.2 \right\} \quad (3.7)$$

$$-2 \leq x_j \leq 2, \quad j = 1, \dots, n,$$

This is an example of a difficult unimodal function having an almost flat region with a very narrow escape path to the minimum point [15]. The 2-dimensional case is presented in Fig. 3.

## 4 Constrained Optimization Test Functions

In this section 10 ( $f_{30} - f_{39}$ ) additional new test functions are proposed to the set of constrained optimization test functions.

- $f_{30}$  - unimodal, *global maximum* value (theoretical)  $f_{30}^* = \frac{n(n+1)}{2}$  at  $\mathbf{x}^* = \left( \frac{n+1}{2}, \frac{n+1}{2}, \dots, \frac{n+1}{2} \right)$ :

$$f_{30}(\mathbf{x}) = \sum_{j=1}^n x_j,$$

$$g_1(\mathbf{x}) = -n + \sum_{j=1}^n \frac{j}{x_j} \leq 0, \quad (4.1)$$

$$g_2(\mathbf{x}) = -n + \sum_{j=1}^n \frac{n-j+1}{x_j} \leq 0,$$

$$n \geq 2, \quad 0.5 \leq x_j \leq n+1, \quad j = 1, \dots, n$$

- $f_{31}$  - unimodal, *global minimum* value (theoretical)  $f_{31}^* = 0$  at  $\mathbf{x}^* = (1, 1, \dots, 1)$ :

$$f_{31}(\mathbf{x}) = \sum_{j=1}^n \frac{x_j^2}{2x_{j-1}x_{j+1} + 1} - \sum_{j=1}^n \frac{x_j}{x_{j-1} + x_jx_{j+1} + 1},$$

$$g(\mathbf{x}) = 1 - \prod_{j=1}^n x_j \leq 0, \quad (4.2)$$

$$n \geq 3, \quad x_{n+1} = x_1, \quad x_0 = x_{n-1}, \quad 0 \leq x_j \leq 2, \quad j = 1, \dots, n$$

- $f_{32}$  - unimodal, *global minimum* value (theoretical)  $f_{32}^* = 1$  at  $\mathbf{x}^* = \left(\frac{1}{n-1}, \frac{1}{n-1}, \dots, \frac{1}{n-1}\right)$ :

$$f_{32}(\mathbf{x}) = \sum_{j=1}^n \frac{1}{1 + (n-1)^2 x_j},$$

$$g(\mathbf{x}) = (n-1) - \sum_{j=1}^n \frac{1}{1+x_j} \leq 0, \tag{4.3}$$

$$n \geq 2, 0 \leq x_j \leq 1, j = 1, \dots, n$$

- $f_{33}$  - unimodal, *global maximum* value (theoretical)  $f_{33}^* = 3$  at  $\mathbf{x}^* = \left(\frac{1}{n}, \frac{1}{n}, \dots, \frac{1}{n}\right)$ :

$$f_{33}(\mathbf{x}) = \sum_{j=1}^n (3x_{j-1}x_j + 4x_jx_j + 2x_jx_{j+1})^2,$$

$$g(\mathbf{x}) = -1 + \sum_{j=1}^n x_j \leq 0, \tag{4.4}$$

$$n \geq 3, x_{n+1} = x_1, x_0 = x_{n-1}, 0 \leq x_j \leq 1, j = 1, \dots, n$$

- $f_{34}$  - unimodal, *global minimum* value (theoretical)  $f_{34}^* = (n-1)^n$  at  $\mathbf{x}^* = \left(\frac{1}{n-1}, \frac{1}{n-1}, \dots, \frac{1}{n-1}\right)$ :

$$f_{34}(\mathbf{x}) = \prod_{j=1}^n \left(\frac{1}{x_j} - 1\right),$$

$$g(\mathbf{x}) = -1 + \sum_{j=1}^n x_j \leq 0, \tag{4.5}$$

$$n \geq 2, 10^{-6} \leq x_j \leq 1, j = 1, \dots, n$$

- $f_{35}$  - unimodal, *global maximum* value (theoretical)  $f_{35}^* = 1$  at  $\mathbf{x}^* = (1, 1, \dots, 1)$ :

$$f_{35}(\mathbf{x}) = \sum_{j=1}^n \frac{1}{\left(1 - x_j^2 + \sum_{j_1=1}^n x_{j_1}^2\right)},$$

$$g(\mathbf{x}) = n - \left(\prod_{j=1}^n x_j\right) \left(\sum_{j=1}^n x_j\right) \leq 0, \tag{4.6}$$

$$n \geq 2, 0 \leq x_j \leq 2, j = 1, \dots, n$$

- $f_{36}$  - unimodal, *global maximum* value (theoretical)  $f_{36}^* = \sum_{j=1}^n \frac{1}{j}$  at

$$\mathbf{x}^* = (1, 2^{1/2}, 3^{1/3}, \dots, n^{1/n}):$$

$$\begin{aligned} f_{36}(\mathbf{x}) &= \sum_{j=1}^n \frac{1}{x_j^j}, \\ g(\mathbf{x}) &= \prod_{j=1}^n x_j - \prod_{j=1}^n j^{\frac{1}{j}} \leq 0, \\ n &\geq 2, \quad 10^{-6} \leq x_j \leq 2, \quad j = 1, \dots, n \end{aligned} \tag{4.7}$$

- $f_{37}$  - unimodal, *global minimum* value (theoretical)  $f_{37}^* = 0$  at  $\mathbf{x}^* = (0, 0, \dots, 0)$ :

$$\begin{aligned} f_{37}(\mathbf{x}) &= \sum_{j=1}^n |x_j|^3 - \prod_{j=1}^n x_j, \\ g(\mathbf{x}) &= \prod_{j=1}^n x_j - \sum_{j=1}^n x_j \leq 0, \\ n &\geq 2, \quad -10 \leq x_j \leq 10, \quad j = 1, \dots, n \end{aligned} \tag{4.8}$$

- $f_{38}$  - unimodal, *global minimum* value (theoretical)  $f_{38}^* = 1$  at  $\mathbf{x}^* = (1, 1, \dots, 1)$ :

$$\begin{aligned} f_{38}(\mathbf{x}) &= \sum_{j=1}^n \frac{x_j}{\left(1 - x_j + \sum_{j_1=1}^n x_{j_1}\right)}, \\ g(\mathbf{x}) &= -1 + \sum_{j=1}^n \frac{1}{\left(1 - x_j + \sum_{j_1=1}^n x_{j_1}\right)} \leq 0, \\ n &\geq 2, \quad 0 \leq x_j \leq 2, \quad j = 1, \dots, n \end{aligned} \tag{4.9}$$

- $f_{39}$  - unimodal, *global maximum* value (theoretical)  $f_{39}^* = n + 1$  at  $\mathbf{x}^* = (1, 1, \dots, 1)$ :

$$\begin{aligned} f_{39}(\mathbf{x}) &= \sum_{j=1}^n x_j + \frac{1}{n} \left( \sum_{j_1=1}^n x_{j_1}^2 \right), \\ g(\mathbf{x}) &= -(n + 1) + \sum_{j=1}^n x_j^2 + \prod_{j=1}^n x_j \leq 0, \\ n &\geq 2, \quad 0 \leq x_j \leq 2, \quad j = 1, \dots, n \end{aligned} \tag{4.10}$$



## 5 Conclusions

The paper proposed two new sets of optimization test functions: a set of 7 continuous unconstrained (or box constrained) test functions, and a set of 10 continuous constrained test functions. All the proposed functions are multidimensional and scalable to the dimension of the search space  $n$ , which is a useful property when the response in performance (efficiency and success rate) of an optimization method is investigated with respect to the increase in search space dimension. It is the hope of the author that the proposed new sets of optimization test functions will represent a valuable addition to the known collections of optimization test functions and will prove useful in investigating the properties of new or existing optimization methods.

## Competing Interests

Author has declared that no competing interests exist.

## References

- [1] Floudas CA, Pardalos PM. A collection of test problems for constrained global optimization algorithms. Springer-Verlag, Berlin; 1990.
- [2] Hock W, Schittkowski K. Test examples for nonlinear programming codes. Springer-Verlag Berlin Heidelberg, Berlin; 1981.
- [3] Schittkowski K. More test examples for nonlinear programming codes. Springer-Verlag Berlin Heidelberg, Berlin; 1987.
- [4] Momin J, Xin-She Yang. A literature survey of benchmark functions for global optimization problems. *Int. Journal of Mathematical Modelling and Numerical Optimisation*. 2013;4:150-194.
- [5] Molga M, Smutnicki C. Test functions for optimization needs. 2005;1-43. Available:<http://www.robertmarks.org/Classes/ENGR5358/Papers/functions.pdf> (Last time accessed in November, 2019)
- [6] Anescu G. Scalable test functions for multidimensional continuous optimization. *U.P.B. Sci. Bull., Series C*. 2017;79(4):27-42.
- [7] Glvez J, Cuevas E, Hinojosa S, Avalos O, Prez-Cisnerosa M. A reactive model based on neighborhood consensus for continuous optimization. *Expert Systems with Applications*. 2019;121:115-141.
- [8] Cuevas E, Glvez J. An optimization algorithm guided by a machine learning approach. *International Journal of Machine Learning and Cybernetics*. 2019;10:129.
- [9] Cuevas E, Glvez J, Avalos O. Knowledge-based optimization algorithm. In Book: *Recent Metaheuristics Algorithms for Parameter Identification*, Springer International Publishing; 2020.
- [10] Cuevas E, Glvez J, Avalos O. Neighborhood based optimization algorithm. In Book: *Recent Metaheuristics Algorithms for Parameter Identification*, Springer International Publishing; 2020.
- [11] Pintér JD. Global optimization: Software, test problems, and applications. Ch. 15 in *Handbook of Global Optimization*, (Ed. P. M. Pardalos and H. F. Romeijn), Kluwer Academic Publishers, Dordrecht, Boston, London. 2002;2:515-569.

- [12] Anescu G, Prisecaru I. NSC-PSO, a novel PSO variant without speeds and coefficients. In: Proceedings of the 17th International Symposium on Symbolic and Numeric Algorithms for Scientific Computing, SYNASC 2015, Timisoara, Romania: September 21-24. 2015;460-467.
- [13] Anescu G. Gradual and cumulative improvements to the classical differential evolution scheme through experiments. Annals of West University of Timisoara - Mathematics and Computer Science. 2016;54(2):13-36.
- [14] Anescu G. A fast artificial bee colony algorithm variant for continuous global optimization problems. U.P.B. Sci. Bull., Series C. 2017;79(1):83-98.
- [15] Anescu G. A heuristic fast gradient descent method for unimodal optimization. Journal of Advances in Mathematics and Computer Science. 2018;26(5):1-20.

---

©2019 Anescu; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution and reproduction in any medium, provided the original work is properly cited.

**Peer-review history:**

The peer review history for this paper can be accessed here (Please copy paste the total link in your browser address bar)

<http://www.sdiarticle4.com/review-history/52793>