

A Green and Scalable Synthesis of 1-Amino Anthraquinone

Bin Lu^{1#}, Yi-Yu Yan^{1#}, Yong-Fu Qiu^{1#}, Tian-Li Zhang^{1#}, Shi Qi¹, Jian-Hua Tian¹,
Wan-Yue Luo¹, Xiao Hu^{2*} and Jin Wang^{1*}

¹School of Pharmacy, Jiangsu Key Laboratory for Bioresources of Saline Soils, Yancheng Teachers University, Hope Avenue South Road No.2, Yancheng, 224007, Jiangsu Province, P.R. China.

²Yancheng Teachers University Library, Hope Avenue South Road No.2, Yancheng, 224007, Jiangsu Province, P.R. China.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJACR/2019/v3i130081

Editor(s):

(1) Dr. Angélica Machi Lazarin, Professor, Department of Chemistry, State University of Maringá, Brazil.

Reviewers:

(1) Hasan Küçükbay, İnönü University, Turkey.

(2) Charles Emeka Umenwa, University of Ibadan, Nigeria.

Complete Peer review History: <http://www.sdiarticle3.com/review-history/47889>

Received 30 December 2018

Accepted 13 March 2019


Published 20 March 2019

Original Research Article

ABSTRACT

1-amino anthraquinone (2) is the most important intermediate in the synthesis of acid dyes. This paper presents a new method for the preparation of title compound (2) in a highly chemo- and regioselective reduction of 1-nitro anthraquinone (1) by NaHS in water under mild conditions. This protocol is clean, operationally simple, easy work-up and could be applied in the industrial production.

Graphical_Abtract



*Corresponding author: Email: jaxdon@126.com, wangj01@yctu.edu.cn;


These authors contributed equally to this work

Keywords: Nitroanthraquinone; aminoanthraquinone; reduction.

1. INTRODUCTION

1-Amino anthraquinone [1] is one of the most important intermediates [2] in the synthesis of functional dyes [1]. To date, a variety of methods for the synthesis of 1-amino anthraquinone [1] have been disclosed [3], most of these protocols employing 1-nitro anthraquinone [2] as a starting material [4]. 1-amino anthraquinone [1] can be obtained by single-step reduction from compound [2]. These reductants include ammonium formate,[5] sodium sulfide(Na_2S),[1] sodium borohydride (NaBH_4),[6] Gold-Catalyzed $\text{CO-H}_2\text{O}$ system [7] and bis(cyclopentadienyl)titanium(IV) dichloride-indium system [8]. However, none of these reagent is suitable for industrial production due to drawbacks like high cost, toxic substance, complex work-up, etc.[9] In recent years, the demand for 1-amino anthraquinone (2) in the dye industry has increased rapidly. Hence, it is important to developed an efficient and scalable method for synthesis of 1-amino anthraquinone [1].

As shown in Scheme 1, we reported here a facial, green and scalable method for the preparation of 1-amino anthraquinone (2) by using NaHS as a reductant, [10] the solvent water meets the requirements of green chemistry and it should be suitable for industrial production.



Scheme 1. Synthesis of compound 2

2. EXPERIMENTAL SECTION

All reactions were monitored by TLC, Melting points were measured on Melting Point M-565 (BUCHI). NMR and mass spectra were recorded on a Bruker Avanc III-HD 400 NMR and a TripleTOF Mass spectrometers, respectively. All reagents: e.g. $\text{Na}_2\text{S}\cdot 9\text{H}_2\text{O}$, NaSH, NaBH_4 , $\text{Na}_2\text{S}_2\text{O}_4$ were purchased from Adamas, P. R. China, and used without further purification [11].

Synthesis of 1-amino anthraquinone (2)

A 250 mL three-necked flask is equipped with a stirrer and thermometer and a dropping funnel. The flask is charged with a solution of NaSH (3.00 g, 0.05 mol) in water (15 mL) and stirred at 60°C for 1h. Then 1-nitro anthraquinone (1) powder (3.04 g, 0.01 mol) was added over 10 minutes and the reaction mixture quickly turned to red. The mixture was stirred at 60°C for another 1h and the progress of the reaction was monitored by TLC. The reaction mixture was filtered and the red precipitates were washed with water, and recrystallization from ethanol to give compound 2 as a red powder (2.79 g, yield 92%), m.p. 253°C (lit[8]253-255 $^\circ\text{C}$)


^1H NMR (400 MHz, CDCl_3): δ 8.30 (d, $J = 8.0$ Hz, 1H), 8.26 (d, $J = 8.0$ Hz, 1H), 7.78 (t, $J = 8.0$ Hz, 1H), 7.73 (t, $J = 8.0$ Hz, 1H), 7.65 (d, $J = 8.0$ Hz, 1H), 7.47 (t, $J = 8.0$ Hz, 1H), 6.98 (d, $J = 8.0$ Hz, 1H), 6.87 (s, 2H).

^{13}C NMR (100 MHz, CDCl_3): δ 185.3, 183.6, 151.0, 134.8 (2C), 134.4, 134.0, 133.2 (2C), 126.8 (2C), 123.1, 117.3, 113.7.

MS(ESI): $m/z = 224$ (M+H)

3. RESULTS AND DISCUSSION

The key factor to obtain compound 2 is how to efficiently and selectively reduce the nitro-group of compound 1 without affecting the carbonyl group [12]. We investigated the effects of different reagents and solvents, the results were shown in Table 1. The reaction solvent plays an important role in this reaction, water is better than alcohols or the alcohol solutions. Both Na_2S and NaHS can be severe as a good reducing agents, but when the reaction scale is kilogram level, we found that Na_2S is difficult to agitate in the 5L three round-bottomed flasks, while NaHS do not have this problem. Based on this point, NaHS is much more suitable in industrial large-scale production. The poor solubility of sulfide in ethanol lead to a decrease of yield. We also examined reagents NaBH_4 and $\text{Na}_2\text{S}_2\text{O}_4$, which gave compound 2 in 45% and 35% yield respectively. The optimal condition was using AgNO_3 (40%), and $\text{K}_2\text{S}_2\text{O}_8$ (2 equiv) in water at 60°C for 2 h (entry 4, Table 1).

Table 1. Reduction of 1-nitroanthraquinone (1) under different conditions

Entry	Reagent	Solvent(v/v)	Temperature(°C)	Yield(%)
1	Na ₂ S	H ₂ O	60	80%
2	Na ₂ S	EtOH	60	35%
3	Na ₂ S	EtOH/ H ₂ O(1/1)	60	63%
4	NaSH	H ₂ O	60	92%
5	NaSH	EtOH	60	24%
6	NaSH	EtOH/ H ₂ O(1/1)	60	59%
7	NaBH ₄	isopropanol	60	45%
8	Na ₂ S ₂ O ₄	EtOH	60	35%

Reaction Conditions: compound 1 (0.05mol), reagent (2 equiv), 2 hour under open air

4. CONCLUSION

In summary, a NaHS-mediated new method for the synthesis of 1-amino anthraquinone (2) has been developed. This protocol is easily operational, efficient, and is amenable to the kilogram-scale synthesis of compound (2). This chemistry also provides a new selective reduction of aromatic nitro-group without using metal catalyst.

ACKNOWLEDGEMENTS

This study was supported by the National Natural Science Foundation of China (Nos. 31600740 and 81803353), the Natural Science Foundation of Jiangsu Province (BK20160443), the Six Talent Peaks Project in Jiangsu Province (SWYY-094), the Jiangsu Provincial Key Laboratory for Bioresources of Saline Soils (Nos. JKLBS2016013 and JKLBS2017010).


COMPETING INTERESTS

Authors have declared that no competing interests exist.


REFERENCES

- Ghaieni H, Sharifi M, Fattollahy M. Dyes and Pigments. 2007;72:97-100.
- Sondhi SM, Sharma VK, Singhal N, Verma RP, Shukla R, Raghubir R, Dubey MP. Synthesis and anti-inflammatory activity evaluation of some acridinyl amino antipyrine, acridinyl amino anthraquinone, acridino thiourea and thiazolino thiourea derivatives. Phosphorus, sulfur, and silicon and the Related Elements. 2000;156:21-33.
- Malik EM, Baqi Y, Müller CE. Syntheses of 2-substituted 1-amino-4-bromoanthraquinones (bromaminic acid analogues) – precursors for dyes and drugs Beilstein Journal of Organic Chemistry. 2015;11: 2326-2333.
- Ghaieni H, Rostamizadeh S, Fattollahy M, Aryan R, Tavangar S. Dyes and Pigments. 2008;77:483-486.
- Lou XB, He L, Qian Y, Liu YM, Cao Y, Fan KN. Cheminform. 2015;353:281-286.
- Morley JO. Synthesis 1976, 1976, 528-530.
- H Lin, W. Lu-Cun, S. Hao, N. Ji, C. Yong, H. He-Yong, F. Kang-Nian, Angewandte Chemie 2010, 48, 9538-9541.
- B. W. Yoo, S. J. Lee, B. S. Yoo, K. I. Choi, J. H. Kim, Synthetic Communications 2002, 32, 2489-2493.
- Wang J, Xia F, Jin WB, Guan JY, Zhao H. Bioorganic Chemistry. 2016;68:214-218.
- Wang J, Gasc F, Prandi J. Samarium Diiodide Mediated Coupling of 2 - Pyridylsulfonyl Furanosides with Aldehydes and Ketones: A General Synthesis of C-Furanosides European Journal of Organic Chemistry. 2015;2691-2697.
- Wang J, Li S, Yang T, Yang J. Tetrahedron. 2014;70:9029-9032.
- Wang J, Li S, Yang T, Yang J. Synthesis and antioxidant activities of Coenzyme Q analogues. European Journal of Medicinal Chemistry. 2014;86:710-713.

APPENDIX



¹H NMR OF COMPOUND 2



¹³C NMR of compound 2

© 2019 Qiu et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<http://www.sdiarticle3.com/review-history/47889>