



Efficacy of Nano Fertilizers on Yield, Attributes and Economics of Wheat

Y. K. Singh ^{a*}, Bal Veer Singh ^{a*}, Drishty Katiyar ^{b++},
D. R. K. Saikanth ^{c#}, Kaushal Kumar ^d, Omkar Singh ^{e#},
Shikhar Verma ^a and Pramod Kumar ^a

^a Department of Agronomy, Chandra Shekhar Azad University of Agriculture and Technology, Kanpur, U.P.-208002, India.

^b Soil Science and Agricultural Chemistry, Chandra Shekhar Azad University of Agriculture and Technology, Kanpur, U.P.-208002, India.

^c ICAR -ATARI, Zone -X, Hyderabad, Telangana, India.

^d Department of SCWM, Chandra Shekhar Azad University of Agriculture and Technology, Kanpur, U.P.-208002, India.

^e Sardar Vallabhbhai Patel University of Agriculture and Technology, Meerut, U.P.-250110, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJECC/2023/v13i71879

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/98586>

Original Research Article

Received: 19/02/2023

Accepted: 21/04/2023

Published: 04/05/2023

ABSTRACT

The present investigation was carried out to study the effect of foliar application of nano-fertilizers N and P on yield, and the economics of wheat in Semi-arid and sub-tropical region of Central Plain Zone of Uttar Pradesh. The experiment was laid out in triplicate following a randomized block design with twelve treatments comprising foliar application semi-different doses of Nano N and P. Foliar application of 1st spray of Nano N and P at 30 days after sowing (DAS) + 2nd spray of Nano N

⁺⁺ PhD Research Scholar;

[#] Senior Research Fellow;

*Corresponding author: E-mail: dryksingh1209@gmail.com, bvs955rajput@gmail.com;

and Zn at 45 DAS along with 75% recommended dose of fertilizers (RDF) significantly ($P=0.05$) increased yield and have better economics. Foliar application of nano-fertilizers leads to significant improvement of crop productivity of wheat in Semi- arid and sub-tropical region of Uttar Pradesh. Moreover, the foliar application of nano-fertilizers, *i.e.*, Nano N has direct role in increasing yield as nutrient get easily available to plant in case of foliar spray.

Keywords: Foliar application; nano-fertilizers; wheat productivity; wheat economics.

1. INTRODUCTION

Wheat (*Triticum aestivum* L.) is the staple food of 40% human population across the globe and second most important cereal after rice. This covers world 222.27 million ha, produces 779.3 million metric tonnes, and yields 3.51 tonnes per hectare [1]. In India, wheat plays a vital role in food and nutritional security with an area of 31.31 million ha and an output of 109.6 million metric tonnes with average productivity of 3.52 Metric tonnes per hectare [2]. The world's 7.7 billion people will grow to 9.7 billion by 2050. India is the second-most populated country (1.3 billion) after China (1.41 billion) and is anticipated to surpass China in seven years, reaching 1.7 billion by 2050 [3]. Wheat will likely remain crucial to global food security.

Low nutrient use efficiency is an issue when building and evaluating agricultural production systems. It can be considerably affected by fertilizer, soil, and water management to maximize crop output and minimize nutrient loss from the field [4]. The indiscriminate use of inorganic fertilizers for the past 50 years has reduced the quality and quantity of agricultural produce. Due to the global energy crisis and the rising cost of chemical fertilizer, nano-fertilizer is gaining popularity as a plant nutrient. A correct blend of nano and inorganic fertilizer is needed to increase yield and maintain soil health. Wheat needs nitrogen and phosphorus for growth and yield. Nano-fertilizer reduces soil contamination and boosts fertility. Nanomaterials alone or with inorganic fertilizers can improve soil nutrition and fertility.

Nano-scale matter has novel, distinct properties than macroscopic matter. Reduced molecular size and changing molecular interactions change characteristics. High reactivity, increased bioavailability and bioactivity, adhesion effects, and surface effects of nanoparticles are important in agricultural revolution [5]. Custom-made things are formed from atoms; their qualities depend on their arrangement. Nano fertilizers increase growth metrics (plant height,

leaf area, number of leaves per plant), dry matter production, chlorophyll production, and photosynthesis rate, resulting in higher photosynthetic production and translocation than standard fertilizers [6,7].

Foliar applications nourish plants by spraying liquid fertilizers or other chemical or natural products directly on the leaves of macro and micronutrients for optimal production and reduced losses [8]. Optimal N management for wheat maximizes productivity, water use, and environmental contamination [9,10].

To improve seed maturation and development, phosphorus is necessary [11]. Liakas et al. [12] P treatment encourages tillering of wheat and decreases lodging in wheat, as well as improving photosynthetic activity and transfer to the maturing grains [13]. This makes the grains heavier [14,15]. Wheat grain production can increase by 20% with proper phosphorus application [16]. Phosphorus applications could rise, enhancing N and P absorption [17]. Various researches suggested various P application rates.

As a result, we wanted to find out how foliar feeding of Nano-fertilizers of N and P compared to the control and traditional fertilizer affected several yield-related parameters.

2. MATERIALS AND METHODS

Study Area: The experiment was conducted during rabi 2021-2022 at the Student's Instructional Farm, Chandra Shekhar Azad University of Agriculture and Technology, Kanpur. The region falls under central plain zone of Uttar Pradesh India. The climate of the study area is semi- arid and sub-tropical with an average minimum and maximum temperature (January- April) ranging between 4.3°C to 36.9°C during winter season.

Experimental design and treatments: The experiment was laid out in a randomized block design and replicated thrice in the plot size of

3.20 m x 2.20 m (7.04 m²). The experiment consisted of 12 treatments viz., T1: Control; T2: 100% NPK (150:60:40 kg ha⁻¹); T3: 75% NPK (112.5: 45: 30 kg ha⁻¹); T4: 75% NPK + Nano N spray at 30 Days After Sowing (DAS); T5: 75% NPK + Nano N spray at 45 DAS; T6: 75% NPK + Nano N spray at 30 and 45 DAS; T7: 75% NPK + Nano P spray at 30 DAS; T8: 75% NPK + Nano P spray at 45 DAS; T9: 75% NPK + Nano P spray at 30 and 45DAS; T10: 75% NPK + Nano N and Nano P spray at 30 DAS; T11: 75% NPK + Nano N and Nano P spray at 45 DAS; T12: 75% NPK + Nano N and Nano P spray at 30 and 45 DAS were applied to wheat variety Karan Vandana (DBW 187) in rabi season of 2021-22. The recommended dose of fertilizer were 150 kg N, 60 kg P₂O₅, 40 kg K₂O ha⁻¹ as was used in T2 and also other agronomic practices recommended for central plain Zone were followed as per package and practices, Government of Uttar Pradesh. The grains obtained from the produce of individual plot were recorded as grain yield kg per plot and later it was converted into kg ha⁻¹.

Application protocol of foliar spray: Foliar sprays were given as 1st application at 30 days after sowing and 2nd application at 45 days after sowing in selected treatments with the help of knapsack sprayer fitted with flat fan nozzle. Nano N @ 4 ml l⁻¹ water and Nano DAP @ 250 ml acre⁻¹ with @ 500 l water ha⁻¹ as per scheduled treatments.

Estimation of yield: Yield character was recorded as per standard methods [18]. The grain and straw yield was recorded from the net plot area of each treatment.

Economic Analysis: Economic analysis i.e. (Cost of Cultivation, Gross Returns, Net Returns, Benefit: Cost ratio) as per standard methods [19]. Minimum support price for grain and local prices for straw were used to calculate monetary return of crop.

Statistical analysis: SPSS Version 10.0, SPSS, Chicago, IL, was used for statistical analysis. SPSS was used to determine the statistical significance of treatment effect at 5% probability level. Further, F- test and significance of difference between treatments was investigated using critical difference (CD) as stated by Gomez and Gomez [20].

3. RESULTS AND DISCUSSION

The application of foliar sprays of nano-fertilizer improved the grain, straw and biological yields of

wheat were significantly influenced (Table 1 & Fig. 1). Application of T12 (75% NPK coupled with spray of Nano N and Nano P spray at 30 and 45 DAS) obtained highest grain yield (61.48 q ha⁻¹), straw yield 94.39 (q ha⁻¹) with increased grain yield by 7.25 q ha⁻¹ (11.8%) and 25.27 q ha⁻¹ (41.10%) over 100% NPK and control. T1 and T2 Followed by T10 (75% NPK + Nano N and Nano P spray at 30 DAS), T6 (75% NPK + Nano N spray at 30 and 45 DAS), T11 (75% NPK + Nano N and Nano P spray at 45 DAS) as compared to control. Wheat yield characteristics improves with foliar treatment of N and P nano-fertilizers in combination with conventional fertilizers. These results accord well with those previously reported by Abdel-Aziz et al. [17].

Although fertilizers are crucial for the growth and development of plants, the majority of applied fertilizers remain inaccessible to plants due to a variety of processes, including leaching, photolysis, hydrolysis, and decomposition. The function of these nutrients in promoting plant growth is responsible for the improvement in yield attribute and yield at foliar feeding of nano N and P fertilizer with commercial fertilizers. These components are necessary for the plant to flourish at its best and finish its life cycle [21,22]. In order to increase the quality and production of wheat, the nutrients nitrogen, phosphorus, and potassium are among the most important macronutrients [8]. These processes include the production of chloroplasts and thylakoids as well as the synthesis of chlorophyll [23]. It also participates in the transfer of energy inside the plant, several enzymatic processes, photosynthesis, respiration, and the synthesis of proteins, and as a result, it is essential for plant growth [6]. Wheat was more productive because Nano fertilizer was easily absorbed by the epidermis of leaves and moved to stems, facilitating the uptake of active chemicals [16]. Nano fertilizers contain large surface areas and particles that are smaller than the pore size of plant leaves, which can boost penetration into plant tissues from the applied surface and improve nutrient uptake and usage efficiency [24,25]. It is generally known that applying foliar nitrogen and phosphorus improves photosynthetic capability, maintains healthy leaf nutrition, and maintains carbon balance [26].

The application of Nano mediated fertilizes significantly influenced the economics of wheat. Economics of any crop directly influences the crop growing by the farmer (Table 2 & Fig. 2). It can be directly determined by the B: C ratio of

crop. Crop grown with T12 (75% NPK + Nano N and Nano P) spray at 30 and 45 DAS got highest B: C ratio (2.36) with cost of cultivation of 72548 (\neq ha⁻¹) and net return of 98529.2 (\neq ha⁻¹) followed T10 (75% NPK + Nano N and Nano P spray at 30 DAS), T6 (75% NPK + Nano N spray at 30 and 45DAS), T11 (75% NPK + Nano N and Nano P spray at 45 DAS and lowest in the crop

grown with Control. The higher economics of the wheat is due to the realization of maximum grain yield achieved by the adoption of comprehensive and integrated nitrogen management practices involving optimum level of basal RDF application and foliar nutrition of nano N and P at 30 and 45 DAS stage of the crop [27] and [28].

Table 1. Effect of foliar application of nano-fertilizers on yield of wheat

Symbol	Treatments	Yield (q ha ⁻¹)			
		Grain	Straw	Biological	Harvest index
T ₁	Control	36.21	50.79	89.21	40.59
T ₂	100% NPK (150:60:40 kg ha ⁻¹)	54.23	81.78	136.01	39.87
T ₃	75% NPK (112.5: 45: 30 kg ha ⁻¹)	46.2	66.30	112.5	41.07
T ₄	75% NPK + Nano N spray at 30 DAS	54.32	83.53	137.85	39.41
T ₅	75% NPK + Nano N spray at 45 DAS	53.75	78.54	132.29	40.63
T ₆	75% NPK + Nano N spray at 30 and 45 DAS	56.55	85.45	142	39.82
T ₇	75% NPK + Nano P spray at 30 DAS	53.00	76.41	129.41	40.96
T ₈	75% NPK + Nano P spray at 45 DAS	52.2	73.68	125.88	41.47
T ₉	75% NPK + Nano P spray at 30 and 45 DAS	55.12	79.99	135.11	40.80
T ₁₀	75% NPK + Nano N and Nano P spray at 30 DAS	59.60	87.20	146.8	40.60
T ₁₁	75% NPK + Nano N and Nano P spray at 45 DAS	56.20	82.38	138.58	40.55
T ₁₂	75% NPK + Nano N and Nano P spray at 30 and 45 DAS	61.48	94.39	155.87	39.44
SEM±		2.0	2.87	4.83	0.01
CD (P = 0.05)		5.60	8.24	13.9	NS




Fig. 1. Effect of foliar application of nano-fertilizers on yield of wheat

Table 2. Effect of foliar application of nano-fertilizers on economics of wheat

Symbol	Treatments	Cost of cultivation (₹/ha)	Gross return (₹/ha)	Net return (₹/ha)	B:C Ratio
T ₁	Control	62304	98358.15	36054.2	1.58
T ₂	100% NPK (150:60:40 kg ha ⁻¹)	68294.6	150163.45	81868.9	2.20
T ₃	75% NPK (112.5: 45: 30 kg ha ⁻¹)	66796	126243	59447.0	1.89
T ₄	75% NPK + Nano N spray at 30 DAS	68164	151219.8	83055.8	2.22
T ₅	75% NPK + Nano N spray at 45 DAS	68164	147576.25	79412.3	2.17
T ₆	75% NPK + Nano N spray at 30 and 45DAS	69532	156673.25	87141.3	2.25
T ₇	75% NPK + Nano P spray at 30 DAS	68304	145000	76696.0	2.12
T ₈	75% NPK + Nano P spray at 45 DAS	68304	142023	73719.0	2.08
T ₉	75% NPK + Nano P spray at 30 and 45DAS	69812	151061.8	81249.8	2.16
T ₁₀	75% NPK + Nano N and Nano P spray at30 DAS	69672	163694	94022.0	2.35
T ₁₁	75% NPK + Nano N and Nano P spray at45 DAS	69672	154433	84761.0	2.22
T ₁₂	75% NPK + Nano N and Nano P spray at30 and 45 DAS	72548	171077.2	98529.2	2.36




Fig. 2. Effect of foliar application of nano-fertilizers on economics of wheat

4. CONCLUSION

Based on the above results, it can be concluded that foliar application of nano-fertilizers not only improves the yield but also have a better economics of crop. Nano-fertilizers can serve as an efficient nutrient delivery system thereby reducing the quantity of nutrients required.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. USDA report. U.S. Department of Agriculture, Agricultural Research Service. 2021;22:11-2.
2. USDA report. United States Department of Agriculture, Agricultural Research Service; 2019.
3. The United Nations. World. Popul Prospects. Online edn Rev;1 (Department of Economic and Social Affairs. Population division); 2019.

4. Fixen P, Brentrup F, Bruulsema T, Garcia F, Norton R, Zingore S. Nutrient/fertilizer use efficiency: measurement, current situation and trends. *Manag Water Fert Sustain Agric Intensif.* 2015;270.
5. Gutierrez FJ, Mussons ML, Gatón P, Rojo R. Nanotechnology and food industry. Scientific, health and social aspects of the food industry. In: Tech, Croatia Book Chapter; 2011.
6. Ali EA. Effect of iron nutrient care sprayed on foliage at different physiological growth stages on yield and quality of some durum wheat (*Triticum durum* L.) varieties in Sandy Soil. *Asian J Crop Sci.* 2012;4(4):139-49.
7. Singh MD, Gautam C, Patidar OP, Meena HM, Prakasha G, Vishwajith. Nano-Fertilizers is a new way to increase nutrients use efficiency in crop production a review article. *Int J Agric Sci.* 2017;9(7):3831-3.
8. Rahman IU, Aftab RA, Zafar I, Shafiul. M. Foliar application of plant mineral nutrients on wheat: a review. *RRJAAS.* 2014;3(2):19-22.
9. Corbeels M, Hofman G, Van Cleemput O. Fate of fertilizer N applied to winter wheat growing on a vertisol in a Mediterranean environment. *Nutr Cycl Agro-Ecosyst.* 1999;53(3):249-58.
10. AL-Taey DKA, Al-Janabi ASH, Rachid AM. Effect of water salinity, Organic and minerals fertilization on growth and some nutrients elements in cabbage *Brassica oleracea* var capitata. *Babylon J Pure Appl Sci.* 2017;25(6):232-48.
11. Ziadi N, Bélanger G, Cambouris AN, Tremblay N, Nolin MC, Claessens A. Relationship between phosphorus and nitrogen concentrations in spring wheat. *Agron J.* 2008;100(1):80-6.
12. Liakas V, Rauckis V, Paltanavičius V. Influence of phosphorus and potash fertilizers on germination, tillering and overwintering of winter wheat. *Mokslo Darbai.* 2001;74:3-12.
13. Molla H, Gashaw M, Wassie H. Response of bread wheat to integrated application of vermicompost and NPK fertilizers. *Afr J Agric Res.* 2018;13(1):14-20.
14. Crista F, Isidora R, Florin S, Laura C, Adina B. Influence of NPK fertilizers upon winter wheat grain quality. *Res J Agric Sci.* 2012;44(3):30-5.
15. Rietra RPJJ, Heinen M, Dimkpa CO, Bindaban PS. Effects of nutrient antagonism and synergism on yield and fertilizer use efficiency. *Commun Soil Sci Plant Anal.* 2017;48(16):1895-920.
16. Abdel-Aziz H, Hasaneen MN, Omar A. Effect of foliar application of Nano chitosan NPK fertilizer on the chemical composition of wheat grains. *Egypt J Bot.* 2018;58(1):87-95.
17. Abdel-Aziz HMM, Hasaneen MNA, Omer AM. Nano chitosan-NPK fertilizer enhances the growth and productivity of wheat plants grown in sandy soil. *Span J Agric Res.* 2016;14(1):1-9.
18. Singh U, Praharaj CS, Singh SS, Hazra KK, Kumar N. Up-scaling nutrient energy and system productivity of pigeonpea-wheat cropping system in Indo-Gangetic plains of India. *J Environ Biol.* 2018;39(5):647-58.
19. Barker G. Archaeology: The cost of cultivation. *Nature.* 2011;473(7346):163-4.
20. Gomez KA, Gomez AA. Statistical procedures for agricultural research. John Wiley & Sons; 1984.
21. Ali NS Et al. Juthery H.W.A. The application of nanotechnology for micronutrient in agricultural production (review article). *Iraqi J Agric Sci.* 2017;9(48):489-41.
22. AL-Taey DKA, AL-Azawi SSM, AL-Shareefi MJH, AL-Tawaha AR. Effect of saline water, NPK and organic fertilizers on soil properties and growth, antioxidant enzymes in leaves and yield of lettuce (*Lactuca sativa* var. Parris Island). *Res Crops.* 2018;19:441-9.
23. Masoud B, Abdolshahi R, Nejad GM, Yousefi K, Tabatabaie SM. Effect of different microelement treatment on wheat (*Triticum aestivum* L.) growth and yield. *Int Res J Appl Basic Sci.* 2012;3(1):219-23.
24. Qureshi A, Singh DK, Dwivedi S. Nano-fertilizers: A novel way for enhancing nutrient use efficiency and crop productivity. *Int J Curr Microbiol Appl Sci.* 2018;7(2):3325-35.
25. Dimkpa CO, McLean JE, Britt DW, Anderson AJ. Nano-CuO and interaction with Nano-ZnO or soil bacterium provide evidence for the interference of nanoparticles in metal nutrition of plants. *Ecotoxicology.* 2015;24(1):119-29.

26. Gosavi AB, Deolankar KP, Chaure JS, Gadekar DA. Response of wheat for NPK foliar sprays under water stress condition. Int J Chem Stud. 2017;5(4):766-8.
27. Ajith kumar K, Kumar Y, Savitha AS, Ajayakumar MY, Narayanaswamy C, Raliya R et al. Effect of IFFCO nanofertilizer on growth, grain yield and managing turcicum leaf blight disease in maize. Int J Plant Soil Sci. 2021;33(16): 19-28.
28. Kumar Y, Singh T, Raliya R, Tiwari KN. Nano fertilizers for sustainable crop production, higher nutrient use efficiency and enhanced profitability. Indian J Fertilisers. 2021;17(11):1206-14.

© 2023 Singh et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/98586>