



## **Comparative Evaluation of Anti-diabetic Potential of Aqueous Seed Extracts of *Momordica charantia*, Seed Kernel Extract of *Mangifera indica* and Its Herbal Formulation – an *in vitro* Study**

**Chevuru Sai Shreya Reddy<sup>a</sup>, R. Gayathri<sup>a\*≡</sup>, V. Vishnu Priya<sup>a∞</sup>, J. Selvaraj<sup>a≡</sup> and S. Kavitha<sup>a#</sup>**

<sup>a</sup> Department of Biochemistry, Saveetha Dental College and Hospitals, Saveetha Institute of Medical and Technical Sciences, Saveetha University, Chennai-600 077, India.

### **Authors' contributions**

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

### **Article Information**

DOI: 10.9734/JPRI/2021/v33i62A35518

### **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/80931>

**Original Research Article**

**Received 12 November 2021**

**Accepted 27 December 2021**

**Published 28 December 2021**

## **ABSTRACT**

**Introduction:** A condition in which the body's ability gets impaired to process blood glucose level is called diabetes, otherwise known as blood sugar. It ranked sixteenth among leading causes of death around the globe. There are estimated 72.96 million cases diagnosed with diabetes in the adult population of India. *Momordica charantia* possess anti-diabetic, anti hyperglycemic and anti inflammatory properties. *Mangifera indica* is traditionally used to treat diarrhoea, cancer, diabetes and tooth aches.

**Aim:** Aim of this study is to analyse comparative evaluation of anti-diabetic potential of aqueous seed extracts of *Momordica charantia*, seed kernel extract of *Mangifera indica* through *in vitro* analysis.

**Methods:** *Momordica charantia* and *Mangifera indica* were collected from local farms, and the extracts were analysed for its phytochemicals. Antioxidant and Anti-diabetic potential of the seed

<sup>≡</sup> Associate Professor;

<sup>∞</sup> Professor;

<sup>#</sup> Lecturer;

\*Corresponding author: E-mail: [gayathri.sdc@saveetha.com](mailto:gayathri.sdc@saveetha.com);

extract were evaluated along with standards. The data were analysed statistically by a one way analysis of variance (ANOVA) followed by Duncan's multiple range test was used to see the statistical significance among the groups. The results with the  $p < 0.05$  level were considered to be statistically significant.

**Results:** *Momordica charantia* and *Mangifera indica* and its herbal formulation showed significant antioxidant and anti-diabetic potential.

**Conclusion:** The results showed both the seed and seed kernel extracts of *Momordica charantia* and *Mangifera indica* possessed anti-diabetic activity but the herbal formulation possessed much more significant anti-diabetic activity than the individual extract.

**Keywords:** *Momordica charantia*; *Mangifera indica*; anti-diabetic; in vitro; therapeutic efficacy; innovative technology; novel method.

## 1. INTRODUCTION

A condition in which the body's ability gets impaired to process blood glucose level is called diabetes, otherwise known as blood sugar. Diabetes Mellitus is a major and rapidly increasing health problem in most of the countries around the world, and is an important cause of prolonged ill health and early death. It ranked sixteenth in the cause of global mortality in 1990, for about 5,71,000 deaths [1]. Recent studies of geographical and ethnical influences have shown and concluded that people of Indian origin are highly prone to diabetes [2]. The number of adults affected from diabetes in India is predicted to increase three times from 19.4 million in 1995 to 57.2 million in 2025 [3]. The global diabetic prevalence in 2019 is estimated to be 9.3%. Different types of diabetes can occur, and managing the condition depends on the type. Not all forms of diabetes occur because of an inactive lifestyle. In fact, some are present from childhood. There are 3 types of diabetes classified into type 1, type 2 and gestational diabetes. Type 1 diabetes, also known as juvenile diabetes, occurs when the body fails to produce insulin. People with type 1 diabetes are insulin dependent, in which people must be injected with artificial insulin to stay alive. Type-2 diabetes is affected by the way the body uses insulin. While the body still produces insulin, unlike in Type-1, the cells present in the body do not respond effectively as they once did. This is the most common type of diabetes, according to the National Institute of Diabetes and Digestive and Kidney Diseases, and it is very much related to obesity. Gestational diabetes occurs in women during pregnancy when the body becomes less sensitive towards insulin. Gestational diabetes does not occur in all women but usually continues in women even after giving birth.

Antioxidant activity is the property of a limitation of proteins, lipids, DNA or other oxidation of molecules. Primary antioxidants show effect directly in scavenging free radicals, and secondary antioxidants indirectly cause prevention of free radicals through Fenton's reaction [4]. Free radicals play a very significant role in tissue development in living organisms [5,6]. *Momordica charantia* and *Mangifera indica* possess various antioxidant properties.

Medicinal plants are used as primary health care, as an alternative option to modern synthetic drugs in developing countries [7]. During the past few years *Momordica charantia* has grabbed attention from researchers. *Momordica charantia* also called as "bitter melon" or African cucumber is a plant of *Cucurbitaceae* family widely cultivated in tropical and sub-tropical regions and is commonly used in Mediterranean traditional medicine for its anti-diabetic properties and antihyperglycemic and anti-inflammatory activities. The fruits, stems, leaves and roots of bitter melon are all used as traditional medicine to treat hyperlipidemia, digestive disorders, microbial infections and menstrual problems [8]. The plant possesses traditional usage and modern scientific evidence of benefits of *Momordica charantia*, it is one of the most promising plants for the usage in treating diabetes [9,10].

*Mangifera indica*, commonly known as mango, belongs to the *Anacardiaceae* family. *Mangifera* genus contains 69 species. The bark of the mango tree is traditionally used to treat diarrhea, cancer, diabetes, toothache, skin infections and used as diuretic, antiseptic, hepatoprotective agent [11]. The mango plant shows both invitro and invivo antioxidant properties [12]. Parts of the plant also show hypoglycemic activity [13]. Plant seeds have also shown antibacterial activity [14]. Though there is several work done

on the various parts of the plants, much work has not been documented on the seed and seed kernel extracts of the plants. The aim of the study is to analyze and compare the anti-diabetic potential of aqueous seed extract of *Momordica charantia*, seed kernel extract of *Mangifera indica* and its herbal formulation as an in vitro study.

Even though there are many drugs, the main disadvantage of current drugs is that they have to be given throughout the life and they produce side effects. By exploring the natural herbs to replace the synthetic drugs, prevention can be done naturally without any side effects if significant research is done. Our team has extensive knowledge and research experience that has translate into high quality publications [15-34]. The present study was undertaken to analyze the anti-diabetic effect of *Momordica charantia* and *Mangifera indica*. Alternate hypothesis is that there is an anti-diabetic activity of *Momordica charantia* and seed kernel extract of *Mangifera indica*.

## 2. MATERIALS AND METHODS

### 2.1 Plant Extract

*Momordica charantia* and seed kernel of *Mangifera indica* were collected from a farm in Chennai, washed, crushed and made into powder. Crushed seed powder was utilized to prepare an 80% aqueous extract. Equal volume of the extract was mixed and a formulation was prepared. The extract was prepared by a hot percolation method.

### 2.2 Phytochemical Screening Test

Test for phlobatannin:

1ml of the extract was treated with 1ml of 1% HCl and boiled for 10 mins. The formation of red color precipitate indicates the presence of phlobatannin.

### 2.3 Test for Carbohydrates

Three to five drops of Molisch reagent was added with 1 mL of the extract and then 1 mL of concentrated sulphuric acid was added carefully through the side of the test tube. The mixture was then allowed to stand for two minutes and diluted with 5 mL of distilled water. The development of a red or dull violet ring at the junction of the liquids showed the presence of carbohydrates.

### 2.4 Test for Flavonoids

Few drops of 1% liquid ammonia were taken in a test tube and along with it 1ml of the extract was added resulting in the formation of yellow color thereby indicating the presence of flavonoids.

### 2.5 Test for Alkaloids

2ml of sample was mixed with 2ml of HCl. Then 6 drops of HCN was added and further 2 drops of picric acid was added that resulted in a creamish pale yellow ppt indicating the presence of alkaloids.

### 2.6 Test for Terpenoids

2 ml of sample along with 2ml of chloroform and 3ml of con. H<sub>2</sub>SO<sub>4</sub> was added. Red color ppt obtained indicates the presence of terpenoids.

### 2.7 Test for Proteins

One milliliter of ninhydrin was dissolved in 1 mL of acetone and then a small amount of extract was added with ninhydrin. The formation of purple colour revealed the presence of protein.

### 2.8 Detection of Saponins

Foam test:

A fraction of the extract was vigorously shaken with water and observed for persistent foam.

### 2.9 Test for Steroids

One milliliter of chloroform was mixed with 1 mL of extract and then ten drops of acetic anhydride and five drops of concentrated sulphuric acid were added and mixed. The formation of dark red colour or dark pink colour indicates the presence of steroids [13].

### 2.10 Antioxidant Activity

DPPH free radical scavenging activity:

Scavenging of 2, 2-Diphenyl-1-picrylhydrazyl (DPPH) radical scavenging activity was assessed by the method of Hatano et al<sup>[1989]</sup>. DPPH solution (1.0 ml) was added to 1.0 ml of extract at different concentrations (0.1 to 0.5mg/ml). The mixture was kept at room temperature for 50 minutes and the activity was measured at 517 nm. Ascorbic acid at the same

concentrations was used as standard. The capability to scavenge the DPPH radical scavenging potential was calculated and expressed in percentage (%) using following formula:

$$\text{DPPH radical scavenging (\%)} = \frac{\text{Control OD} - \text{Sample OD}}{\text{Control OD}} \times 100$$

### 2.11 Alpha Amylase Inhibitory Activity of Aqueous Seed Extract of *Momordica charantia*, Seed Kernel Extract of *Mangifera indica*

Alpha amylase inhibitory activity of extract was carried out according to the standard method of ademiluyi et al<sup>[2013]</sup>. In a test tube a reaction mixture containing 500 mu/l phosphate buffer (100mM ; pH=6.8), 100 mu alpha amylase (2 mu/l) and varying concentration of extract (0.1 - 0.5 mg/ml) was incubated at 37° Celsius for 20 minutes. Then the 200 mu/l of 1% soluble starch (100 MM phosphate buffer 6.8) was added as a substrate and incubated further at 37° degree Celsius for 30 minutes. 1000 mu/l of the 3,5 Dinitrosalicylic acid [DNS], DNS colour reagent was then added and boiled for 10 minutes. The absorbance of the resulting mixture was measured at 540 nm using a multi plate reader. Acarbose at various concentrations (0.1-0.5 mg/ml) was used as a standard.

$$\text{Inhibitory activity [\%]} = (1-AS/AC) \times 100$$

AS = Absorbance in the presence of test substance

AC = Absorbance of control

### 2.12 Alpha Glucosidase Inhibitory Activity of Aqueous Seed Extract of *Momordica charantia*, Seed Kernel Extract of *Mangifera indica*

Alpha glucosidase inhibitory activity of extract was carried out according to the method of (35).

Reaction mixture containing 500 mu/l phosphate buffer(100mM pH 6.8), 100mu/l glucosidase (10 ml) and varying concentration of extract (0.1 to 0.5 mg /ml) was pre- incubated at 37 degree Celsius for 15 minutes. Then 200µl of p-NPG(5mM) was added as a substrate and incubated further at 37degree Celsius for 30 minutes. The reaction was stopped by adding 50 mu/l sodium carbonate (0.1M). The absorbance of the released p- nitrophenol was measured at 405 nm using multiple readers. Acarbose at various concentrations (0.1-0.5mg/ml)was used as a standard.

$$\text{Inhibitory activity [\%]} = (1-AS/AC) \times 100$$

AS=absorbance of test substance;

AC= absorbance of control.

### 2.13 Statistical Analysis

The data were subjected to statistical analysis using one-way analysis of variance (ANOVA) and Duncan's multiple range test to assess the significance of individual variations between the groups. In Duncan's test, significance was considered at the level of p<0.05.


## 3. RESULTS AND DISCUSSION

Flavonoides are more concentrated in *Momordica charantia* than in *Mangifera indica*. Alkaloids are equal in concentration in *Momordica charantia* and *Mangifera indica*. Terpenoids are more concentrated in *Mangifera indica* than in *Momordica charantia*. Saponins are equal in concentration in *Momordica charantia* and *Mangifera indica*. Steroids are equal in concentration in *Momordica charantia* and *Mangifera indica*. Carbohydrates are equal in concentration in *Momordica charantia* and *Mangifera indica*.

**Table 1. Phytochemical analysis of aqueous seed extract of *Momordica charantia* and seed kernel extract of *Mangifera indica***


Phytochemicals	<i>Momordica charantia</i>	<i>Mangifera indica</i>
Flavonoides	++	+
Alkaloids	+	+
Terpenoides	+	++
Saponins	+	+
Steroids	+	+
Carbohydrates	-	+

Antioxidant potential of *Momordica charantia*, *Mangifera indica* and its combination




**Fig. 1.** Represents antioxidant potential of aqueous seed extracts of *Momordica charantia*, seed kernel extract of *Mangifera indica* and its herbal formulation against the standard Ascorbic acid- DPPH Assay. X axis represents the concentration in µg/ml and Y axis represents the inhibitory potential of the extracts. Blue bar represents standard Ascorbic acid, orange bar represents aqueous seed extracts of *Momordica charantia* green bar represents seed kernel extract of *Mangifera indica* and yellow bar represents its herbal formulation. Each bar represents Mean ± SEM of 3 independent observations. Significance at  $p \leq 0.05$  X axis represents the concentration in µg/ml and Y axis represents the inhibitory potential of the extracts

αAmylase potential of *Momordica charantia*, *Mangifera indica* and its combination



**Fig. 2.** Represents Alpha Amylase inhibitory potential of aqueous seed extracts of *Momordica charantia*, seed kernel extract of *Mangifera indica* and its herbal formulation against the standard Acarbose. X axis represents the concentration in µg/ml and Y axis represents the inhibitory potential of the extracts. Purple bar represents standard Acarbose, orange bar represents aqueous seed extracts of *Momordica charantia* green bar represents seed kernel extract of *Mangifera indica* and yellow bar represents its herbal formulation. Each bar represents Mean ± SEM of 3 independent observations. Significance at  $p \leq 0.05$



**Fig. 3. Represents Alpha Glucosidase inhibitory potential of aqueous seed extracts of *Momordica charantia*, seed kernel extract of *Mangifera indica* and its herbal formulation against the standard Acarbose. X axis represents the concentration in µg/ml and Y axis represents the inhibitory potential of the extracts. Blue bar represents standard Acarbose, orange bar represents aqueous seed extracts of *Momordica charantia* green bar represents seed kernel extract of *Mangifera indica* and yellow bar represents its herbal formulation. Each bar represents Mean  $\pm$  SEM of 3 independent observations. Significance at  $p \leq 0.05$**

The results revealed a strong presence of phytochemicals such as alkaloid, flavonoids, terpenoids, saponins and steroids in both the seed extract of *Momordica charantia* and seed kernel extract of *Mangifera indica* (Table:1). Phytochemicals are secondary metabolites which are present only among plants. They possess various biologically active compounds that protect and help in normal functioning of the human body. The presence of phytochemicals like alkaloids, terpenoids, saponins, steroids and flavonoids indicate that the extract has potential for further in vitro analysis like antioxidant activity, antidiabetic activity.

Antioxidant activity of aqueous seed extract of *Momordica charantia* and seed kernel of *Mangifera indica* was determined by DPPH free radical scavenging assay. Free radicals / molecules possessing an unpaired electron leads to oxidative stress. Phenolic compounds have great importance in free radical scavenging activity. Similar findings by G. Leela Prakash et al suggest a significant total antioxidant activity possessed by both the aqueous and methanol extract of *Momordica charantia*. Samba Fama Ndoye et al, 2018, has analysed the total antioxidant activity by DPPH radical scavenging activity. The effects of the antioxidants on DPPH free radical scavenging was considered to be due to their hydrogen donating ability. The

results obtained in the study show that aqueous seed extract of *Momordica charantia*, seed kernel extract of *Mangifera indica* and its herbal formulation exhibit significant antioxidant activity with IC<sub>50</sub> of 450, 410 and 350 µg/ml respectively as compared with the standard Vitamin C (Fig. 1). Herbal formulation was found to exhibit a significantly more antioxidant potential than the individual extracts. Further studies may be needed to find out the potential health benefits of the extracts in prevention and scavenging of free radicals.

In the study, *in vitro*  $\alpha$ -amylase inhibitory activity and  $\alpha$ -glucosidase inhibitory activity of aqueous seed extracts of *Momordica charantia*, seed kernel extract of *Mangifera indica* and its formulation were studied. Results revealed a dose dependent increase in % of inhibitory activity. The extracts showed potent anti diabetic activity in a dose dependent manner, with a IC<sub>50</sub> for  $\alpha$ -amylase inhibitory activity was found to be 380, 400, 360 µg/ml respectively (Fig. 2) and IC<sub>50</sub> for  $\alpha$ -glucosidase inhibitory activity was found to be of 420, 400, 320 µg/ml respectively (Fig. 3) and increased in a dose dependent manner as compared to the standard. Acarbose is a standard drug used for treating diabetes, and used for the comparison of anti diabetic activity of the extract. The extracts might be used as starch blocker since it prevents or slows the

absorption of starch into the body, mainly by blocking the hydrolysis of glycosidic linkage of starch. There is a positive relationship between phytoconstituent content and ability to inhibit  $\alpha$ -glucosidase and  $\alpha$ -amylase. Outcome of this study indicates that aqueous extracts of *Momordica charantia*, *Mangifera indica* and its herbal formulation could be used as potential antidiabetic agents. Similar to antioxidant potential, formulation exhibited an increased antidiabetic potential compared to its individual extract.

The increased antioxidant and anti-diabetic potential of the herbal formulation compared to the individual extract can be because of the synergistic effect. More studies need to be done to check the medicinal value of polyherbal formulations. Many indigenous medicines come as a formulation and research is needed to check and document the role of each extract and the synergism shown by the herbal extracts when made into a formulation. Only a proper research on this avenue can help spread awareness on the importance of polyherbal drug preparation and its potential in curing various ailments.

In future, further purification and toxicity studies followed by in vivo studies on the herbal formulation needs to be done. Exploring natural drugs by various research can replace synthetic drugs and its side effects. Prevention can be done even before getting diagnosed with the disease. As there is no significant research done on anti-diabetic properties of seed extract of *Momordica charantia*, seed kernel extract of *Mangifera indica* and its formulation, the current can be a feasible option for further research for drug formulation and diabetic cure.

#### 4. CONCLUSION

From the obtained results, it can be concluded that herbal formulation of aqueous seed extract of *Momordica charantia*, seed kernel extract of *Mangifera indica* possessed a significant antioxidant and antidiabetic potential than the individual extract. In future, more research is needed to throw light on the synergistic role of herbal extracts in a formulation to create awareness on the rich medicinal value of the folk and hereditary medicines.

#### CONSENT

It is not applicable.

#### ETHICAL APPROVAL

It is not applicable.

#### SOURCE OF FUNDING

The present study was supported by the following agencies:

- Saveetha Dental College,
- Saveetha Institute of Medical and Technical Science,
- Saveetha University,
- The Kalpana Indane Gas Agency.

#### ACKNOWLEDGEMENT

The authors would like to thank Saveetha Dental College and Hospitals, SIMATS, Saveetha University for providing laboratory facilities to conduct the study.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. Murray CJL, Lopez AD. Global mortality, disability, and the contribution of risk factors: Global Burden of Disease Study [Internet]. The Lancet. 1997;349: 1436–42. Available: [http://dx.doi.org/10.1016/s0140-6736\(96\)07495-8](http://dx.doi.org/10.1016/s0140-6736(96)07495-8)
2. Shaw JE, Zimmet PZ, de Courten M, Dowse GK, Chitson P, Gareeboo H, et al. Impaired fasting glucose or impaired glucose tolerance. What best predicts future diabetes in Mauritius? [Internet]. Diabetes Care. 1999;22:399–402. Available: <http://dx.doi.org/10.2337/diacare.22.3.399>
3. Madhan M, Gunasekaran S, Arunachalam S. Evaluation of research in India – are we doing it right? [Internet]. Indian Journal of Medical Ethics; 2018. Available: <http://dx.doi.org/10.20529/ijme.2018.024>
4. Huang D, Ou B, Prior RL. The Chemistry behind Antioxidant Capacity Assays [Internet]. Journal of Agricultural and Food Chemistry. 2005;53:1841–56. Available: <http://dx.doi.org/10.1021/jf030723c>

5. Kehrer JP. Free Radicals as Mediators of Tissue Injury and Disease [Internet]. *Critical Reviews in Toxicology*. 199;323: 21–48. Available:<http://dx.doi.org/10.3109/10408449309104073>
6. Free Radicals in Biology and Medicine. Fifth Edition. By Barry Halliwell and John M. C. Gutteridge. Oxford and New York: Oxford University Press. \$115.00 (paper). xxxviii 905 p.; ill.; index. ISBN: 978-0-19-871748-5. 2015 [Internet]. *The Quarterly Review of Biology*. 2016;91: 371–371. Available: <http://dx.doi.org/10.1086/688138>
7. Fachinan R, Yessoufou A, Nekoua MP, Moutairou K. Effectiveness of Antihyperglycemic Effect of *Momordica charantia*: Implication of T-Cell Cytokines [Internet]. *Evidence-Based Complementary and Alternative Medicine*. 2017;2017:1–8. Available:<http://dx.doi.org/10.1155/2017/3707046>
8. Joseph B, Jini D. Antidiabetic effects of *Momordica charantia* (bitter melon) and its medicinal potency [Internet]. *Asian Pacific Journal of Tropical Disease*. 2013;3:93–102. Available: [http://dx.doi.org/10.1016/s2222-1808\(13\)60052-3](http://dx.doi.org/10.1016/s2222-1808(13)60052-3)
9. Cefalu WT, Ye J, Zuberi A, Ribnicky DM, Raskin I, Liu Z, et al. Botanicals and the metabolic syndrome [Internet]. *The American Journal of Clinical Nutrition*. 2008;87:481S – 487S. Available:<http://dx.doi.org/10.1093/ajcn/87.2.481s>
10. Leung L, Birtwhistle R, Kotecha J, Hannah S, Cuthbertson S. Anti-diabetic and hypoglycaemic effects of *Momordica charantia* (bitter melon): a mini review [Internet]. *British Journal of Nutrition*. 2009;102: 1703–8. Available:<http://dx.doi.org/10.1017/s000714509992054>
11. Akram M, Ahmed R, Shakir I, Ibrahim WAW, Hussain R. Extracting hydroxyapatite and its precursors from natural resources [Internet]. *Journal of Materials Science*. 2014;49:1461–75. Available:<http://dx.doi.org/10.1007/s10853-013-7864-x>
12. Nguyen HX, Van Do TN, Le TH, Nguyen MTT, Nguyen NT, Esumi H, et al. Chemical Constituents of *Mangifera indica* and Their Antiausterity Activity against the PANC-1 Human Pancreatic Cancer Cell Line. *J Nat Prod*. 2016; 79(8):2053–9.
13. Phytochemical Analysis, Antibacterial and Antioxidant Activity of the Leave Extracts of *Ruta Chalepensis* [Internet]. *Chemistry and Materials Research*; 2019. Available: <http://dx.doi.org/10.7176/cmr/11-6-01>
14. Saleem A, Saleem M, Akhtar MF, Shahzad M, Jahan S. Correction to: *Moringa oleifera* leaf extracts attenuate Complete Freund’s adjuvant-induced arthritis in Wistar rats via modulation of inflammatory and oxidative stress biomarkers [Internet]. *Inflammopharmacology*. 2020;28:341–3. Available:<http://dx.doi.org/10.1007/s10787-019-00636-y>
15. Wu F, Zhu J, Li G, Wang J, Veeraraghavan VP, Krishna Mohan S, et al. Biologically synthesized green gold nanoparticles from Siberian ginseng induce growth-inhibitory effect on melanoma cells (B16). *Artif Cells Nanomed Biotechnol*. 2019;47(1):3297–305.
16. Chen F, Tang Y, Sun Y, Veeraraghavan VP, Mohan SK, Cui C. 6-shogaol, a active constituents of ginger prevents UVB radiation mediated inflammation and oxidative stress through modulating Nrf2 signaling in human epidermal keratinocytes (HaCaT cells). *J Photochem Photobiol B*. 2019;197:111518.
17. Li Z, Veeraraghavan VP, Mohan SK, Bolla SR, Lakshmanan H, Kumaran S, et al. Apoptotic induction and anti-metastatic activity of eugenol encapsulated chitosan nanopolymer on rat glioma C6 cells via alleviating the MMP signaling pathway [Internet]. *Journal of Photochemistry and Photobiology B: Biology*. 2020;203: 111773. Available:<http://dx.doi.org/10.1016/j.jphoto-biol.2019.111773>
18. Babu S, Jayaraman S. An update on  $\beta$ -sitosterol: A potential herbal nutraceutical for diabetic management. *Biomed Pharmacother*. 2020 ;131:110702.
19. Malaikolundhan H, Mookkan G, Krishnamoorthi G, Matheswaran N, Alsawalha M, Veeraraghavan VP, et al. Anticarcinogenic effect of gold nanoparticles synthesized from *Albizia lebeck* on HCT-116 colon cancer cell lines. *Artif Cells Nanomed Biotechnol*. 2020 ;48(1):1206–13.
20. Han X, Jiang X, Guo L, Wang Y, Veeraraghavan VP, Krishna Mohan S, et



- al. Anticarcinogenic potential of gold nanoparticles synthesized from *Trichosanthes kirilowii* in colon cancer cells through the induction of apoptotic pathway. *Artif Cells Nanomed Biotechnol.* 2019; 47(1):3577–84.
21. Gothai S, Muniandy K, Gnanaraj C, Ibrahim IAA, Shahzad N, Al-Ghamdi SS, et al. Pharmacological insights into antioxidants against colorectal cancer: A detailed review of the possible mechanisms. *Biomed Pharmacother.* 2018;107:1514–22.
  22. Veeraraghavan VP, Hussain S, Balakrishna JP, Dhawale L, Kullappan M, Ambrose JM, et al. A Comprehensive and Critical Review on Ethnopharmacological Importance of Desert Truffles: *Terfezia claveryi*, *Terfezia boudieri*, and *Tirmania nivea* [Internet]. *Food Reviews International.* 2021;1–20. Available:<http://dx.doi.org/10.1080/87559129.2021.1889581>
  23. Sathya S, Ragul V, Veeraraghavan VP, Singh L, Niyas Ahamed MI. An in vitro study on hexavalent chromium [Cr(VI)] remediation using iron oxide nanoparticles based beads. *Environmental Nanotechnology, Monitoring & Management.* 2020;14:100333.
  24. Yang Z, Pu M, Dong X, Ji F, Priya Veeraraghavan V, Yang H. Piperine loaded zinc oxide nanocomposite inhibits the PI3K/AKT/mTOR signaling pathway via attenuating the development of gastric carcinoma: In vitro and in vivo studies. *Arabian Journal of Chemistry.* 2020;13(5): 5501–16.
  25. Rajendran P, Alzahrani AM, Rengarajan T, Veeraraghavan VP, Krishna Mohan S. Consumption of reused vegetable oil intensifies BRCA1 mutations. *Crit Rev Food Sci Nutr.* 2020;1–8.
  26. Barma MD, Muthupandiyani I, Samuel SR, Amaechi BT. Inhibition of *Streptococcus mutans*, antioxidant property and cytotoxicity of novel nano-zinc oxide varnish. *Arch Oral Biol.* 2021;126: 105132.
  27. Samuel SR. Can 5-year-olds sensibly self-report the impact of developmental enamel defects on their quality of life? *Int J Paediatr Dent.* 2021;31(2): 285–6.
  28. Samuel SR, Kuduruthullah S, Khair AMB, Shayeb MA, Elkaseh A, Varma SR. Dental pain, parental SARS-CoV-2 fear and distress on quality of life of 2 to 6 year-old children during COVID-19. *Int J Paediatr Dent.* 2021;31(3): 436–41.
  29. Tang Y, Rajendran P, Veeraraghavan VP, Hussain S, Balakrishna JP, Chinnathambi A, et al. Osteogenic differentiation and mineralization potential of zinc oxide nanoparticles from *Scutellaria baicalensis* on human osteoblast-like MG-63 cells [Internet]. *Materials Science and Engineering: C.* 2021;119: 111656. Available:<http://dx.doi.org/10.1016/j.msec.2020.111656>
  30. Yin Z, Yang Y, Guo T, Veeraraghavan VP, Wang X. Potential chemotherapeutic effect of betalain against human non-small cell lung cancer through PI3K/Akt/mTOR signaling pathway. *Environ Toxicol.* 2021;36(6):1011–20.
  31. Veeraraghavan VP, Periadurai ND, Karunakaran T, Hussain S, Surapaneni KM, Jiao X. Green synthesis of silver nanoparticles from aqueous extract of *Scutellaria barbata* and coating on the cotton fabric for antimicrobial applications and wound healing activity in fibroblast cells (L929). *Saudi J Biol Sci.* 2021;28(7): 3633–40.
  32. Mickymaray S, Alfaiz FA, Paramasivam A, Veeraraghavan VP, Periadurai ND, Surapaneni KM, et al. Rhaponticin suppresses osteosarcoma through the inhibition of PI3K-Akt-mTOR pathway. *Saudi J Biol Sci.* 2021;28(7): 3641–9.
  33. Teja KV, Ramesh S. Is a filled lateral canal – A sign of superiority? [Internet]. *5 Journal of Dental Sciences.* 2020;15:562–3. Available: <http://dx.doi.org/10.1016/j.jds.2020.02.009>
  34. Kadanakuppe S, Hiremath S. Social and Behavioural Factors Associated with Dental Caries Experience among Adolescent School Children in Bengaluru City, India [Internet]. *British Journal of Medicine and Medical Research.* 2016;14:1–10. Available:<http://dx.doi.org/10.9734/bjmmr/2016/24021>

35. Ademiluyi AO, Oboh G, Owoloye TR, Agbebi OJ. Modulatory effects of dietary inclusion of garlic (*Allium sativum*) on gentamycin-induced hepatotoxicity and oxidative stress in rats. Asian J Trop Biomed. 2013;3(6):470–5.

© 2021 Reddy et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*  
<https://www.sdiarticle5.com/review-history/80931>