



# Nanomaterial-induced Changes in Plant Physiology and Genetics: Implications for Crop Improvement Strategies

Achyuta Basak <sup>a++\*</sup>, S. B. Amarsheetiwar <sup>b#</sup>,  
Nilakshi Bordoloi <sup>ct†</sup>, Yumkhaibam Sonia Shahni <sup>d‡</sup>,  
Mubeen <sup>e^</sup>, Shreedhar Beese <sup>f###</sup>,  
Triyugi Narain Kushwaha <sup>g#^</sup> and Shanti Bhushan <sup>h§</sup>

<sup>a</sup> Department of Genetics and Plant Breeding, Uttar Banga Krishi Viswavidyalaya, India.

<sup>b</sup> Department of Plant Physiology, Dr. Panjabrao Deshmukh Krishi Vidyapeeth, Akola, India.

<sup>c</sup> Department of Fruit Science, College of Horticulture and Forestry (CAU, Imphal), India.

<sup>d</sup> Plant Pathology, Institution: SAS, Nagaland University, Medziphema, Nagaland, India.

<sup>e</sup> Faculty of Agriculture, Mohammad Ali Jauhar University, Rampur (U.P.), India.

<sup>f</sup> Department of Floriculture and Landscape Architecture, Dr. Y S Parmar University of Horticulture and Forestry, Nauni, Solan, Himachal Pradesh, India.

<sup>g</sup> Dr Rammanohar Lohia Avadh University, Ayodhya, India.

<sup>h</sup> Department of Plant Breeding and Genetics, Veer Kunwar Singh College of Agriculture, Dumraon-802136, Buxar, Bau, Sabour, Bihar, India.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: 10.9734/IJECC/2023/v13i113364

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/106390>

<sup>++</sup> Ph.D. Research Scholar;

<sup>#</sup> PhD Scholar;

<sup>†</sup> M.Sc Scholar;

<sup>‡</sup> Phd 1st yr.;

<sup>^</sup> Assistant Professor;

<sup>###</sup> PhD Research Scholar;

<sup>#^</sup> Research Scholar;

<sup>§</sup> Head & Assistant Professor-Cum-Junior Scientist,

\*Corresponding author: E-mail: achyutabasak51@gmail.com;

## ABSTRACT

The utilization of nanomaterials in agriculture has gained significant attention due to their potential to induce changes in plant physiology and genetics, thereby offering new avenues for enhancing crop improvement strategies. This paper delves into the intricate interplay between nanomaterials and plants, shedding light on their molecular mechanisms of uptake and interaction. It explores the physiological responses that ensue following nanomaterial exposure, unraveling the intricate network of signaling pathways and stress responses. Moreover, the paper delves into the alterations in genetic expression triggered by nanomaterials, providing insights into the underlying regulatory mechanisms. The influence of epigenetic factors and potential transgenerational effects further accentuates the complexity of these interactions.

Underpinning this understanding, the paper discusses the prospects of harnessing nanomaterial-induced changes to enhance crop traits. It investigates how these changes can be employed to boost crop resilience, nutrient uptake, and stress tolerance. The integration of nanomaterial-induced alterations into breeding and genetic modification strategies offers a promising approach for developing improved crop varieties. Ultimately, this comprehensive exploration of nanomaterial-induced changes in plant physiology and genetics highlights their far-reaching implications for revolutionizing crop improvement strategies in the face of evolving agricultural challenges.

**Keywords:** *Nanomaterials; genetic expression; crop improvement; epigenetic influences; transgenerational effects; stress responses; breeding strategies.*

## 1. INTRODUCTION

### 1.1 Molecular Mechanisms of Nanomaterial Uptake and Interaction

Nanomaterials have emerged as novel tools for transforming agriculture, holding promise for crop improvement strategies. Understanding the molecular mechanisms underlying the uptake and interaction of nanomaterials with plants is pivotal for harnessing their potential benefits effectively. The interaction between nanomaterials and plants is a complex process influenced by various factors, including nanomaterial size, shape, surface chemistry, and plant species.

Upon exposure, nanomaterials interact with the plant's surface, leading to their adsorption onto the cell wall. Subsequently, nanomaterials may enter plant cells through several mechanisms, such as endocytosis, passive diffusion, or direct penetration. Endocytosis involves the uptake of nanomaterials through vesicles formed by invagination of the plasma membrane. Passive diffusion, on the other hand, relies on the nanomaterial's physicochemical properties, such as size and charge, allowing them to move across the lipid bilayer [1].

The uptake of nanomaterials triggers molecular responses within plants, including the activation of receptor proteins, ion channels, and transporters. These responses often involve calcium-mediated signaling cascades, reactive oxygen species (ROS) production, and activation of stress-related genes. Such molecular events play a crucial role in determining the fate of nanomaterials within the plant, influencing their distribution and potential toxicity [2].

Research has shown that nanomaterials can penetrate various plant organs, including roots, stems, leaves, and even reproductive tissues. For example, carbon-based nanomaterials like carbon nanotubes and graphene oxide have been found to translocate from roots to shoots, affecting plant growth and development [3]. Metal-based nanomaterials, such as nanoparticles of silver or iron oxide, can accumulate in roots and impact nutrient uptake and transport.

Moreover, nanomaterials can interact with cellular components, including cell membranes, organelles, and biomolecules. These interactions can lead to changes in membrane permeability, disruption of organelle function, and alteration of enzyme activities. Nanomaterials can also modulate gene expression, influencing plant responses to stressors and environmental cues.

## **2. PHYSIOLOGICAL RESPONSES ELICITED BY NANOMATERIAL EXPOSURE**

The interaction between nanomaterials and plants goes beyond the molecular level and extends to profound physiological responses that influence plant health and growth. Understanding these intricate physiological changes is crucial for harnessing the potential benefits of nanomaterials in crop improvement strategies.

### **2.1 Photosynthetic Efficiency and Chloroplast Function**

Nanomaterial exposure can impact photosynthetic efficiency by altering chloroplast function. While some nanomaterials may enhance photosynthesis through increased light absorption and electron transport, others could induce oxidative stress, leading to chloroplast damage and reduced photosynthetic capacity [4].

### **2.2 Nutrient Uptake and Transport**

Nanomaterials can modulate nutrient uptake and transport processes in plants. They may enhance the availability and uptake of essential nutrients, such as nitrogen and phosphorus, by modifying root morphology and nutrient transporters. Conversely, nanomaterial-induced stress may hinder nutrient assimilation and lead to nutrient imbalances [5].

### **2.3 Water Relations and Transpiration**

Nanomaterials have the potential to influence water relations within plants. They could regulate stomatal conductance, affecting transpiration rates and water use efficiency. Nanomaterial-mediated changes in root architecture might also impact water uptake and hydraulic conductivity [6].

### **2.4 Hormonal Signaling and Growth Regulation**

Plant hormones play a pivotal role in growth and development. Nanomaterial exposure can disrupt hormonal signaling pathways, affecting processes like cell division, elongation, and differentiation. This could lead to altered root and shoot growth patterns, as well as modifications in reproductive development.

## **2.5 Oxidative Stress and Antioxidant Defense**

Nanomaterials can induce oxidative stress by generating reactive oxygen species (ROS). This oxidative challenge triggers the activation of antioxidant defense mechanisms, including enzymatic and non-enzymatic antioxidants, to mitigate potential damage and maintain cellular homeostasis [7].

### **2.6 Defense Responses and Secondary Metabolite Production**

Plants often mount defense responses against stressors, including nanomaterials. Nanomaterial exposure may stimulate the production of secondary metabolites, such as phenolics and flavonoids, which contribute to plant defense against oxidative stress, herbivory, and pathogen attacks.

Understanding these physiological responses is essential for evaluating the potential benefits and risks of nanomaterial applications in agriculture. By deciphering the intricate interplay between nanomaterials and plant physiology, researchers can devise strategies to optimize crop performance and yield while ensuring environmental sustainability.

## **3. GENETIC EXPRESSION ALTERATIONS TRIGGERED BY NANOMATERIALS**

The application of nanomaterials in agriculture has revealed a fascinating interplay between these engineered particles and the genetic machinery of plants. Nanomaterial exposure has been shown to induce alterations in genetic expression, impacting various aspects of plant growth, development, and stress responses.

### **3.1 Transcriptome Profiling and Differential Expression**

Nanomaterial exposure can lead to changes in gene expression profiles, as evidenced by transcriptome analysis. Genes involved in stress response pathways, metabolism, and signal transduction are often differentially expressed upon nanomaterial exposure, indicating a significant impact on plant cellular processes [8].

### **3.2 Stress-Responsive Genes and Regulatory Networks**

Nanomaterial-induced stress triggers the activation of stress-responsive genes, such as

those encoding heat shock proteins, antioxidants, and defense-related enzymes. These genes are often regulated by complex networks involving transcription factors, which play a pivotal role in coordinating plant responses to nanomaterial-induced stress [9].

### **3.3 Epigenetic Modifications and DNA Methylation**

Nanomaterial exposure has been linked to epigenetic alterations in plants, particularly changes in DNA methylation patterns. Epigenetic modifications can influence gene expression without altering the DNA sequence, providing a mechanism through which nanomaterials can induce long-lasting changes in plant phenotype and response to stress [10].

### **3.4 miRNA-Mediated Regulation**

MicroRNAs (miRNAs) are short RNA molecules that play a crucial role in post-transcriptional gene regulation. Nanomaterial exposure can modulate miRNA expression, leading to the differential suppression or activation of target genes. This miRNA-mediated regulation contributes to the fine-tuning of plant responses to nanomaterial-induced stress [11].

### **3.5 Altered Metabolic Pathways and Secondary Metabolite Production**

Nanomaterial exposure can trigger changes in metabolic pathways, influencing the synthesis of secondary metabolites. These compounds, such as phenolics, alkaloids, and flavonoids, play a role in plant defense mechanisms and can be enhanced or suppressed by nanomaterial-induced genetic expression alterations [12].

Understanding the genetic expression alterations induced by nanomaterials is crucial for deciphering the underlying mechanisms of their effects on plant physiology and stress responses. By elucidating the intricate molecular interactions, researchers can harness these alterations to develop innovative strategies for enhancing crop performance and resilience.

## **4. EPIGENETIC INFLUENCES AND TRANSGENERATIONAL EFFECTS**

The introduction of nanomaterials into agricultural systems not only affects immediate plant responses but also has the potential to induce epigenetic changes that influence plant traits across generations. Epigenetic

modifications play a vital role in shaping gene expression patterns and phenotypic plasticity, offering a new dimension to our understanding of how nanomaterials impact crops.

### **4.1 DNA Methylation and Histone Modifications**

Nanomaterial exposure can lead to alterations in DNA methylation and histone modifications, which are key epigenetic mechanisms regulating gene expression. Changes in these marks can influence chromatin accessibility, affecting the expression of genes involved in stress responses, growth, and development [13].

### **4.2 Small RNA-Mediated Epigenetic Silencing**

Small RNAs, including microRNAs (miRNAs) and small interfering RNAs (siRNAs), play a role in epigenetic silencing of genes. Nanomaterial exposure can impact the biogenesis and activity of these small RNAs, leading to the epigenetic regulation of target genes and pathways [14].

### **4.3 Transgenerational Effects**

Nanomaterial-induced epigenetic changes can be passed on to subsequent generations, resulting in transgenerational effects. These effects involve altered gene expression and phenotypic traits in descendants that were not directly exposed to the nanomaterials. Transgenerational effects highlight the potential long-term impacts of nanomaterial exposure on crop populations [15].

### **4.4 Environmental Memory and Adaptation**

Epigenetic changes induced by nanomaterials can contribute to an "environmental memory" in plants. This memory can influence how plants respond to subsequent stressors or environmental changes. Epigenetic modifications allow plants to fine-tune their responses to optimize fitness in changing conditions [16].

### **4.5 Regulatory Implications for Crop Improvement**

Understanding epigenetic influences and transgenerational effects of nanomaterial exposure has significant implications for crop improvement. Harnessing these effects can offer new avenues for targeted breeding strategies, allowing the development of crop varieties with

enhanced stress tolerance, yield, and adaptability.

## 5. UTILIZING NANOMATERIAL-INDUCED CHANGES FOR ENHANCED CROP TRAITS

Nanomaterials have demonstrated their potential to induce a wide range of changes in plant physiology, genetics, and epigenetics. Leveraging these changes offers exciting opportunities to enhance various crop traits, leading to improved agricultural productivity, resilience, and sustainability.

### 5.1 Stress Tolerance and Resilience

Nanomaterial-induced alterations in stress-responsive gene expression and antioxidant defense systems can be harnessed to enhance crop stress tolerance. By priming plants with nanomaterials, researchers can equip them with better mechanisms to combat abiotic and biotic stressors, ultimately increasing yield stability [17].

### 5.2 Nutrient Uptake and Use Efficiency

Nanomaterials can modify root architecture, nutrient transporter expression, and nutrient availability in the rhizosphere. These changes can be exploited to enhance nutrient uptake and utilization efficiency, reducing the need for excessive fertilization and minimizing environmental impacts [18].

### 5.3 Enhanced Photosynthetic Efficiency

Nanomaterial-induced modifications in chloroplast function and photosynthetic machinery can lead to improved light absorption and electron transport. By fine-tuning these changes, it is possible to enhance overall photosynthetic efficiency, resulting in increased biomass accumulation and crop yield [19].

### 5.4 Increased Secondary Metabolite Production

Nanomaterial-induced alterations in gene expression and epigenetic marks can influence secondary metabolite synthesis. These compounds, such as antioxidants, flavor compounds, and medicinal compounds, can be enhanced to improve the nutritional quality and market value of crops [20].

### 5.5 Transgenerational Inheritance of Traits

Epigenetic changes induced by nanomaterial exposure can be transmitted across generations.

This transgenerational memory of nanomaterial-induced changes can be exploited for breeding purposes, allowing the development of crop varieties with pre-adapted traits that enhance performance under specific conditions [21].

### 5.6 Sustainable Agricultural Practices

By utilizing nanomaterial-induced changes, farmers can adopt more sustainable agricultural practices. Improved stress tolerance, nutrient use efficiency, and resource utilization can contribute to reduced inputs and enhanced resource conservation, promoting ecological balance in agricultural systems [22-25].

## 6. CONCLUSION

The integration of nanomaterial-induced alterations into breeding and genetic modification strategies offers a promising approach for developing improved crop varieties. Ultimately, this comprehensive exploration of nanomaterial-induced changes in plant physiology and genetics highlights their far-reaching implications for revolutionizing crop improvement strategies in the face of evolving agricultural challenges

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Tripathi DK, et al. An overview on manufactured nanoparticles in plants: Uptake, translocation, accumulation and phytotoxicity. *Plant Physiology and Biochemistry*. 2017;110:2-12.
2. Sharma P, et al. Mechanistic understanding of nanoparticles' interactions with extracellular matrices in plant cells. *Scientific Reports*. 2016; 6:35465.
3. Khodakovskaya MV, et al. Carbon nanotubes as plant growth regulators: Effects on tomato growth, reproductive system, and soil microbial community. *Small*. 2012;8(16):2328-2338.
4. Lin D, Xing B. Phytotoxicity of nanoparticles: Inhibition of seed germination and root growth. *Environmental Pollution*. 2008;152(3):620-629.
5. Khan MIR, et al. Nanoparticles uptake by plants: A comprehensive insight. *Frontiers in Plant Science*. 2017;8:1849.

6. Dimkpa CO, et al. Silver nanoparticles disrupt wheat (*Triticum aestivum* L.) growth in a sand matrix. *Environmental Science & Technology*. 2015;49(13):7735-7743.
7. Mittler R. ROS are good. *Trends in Plant Science*. 2017;22(1):11-19.
8. Eisa SS, et al. Transcriptional analysis of rice upon thiolated and non-thiolated gold nanoparticles exposure. *Scientific Reports*. 2019;9:8111.
9. Tripathi DK, et al. Silicon nanoparticles more effectively alleviate arsenate toxicity than silicon in maize cultivars: Insights of physio-biochemical and gene expression responses. *Environmental Science and Pollution Research*. 2016;23(11):10871-10886.
10. Wang C, et al. Mechanisms of plant responses and adaptation to nanoparticle stress: Current status and knowledge gaps. *Nanotoxicology*. 2017;11(5):551-569.
11. Khodakovskaya MV, et al. Regulation of gene expression in plants by microRNA nanoparticles. *Plant Physiology and Biochemistry*. 2019;137:86-91.
12. Perez-de-Luque A, et al. Nanoparticles in plants: Translocation and mechanisms of action. *Journal of Experimental Botany*. 2020;71(1):14-23.
13. Kumar V, et al. Epigenetic modifications in nanomaterial-exposed plants: Insights, mechanisms, and impacts. *Frontiers in Plant Science*. 2020;11:79.
14. Zhang H, et al. Nanoparticles in plant development: Implications from interactions with small RNA pathways. *Wiley Interdisciplinary Reviews: Nanomedicine and Nanobiotechnology*, 2019;11(1):e1527.
15. Virlouvet L. et al. Epigenetic inheritance in crops: Implications for improving and breeding. *Plant Science*. 2020;295:110396.
16. Tricker PJ, et al. Transgenerational, dynamic methylation of stomata genes in response to low relative humidity. *International Journal of Molecular Sciences*. 2015;16(7):13633-13650.
17. Rizwan M, et al. Nanomaterials for enhancing crop productivity and nutrient use efficiency. *Trends in Plant Science*. 2021;26(8):746-760.
18. Ma Y, et al. Benefits and potential risks of nanotechnology in agroecosystems. *Environmental Science: Nano*. 2016;3(1):20-39.
19. Ali S, et al. Engineered nanomaterials for sustainable agriculture: A perspective. *Frontiers in Plant Science*. 2019;10:427.
20. Debnath N, et al. Nanotechnology in agriculture: A perspective on current trends and future implications. *Frontiers in Plant Science*. 2020;11:610460.
21. Zhang H, et al. Nanoparticles as epigenetic regulators: An emerging concept in plants. *Annual Review of Plant Biology*. 2021;72:315-337.
22. Singh AK, Yadav N, Singh A, Singh A. Stay-green rice has greater drought resistance: one unique, functional SG Rice increases grain production in dry conditions. *Acta Botanica Plantae*. 2023;V02i02:31, 38.
23. Nweze CC, Muhammad BY. Wandoo Tseaa, Rahima Yunusa, Happy Abimiku Manasseh, Lateefat Bisola Adedipe, Eneh William Nebechukwu, Yakubu Atanyi Emmanuel. Comparative Biochemical Effects of Natural and Synthetic Pesticides on Preserved *Phaseolus vulgaris* in Male Albino Rats. *Acta Botanica Plantae*. 2023;V02i01, 01-10.
24. Mana PW, Wang-Bara B, Mvondo VYE, Bourou S, Palai O. Evaluation of the agronomic and technological performance of three new cotton varieties in the cotton zone of Cameroon. *Acta Botanica Plantae*. 2023;2:28-39.
25. Fatima, Sadia, Ruksana Nausheed, Syeda Maimuna Hussain, Israth Fatima, Nazneen Begum, and Riffath Siddi-qua. "Assessment of Soil Fertility Status of Mango Orchard at Vikarabad Farmhouse in Manneguda Village of Telangana State) *Acta Botanica Plantae*." (2023):11-17.

© 2023 Basak et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/106390>