



# Influence of Weather Parameters on the Incidence and Severity of Guava Scab Caused by *Pestalotiopsis psidii* (Pat.)

Sirisha M. <sup>a</sup>, Aswathanarayana D. S. <sup>a\*</sup>, Yenjerappa S. T. <sup>a</sup>,  
Gowdar S. B. <sup>a</sup>, Pampanna Y. <sup>b</sup> and Udaykumar Nidoni <sup>c</sup>

<sup>a</sup> Department of Plant Pathology, College of Agriculture, UAS, Raichur, India.

<sup>b</sup> Department of Horticulture, MARS, UAS, Raichur, India.

<sup>c</sup> Department of Processing and Food Engineering, CAE, Raichur, India.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: 10.9734/IJECC/2023/v13i113646

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/110043>

Original Research Article

Received: 02/10/2023

Accepted: 06/12/2023

Published: 11/12/2023

## ABSTRACT

Guava (*Psidium guajava*) is an important tropical fruit crop of India and is known as "apple of the tropics". It is grown and utilized as an important fruit in tropical countries like India, Indonesia, Pakistan, Bangladesh and South America. Among the biotic and abiotic diseases of guava, guava infected by scab disease [*Pestalotiopsis psidii* (Pat.) Mordue] is an economically important and reported average yield losses in the range of 12-18%. An experiment was conducted during 2023 at Horticulture garden, Main Agricultural Research Station, University of Agricultural Sciences, Raichur during 2023 to understand the influence of various weather parameters on the guava scab disease development. The results revealed that, irrespective of the varieties, interaction among maximum temperature, minimum temperature, maximum relative humidity, minimum relative humidity, rainy

\*Corresponding author: E-mail: ashanpathos@gmail.com;

days and rainfall showed significant positive correlation with guava scab disease. And the incidence and severity of guava scab was found to be 23.21, 56.00 (Allahabad Safed), 32.45, 68.50 (Lucknow 49) and 85.00, 74.51 (Arka Kiran) respectively.

**Keywords:** Guava; epidemiology; temperature; relative humidity; rainfall; *Pestalotiopsis psidii*.

## 1. INTRODUCTION

Guava (*Psidium guajava*) is an important fruit crop of India that belongs to the genus *Psidium* and family Myrtaceae [1]. The crop was originated from Tropical America or the Mexico [2] or the Brazil. It is known as “apple of the tropics”. Globally, the production of guava is 55 million tonnes [3]. Guava has been grown and utilized as an important fruit in tropical countries like India, Indonesia, Pakistan, Bangladesh and South America [4]. The total area under guava cultivation in India is around 358.82 thousand hectares with annual production of 5.59 million tonnes. In Karnataka, area under guava cultivation is 8.10 thousand ha with production of 0.16 mt [5]. It has gained considerable prominence an account of its high nutritive value, availability at moderate prices, pleasant aroma and good flavor. It is the fifth most widely grown fruit crop of India [4]. Several biotic and abiotic factors affect the growth and yield of guava. Among the biotic factors, diseases such as *Fusarium* wilt, anthracnose seedling damping off and fruit rot, algal leaf and fruit spot stem canker or bark canker Leaf blight are the most important causing major losses in the guava. Scabby fruit canker caused by *Pestalotiopsis* spp. is one of the most common fruit diseases in guava growing areas and appears at all developmental stages of guava fruit. Scaby canker caused by *Pestalotiopsis* spp. was reported in India and cause drastic reduction of qualitative and quantitative yield (20-25 %) in the fields during the pre-harvest stage and can also lead to fruit losses during post-harvest storage [6]. A study was undertaken to observe impact of weather conditions on the development of scab disease of guava. An understanding of the role of environmental factors and their consequences on epidemiology, infection, development and spread of the pathogen or diseases are required to develop sustainable disease management practices. Keeping in view all of the above points, present investigation was planned and conducted to study the effect of various weather parameters on disease incidence and severity of

guava scab at Horticulture garden, Raichur during 2022-23.

## 2. MATERIALS AND METHODS

To study the effect of different weather parameters on the development of guava scab disease, an experiment was conducted during 2022-23 in the Horticulture garden, Main agricultural research station, University of Agricultural Sciences, Raichur. Ten guava plants, each from three varieties of guava (Allahabad Safed, Lucknow 49 and Arka Kiran) were selected randomly and tagged for future observation in the orchard of Department of Horticulture. From each plant, four branches were selected and in each branch five fruits were selected and tagged. Disease development was monitored at weekly intervals and disease severity was recorded using the 1-4 scale given by Keith et al [6] and per cent disease index (PDI) was calculated. The data of Standard Meteorological week pertaining to maximum and minimum temperature, relative humidity (maximum and minimum), amount of rainfall, number of rainy days and sunshine hours were collected from Meteorological Division, Main Agricultural Research Station, (MARS), Raichur. Correlation co-efficient between disease incidence or disease severity in relation to different weather parameters were determined by Karl Pearson’s formula and tested individually for their significance at 5 per cent probability level using following formula. At the end of the experiment weather parameters which are favourable for disease development can be known.

$$t = \frac{r\sqrt{n-2}}{\sqrt{1-r^2}}$$

Where,

t = test of significance  
r = correlation co-efficient  
n = number of observations

**Chart 1. Disease severity scale for guava scab given by Keith et al [6]**

Grade	Description
1	0 to 25 per cent area of canopy showing typical symptoms of scabby canker
2	26 to 50 per cent area of canopy showing typical symptoms of scabby canker
3	51 to 75 per cent area of canopy showing typical symptoms of scabby canker
4	More than 76 per cent area of canopy showing typical symptoms of scabby canker

**Disease incidence was also calculated using the formula given below**

$$\text{Disease Incidence (\%)} = \frac{\text{Number of fruits infected}}{\text{Total number of fruits observed}} \times 100$$

**Disease severity was calculated using the formula given below**

$$\text{Disease severity (\%)} = \frac{\text{Sum of the individual disease ratings}}{\text{Number of plants / fruits observed}} \times \frac{100}{\text{Maximum disease grade}}$$

The rate of disease development/ unit / day was estimated according to the method given by Van [7]. The apparent infection rate (r) for total period was calculated by using following formula,

$$r = \frac{1}{t_2 - t_1} \log_e \frac{X_2 (1 - X_1)}{X_1 (1 - X_2)}$$

Where,

- r = rate of disease development
- t<sub>1</sub> = date of first observation
- t<sub>2</sub> = date of second observation
- X<sub>1</sub> = disease severity on first observation
- X<sub>2</sub> = disease severity on second observation

### 3. RESULTS AND DISCUSSION

#### 3.1 Influence of Weather Parameters on Guava Scab Disease Severity on Allahabad Safed Variety

The initial guava scab symptoms appeared on 49<sup>th</sup> standard meteorological on the variety Allahabad Safed (2.50 % PDI) and incidence of 1.76 per cent gradually increased week by week until the end of fruit bearing season. Maximum severity (56.00 %) and incidence (23.21 %) was recorded in the first week of March (10<sup>th</sup> SMW) 2023 and the minimum severity and incidence (Table 1 & Fig. 1). Correlation coefficient analysis presented in Table 3 revealed that, as individual parameters maximum temperature (0.587) and sunshine hours (0.753) has positive significant

correlation with guava scab disease development. Whereas, minimum temperature (0.023) was non-significant with positive effect. Rainfall (- 0.377), rainy day (- 0.377), maximum relative humidity (- 0.482) were non-significant and had negative effect with disease. As the temperature increases, relative humidity reduces and that weather condition may not be suitable for the disease development, hence it showed negative correlation. Whereas interaction of sunshine hours with minimum temperature (- 0.419), rainfall (- 0.646), rainy day (- 0.679) and minimum relative humidity (- 0.853) have showed negative significant correlation with the disease development of guava scab. Because of increase in sunshine hours quick evaporation of morning dew and decrease in relative humidity were not favourable for disease development. The multiple linear regression of PDI of guava scab (Allahabad Safed variety) in relation to weather parameters during 2022-23 indicated that, the regression coefficients for maximum temperature (X<sub>1</sub>), minimum temperature (X<sub>2</sub>), rainfall (X<sub>3</sub>), rainy day (X<sub>4</sub>), morning relative humidity (X<sub>5</sub>), evening relative humidity (X<sub>6</sub>) and sunshine hours (X<sub>7</sub>) were found to be 0.045, 5.563, 0.060, 0.060, - 0.133, - 0.216 and 9.481 respectively. The multiple linear regression equation was fitted to the data and the equation arrived for the weather parameters is  $Y = 122.870 + 0.045 X_1 + 5.563X_2 + 0.060 X_3 + 0.060 X_4 - 0.133 X_5 - 0.216 X_6 + 9.481X_7$  (Table 4). This analysis showed that, when there was increase in one unit of maximum temperature, minimum temperature, rainfall, rainy day and sunshine hours, the per cent disease severity

was also increased by 0.045, 5.563, 0.060, 0.060 and 9.481 units. Whereas increase in one unit of morning relative humidity and evening relative humidity, the per cent disease severity was decreased by 0.133 and 0.216 units, respectively.

### 3.2 Influence of Weather Parameters on Guava Scab Disease Severity on Lucknow 49 Variety

In Lucknow 49 variety, the first onset of symptom was noticed during 19<sup>th</sup> November to 25<sup>th</sup> November with the disease severity of 3.50 and incidence of 1.21 (47<sup>th</sup> SMW). Over a period of time, per cent disease index (PDI) reached peak of 68.50 and incidence of 32.45 per cent at 8<sup>th</sup> SMW (3<sup>rd</sup> week of February) due to congenial weather conditions (Table 1 and Fig. 2). The results revealed that, the disease development has positive correlation co-efficient with individual parameters such as, maximum temperature (0.533) and sunshine hours (0.795) but has significant negative correlation co-efficient with minimum relative humidity (- 0.707) (Table 5). Among the interaction between different weather parameters, the positive significant correlation was recorded by maximum temperature and sunshine hours (0.624), rainfall and rainy days (1.000), minimum relative humidity and rainfall (0.691), minimum relative humidity and rainy days (0.680) and minimum relative humidity and

sunshine hours (0.904). The interaction of maximum temperature with rainfall (- 0.565), rainy days (- 0.551), maximum relative humidity (- 0.520) and minimum relative humidity (- 0.602) sunshine hours with rainfall (- 0.567) and rainy day (- 0.556) showed significant negative correlation with guava scab disease incidence. The multiple linear regression of PDI of guava scab on Lucknow 49 variety in relation to weather parameters during 2022-23 indicated that, the regression coefficients for maximum temperature (X<sub>1</sub>), minimum temperature (X<sub>2</sub>), rainfall (X<sub>3</sub>), rainy day (X<sub>4</sub>), morning relative humidity (X<sub>5</sub>), evening relative humidity (X<sub>6</sub>) and sunshine hours (X<sub>8</sub>) were found to be 0.016, 0.273, 0.313, 0.312, 0.117, - 0.068 and 10.505 respectively. The multiple linear regression equation was fitted to the data and the equation arrived for the weather parameters is  $Y = 24.918 + 0.016 X_1 + 0.273X_2 + 0.313 X_3 + 0.312X_4 + 0.117X_5 - 0.068 X_6 + 10.505X_7$  (Table 6). This analysis showed that, when there was increase in one unit of maximum temperature, minimum temperature, rainfall, rainy day, morning relative humidity and sunshine hours, the per cent disease severity was increased by 0.016, 0.273, 0.313, 0.312, 0.117 and 10.505 units respectively. Whereas increase in one unit of evening relative humidity, the per cent disease severity of guava scab was decreased by 0.068 units.




Fig. 1. Progression of guava scab in relation to weather parameters on Allahabad Safed variety

**Table 1. Progression of guava scab in relation to weather parameters on different varieties of guava**

Sl. No.	SMW	Date	Temp. (°C)		RH (%)		Rainfall (mm)	Rainy days	Sunshine hours	Allahabad Safed		Lucknow 49		Arka Kiran	
			Max.	Min.	Max.	Min.				Severity (%)	Incidence (%)	Severity (%)	Incidence (%)	Severity (%)	Incidence (%)
1.	45	05.11.2022 -11.11.2022	30.97	19.86	83.86	38.14	0.00	0.00	7.10	-	-	0.00	0.00	-	-
2.	46	12.11.2022 - 18.11.2022	30.60	17.43	78.29	40.00	0.00	0.00	5.13	-	-	0.00	0.00	-	-
3.	47	19.11.2022- 25.11.2022	29.20	17.63	78.43	47.43	0.06	0.00	3.89	0.00	0.00	3.50	1.21	-	-
4.	48	26.11.2022 - 02.12.2022	31.00	17.80	72.71	47.14	0.00	0.00	5.23	0.00	0.00	10.00	4.82	-	-
5.	49	03.12.2022 - 09.12.2022	30.74	17.14	71.14	37.57	0.00	0.00	4.16	2.50	1.76	15.50	8.01	-	-
6.	50	10.12.2022- 16.12.2022	28.71	18.94	85.00	59.29	1.91	0.43	2.31	6.50	3.09	19.50	10.12	0.00	0.00
7.	51	17.12.2022 - 23.12.2022	30.97	15.23	84.57	35.86	0.00	0.00	6.06	11.50	3.98	27.50	14.34	0.00	0.00
8.	52	24.12.2023-31-12-2023	31.55	18.20	83.63	41.75	0.00	0.00	6.26	22.50	12.38	35.50	19.83	0.00	0.00
9.	1	01.01.2023 – 07.01.2023	30.66	18.86	82.00	41.00	0.00	0.00	4.87	35.50	17.25	47.00	24.47	0.00	0.00
10.	2	08.01.2023 -14.01.2023	30.31	13.17	74.00	25.29	0.00	0.00	8.39	37.00	18.14	55.00	27.00	0.00	0.00
11.	3	15.01.2023-21.01.2023	31.06	16.00	75.57	29.43	0.00	0.00	9.24	40.00	18.14	61.50	28.27	0.00	0.00
12.	4	22.01.2023 – 28.01.2023	30.43	17.49	77.71	31.14	0.00	0.00	7.94	44.00	19.46	65.50	29.53	4.50	2.91
13.	5	29.01.2023 – 04.02.2023	31.34	17.17	76.29	35.00	0.00	0.00	6.81	47.50	19.75	66.00	30.37	10.50	12.75
14.	6	05. 02.2023-11. 02.2023	33.49	18.31	68.29	29.29	0.00	0.00	8.26	51.50	20.21	66.00	31.25	34.50	26.31
15.	7	12. 02.2023-18.02.2023	31.26	17.03	76.78	33.27	0.00	0.00	7.40	52.50	20.21	67.50	32.00	55.50	46.66
16.	8	19.02.2023 – 25.02.2023	31.22	16.86	75.81	32.06	0.00	0.00	7.56	55.00	21.00	68.50	32.45	63.97	67.00
17.	9	26.02.2023 – 04.03.2023	34.60	18.09	53.14	21.71	0.00	0.00	8.01	56.00	22.64	3.50	1.21	68.21	79.00
18.	10	05.03.2023-11.03.2023	34.66	18.54	51.00	23.14	0.00	0.00	7.37	56.00	23.21	10.00	4.82	69.98	81.00
19.	11	12.03.2023-18.03.2023	34.89	20.40	66.43	31.29	0.43	0.14	5.54	-	-	-	-	72.00	81.00
20.	12	19.03.2023-25.03.2023	34.40	19.63	64.71	24.43	1.34	0.14	7.83	-	-	-	-	74.51	85.00




Fig. 2. Progression of guava scab in relation to weather parameters on Lucknow 49 variety

Table 2. Correlation coefficient between weather parameters with guava scab on Allahabad Safed, Lucknow 49 and Arka Kiran

Parameters	Correlation coefficient ( r ) of Allahabad safed	Correlation coefficient ( r ) of Lucknow 49	Correlation coefficient ( r ) of Arka Kiran
X <sub>1</sub> Maximum temperature (°C)	0.045	0.016	12.233
X <sub>2</sub> Minimum temperature (°C)	5.583	0.273	0.096
X <sub>3</sub> Rainfall (mm)	0.060	0.313	0.127
X <sub>4</sub> Rainy days	0.060	0.312	0.114
X <sub>5</sub> Relative humidity (max) (%)	-0.133	0.117	0.038
X <sub>6</sub> Relative humidity (min) (%)	-0.216	-0.068	0.062
X <sub>7</sub> Sunshine hours (hrs/day)	9.481	10.505	-0.215

Table 3. Correlation coefficient between weather parameters with guava scab on Allahabad Safed variety

Parameters	Y	X <sub>1</sub>	X <sub>2</sub>	X <sub>3</sub>	X <sub>4</sub>	X <sub>5</sub>	X <sub>6</sub>	X <sub>7</sub>
Y PDI	1.000							
X <sub>1</sub> Maximum temperature (°C)	0.587*	1.000						
X <sub>2</sub> Minimum temperature (°C)	0.023	0.261	1.000					
X <sub>3</sub> Rainfall (mm)	- 0.377	- 0.475	0.305	1.000				
X <sub>4</sub> Rainy days	- 0.377	- 0.484	0.305	1.000**	1.000			
X <sub>5</sub> Relative humidity (max) (%)	- 0.482	- 0.858**	- 0.171	0.306	0.306	1.000		
X <sub>6</sub> Relative humidity (min) (%)	- 0.747**	- 0.693**	0.367	0.694**	0.694**	0.665**	1.000	
X <sub>7</sub> Sunshine hours (hrs/day)	0.753**	0.506	- 0.419	- 0.646**	- 0.679**	- 0.396	- 0.853**	1.000

\*\* Correlation is significant at the 0.01 level (2-tailed)

\* Correlation is significant at the 0.05 level (2-tailed)

**Table 4. Multiple regression analysis for guava scab severity in relation to weather parameters on Allahabad Safed variety**

Location	Constant	X <sub>1</sub>	X <sub>2</sub>	X <sub>3</sub>	X <sub>4</sub>	X <sub>5</sub>	X <sub>6</sub>	X <sub>7</sub>	R	R <sup>2</sup>
MARS, Raichur	122.870	0.045	5.563	0.060	0.060	- 0.133	- 0.216	9.481	0.757	0.572
<b>Multiple linear regression equation</b>										
$Y = 122.870 + 0.045X_1 + 5.563X_2 + 0.060 X_3 + 0.060X_4 - 0.133X_5 - 0.216X_6 + 9.481X_7$										
<i>X<sub>1</sub>: Max. Temp, X<sub>2</sub>: Min. Temp, X<sub>3</sub>: Max. RH, X<sub>4</sub>: Min. RH, X<sub>5</sub>: Rainfall (mm), X<sub>6</sub>: Rainy days and X<sub>7</sub>: Sunshine hours</i>										

**Table 5. Correlation coefficient between weather parameters with guava scab on Lucknow 49 variety**

Parameters	Y	X <sub>1</sub>	X <sub>2</sub>	X <sub>3</sub>	X <sub>4</sub>	X <sub>5</sub>	X <sub>6</sub>	X <sub>7</sub>
Y PDI	1.000							
X <sub>1</sub> Maximum temperature (°C)	0.533*	1.000						
X <sub>2</sub> Minimum temperature (°C)	- 0.207	0.012	1.000					
X <sub>3</sub> Rainfall (mm)	- 0.239	- 0.565*	0.340	1.000				
X <sub>4</sub> Rainy days	- 0.226	- 0.551*	0.336	1.000**	1.000			
X <sub>5</sub> Relative humidity (max) (%)	- 0.244	- 0.520*	0.176	0.436	0.433	1.000		
X <sub>6</sub> Relative humidity (min) (%)	- 0.707**	- 0.602*	0.621	0.691**	0.680**	0.510	1.000	
X <sub>7</sub> Sunshine hours (hrs/day)	0.795**	0.624*	- 0.513	- 0.567*	- 0.556*	- 0.427	0.904**	1.000

\*\* Correlation is significant at the 0.01 level (2-tailed)  
\* Correlation is significant at the 0.05 level (2-tailed)

**Table 6. Multiple regression analysis for guava scab severity in relation to weather parameters on Lucknow 49 variety**

Location	Constant	X <sub>1</sub>	X <sub>2</sub>	X <sub>3</sub>	X <sub>4</sub>	X <sub>5</sub>	X <sub>6</sub>	X <sub>7</sub>	R	R <sup>2</sup>
MARS, Raichur	24.918	0.016	0.273	0.313	0.312	0.117	-0.068	10.505	0.817	0.668
<b>Multiple linear regression equation</b>										
$Y = 24.918 + 0.016X_1 + 0.273X_2 + 0.313X_3 + 0.312X_4 + 0.117X_5 - 0.068X_6 + 10.505X_7$										
<i>X<sub>1</sub>: Max. Temp, X<sub>2</sub>: Min. Temp, X<sub>3</sub>: Max. RH, X<sub>4</sub>: Min. RH, X<sub>5</sub>: Rainfall (mm), X<sub>6</sub>: Rainy days and X<sub>7</sub>: Sunshine hours</i>										

**3.3 Influence of Weather Parameters on Guava Scab Disease Severity on Arka Kiran Variety**

The first symptoms were observed on the Arka Kiran variety at 4<sup>th</sup> SMW (22<sup>nd</sup> January) with disease severity of 4.50 PDI and incidence of 2.91 per cent, the disease progressed and reached maximum at 12<sup>th</sup> SMW (19<sup>th</sup> March to 25<sup>th</sup> March) with severity and incidence of 74.51 and 85.00 per cent respectively (Table 1 & Fig. 3). The data on correlation of weather parameters revealed that, there was a significant positive correlation between individual parameters such as, maximum temperature (0.736), minimum temperature (0.643) per cent disease severity. Other parameters such as rainfall (0.432), rainy day (0.487) also had a positive correlation with disease development. However, the interaction between minimum temperature and maximum temperature (0.823), minimum temperature and rainy day (0.816),

rainfall and rainy day (0.867) and maximum and minimum relative humidity (0.886) had a significant positive correlation with the disease development and the interaction between minimum relative humidity and maximum temperature (- 0.711) had a significant negative correlation with the disease development (Table 7).The multiple linear regression of PDI of guava scab on Arka Kiran variety in relation to weather parameters during 2022-23 indicated that, the regression coefficients for maximum temperature (X<sub>1</sub>), minimum temperature (X<sub>2</sub>), rainfall (X<sub>3</sub>), rainy day (X<sub>4</sub>), morning relative humidity (X<sub>5</sub>), evening relative humidity (X<sub>6</sub>) and sunshine hours (X<sub>8</sub>) were found to be 12.233, 0.096, 0.127, 0.114, 0.038, + 0.062 and - 0.215 respectively. The multiple linear regression equation was fitted to the data and the equation arrived for the weather parameters is  $Y = 355.084 + 12.233 X_1 + 0.096X_2 + 0.127 X_3 + 0.114X_4 + 0.038X_5 + 0.062X_6 - 0.215$  (Table 8). This analysis revealed that, when there was

increase in one unit of maximum temperature, minimum temperature, rainfall, rainy day, morning relative humidity and evening relative humidity, the per cent disease severity was

increased by 12.233, 0.096, 0.127, 0.114, 0.038 and 0.062 units. Whereas increase in one unit of sunshine hours, the per cent disease severity was decreased by 0.215 units.




Fig. 3. Progression of guava scab in relation to weather parameters on Arka Kiran variety

Table 7. Correlation coefficient between weather parameters with guava scab severity on Arka Kiran variety

Parameters	Y	Y	X <sub>1</sub>	X <sub>2</sub>	X <sub>3</sub>	X <sub>4</sub>	X <sub>5</sub>	X <sub>6</sub>
Y PDI	1.000							
X <sub>1</sub> Maximum temperature (°C)	0.736*	1.000						
X <sub>2</sub> Minimum temperature (°C)	0.643*	0.823**	1.000					
X <sub>3</sub> Rainfall (mm)	0.432	0.449	0.640	1.000				
X <sub>4</sub> Rainy days	0.487	0.554	0.816**	0.867**	1.000			
X <sub>5</sub> Relative humidity (max) (%)	-0.631	-0.870**	-0.537	-0.159	-0.160	1.000		
X <sub>6</sub> Relative humidity (min) (%)	-0.493	-0.711*	-0.375	-0.303	-0.143	0.886**	1.000	
X <sub>7</sub> Sunshine hours (hrs/day)	-0.438	-0.336	-0.604	-0.152	-0.496	0.088	-0.294	1.000

\*\* Correlation is significant at the 0.01 level (2-tailed)

\* Correlation is significant at the 0.05 level (2-tailed)

Table 8. Multiple regression analysis for guava scab severity in relation to weather parameter on Arka Kiran variety

Location	Constant	X <sub>1</sub>	X <sub>2</sub>	X <sub>3</sub>	X <sub>4</sub>	X <sub>5</sub>	X <sub>6</sub>	X <sub>7</sub>	R	R <sup>2</sup>
MARS, Raichur	355.084	12.233	0.096	0.127	0.114	0.038	0.062	-0.215	0.736	0.542

Multiple linear regression equation

$$Y = 355.084 + 12.233X_1 + 0.096X_2 + 0.127X_3 + 0.114X_4 + 0.038X_5 + 0.062X_6 - 0.215X_7$$

X<sub>1</sub>: Max. Temp, X<sub>2</sub>: Min. Temp, X<sub>3</sub>: Max. RH, X<sub>4</sub>: Min. RH, X<sub>5</sub>: Rainfall (mm), X<sub>6</sub>: Rainy days and X<sub>7</sub>: Sunshine hours

Temperature has a major influence on the mycelial growth and sporulation of the pathogen. Temperature influences the conidial germination, appressoria formation and the germ-tube penetration into the host. More frequent rains help in inoculum dispersal, thus favourable weather condition set for maximum disease development. The influence of rain splashes on inoculum dispersal from an infection focus was well supported by the outcome of correlation and regression analysis when disease development was seen as dependent variable in relation with weather of preceding first week. Bright sunshine hours has a positive correlation with the disease development of guava scab which influenced the disease severity. These results are in confirmation with the findings of Khan and Hossain [8] who reported that, the lowest incidence and severity of leaf spot disease of coconut caused by *Pestalotiopsis palmarum* disease were recorded in January at temperature, relative humidity and rainfall of 16 °C, 77 per cent and no rain, respectively. The highest prevalence of incidence and severity were recorded in October at temperature of 27.9 °C, 82 per cent relative humidity and 122 mm rainfall. Based on the correlation and regression analyses they reported that, temperature was positively correlated with severity and incidence of the disease. As per the findings of Ismail et al [9] who studied on the effect of different epidemiological factors on severity and incidence of guava scab in Bangladesh, their study revealed that, in different growing seasons of guava seedlings, the maximum disease incidence (50.30 % and 66.84 %) and the highest severity (43.17 % and 56.19 %) of scab disease were recorded in January in which average temperature, relative humidity and rainfall were 16.88 °C, 73.80 % and 0.52 cm, and 30.50 °C, 83.50 %, 5.50 cm, respectively and the results are in contradictory to the findings made by Pan and Mishra [10]. The results are in agreement with the findings of Sharma [11] who reported the influence of weather parameters on northern leaf blight development of maize incited by *Exserohilum turcicum* (Pass.).

#### 4. CONCLUSION

In epidemiological studies, irrespective of the varieties, interaction among maximum temperature, minimum temperature, maximum relative humidity, minimum relative humidity, rainy days and rainfall showed significant positive

correlation with guava scab disease. Temperature has a major influence on the mycelial growth and sporulation. The optimum temperature influences conidial germination, appressoria formation and the germ-tube penetration into the host. The influence of rain splashes on inoculum dispersal from an infection focus was well supported by the outcome of correlation and regression analysis during disease development. Maximum scab disease severity and incidence was reported in Arka Kiran followed by Lucknow 49.

#### ACKNOWLEDGEMENT

I thank Department of Plant Pathology, College of Agriculture, Raichur, Centre for agroclimatic studies, MARS, Raichur and Department of Horticulture for providing all the facilities for fruitful completion of my research.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. Gutierrez RMP, Mitchell S, Solis VR. *Psidium guajava*: A review of its traditional uses, phytochemistry and pharmacology. J. Ethnopharmacol. 2008;117(1):1-27.
2. Candolle DAP. Origin of cultivated plants. Kegan Paul, London. 1904;48:376-384.
3. Anonymous, Area, production, productivity. Ministry of Agriculture and Farmers Welfare; 2021.
4. Dinesh MR, Vasugi C. Guava improvement in India and future needs. J. Horti. Sci. 2010;5(2):94-108.
5. Anonymous, Area, production, Productivity Of Guava; 2023. Available: <https://www.indiastat.com>
6. Keith LM, Velasquez ME, and Zee FT. Identification and Characterization of *Pestalotiopsis* spp. causing scab disease of guava, *Psidium guajava*, in Hawaii. Plant Dis. 2006;90(1):16-23.
7. Van der plank JE. Plant diseases: epidemics and control. Academic Press; 1963.
8. Khan MAH, Hossain I. Leaf spot disease of coconut seedling and its eco-friendly management, J. Bangladesh Agril. Univ. 2014;11(2):199-208.

9. Ismail AM, Cirvilleri G, Polizzi G. Characterisation and pathogenicity of *Pestalotiopsis uvicola* and *Pestalotiopsis clavisporea* causing grey leaf spot of mango (*Mangifera indica* L.) in Italy. European J. Plant Pathol. 2013;135:619-625.
10. Pan S, Mishra NK. Epidemiological studies on some diseases of guava (*Psidium guajava* L.). J. Plant Prot. Sci. 2014;2(4):49-52.
11. Sharma S. Influence of weather parameters on northern leaf blight development of maize incited by *Exserohilum turcicum* (Pass.) Leonard and Suggs in Nepal. Cogent Food & Agriculture. 2023;9(1):1-10.

---

© 2023 Sirisha et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/110043>