



# Polypropylene Packaged Ten Days Cold Stored Rose Cut Flowers cv. Bordeaux Influenced by Different Pulsing Solution

R. J. Makwana <sup>a++\*</sup>, Alka Singh <sup>b#</sup> and U. C. Chaudhary <sup>c++</sup>

<sup>a</sup> Sheth D. M. Polytechnic in Horticulture Model Farm, AAU, Vadodara, 390003, India.

<sup>b</sup> Department of Floriculture and Landscaping, ACHF, NAU, Navsari, India.

<sup>c</sup> College of Horticulture, AAU, Anand, 390003, India.

## Authors' contributions

*This work was carried out in collaboration among all authors. Author RJM designed the study, performed the statistical analysis, wrote the protocol, collected the data, data analysis and wrote the first draft of the manuscript. Author AS managed the analysis of the study, Author UCC managed the literature search. All authors read and approved the final manuscript.*

## Article Information

DOI: <https://doi.org/10.9734/ijbcrr/2024/v33i6914>

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/124707>

**Original Research Article**

**Received: 01/08/2024**

**Accepted: 03/10/2024**

**Published: 14/10/2024**

## ABSTRACT

The study included different pulsing solution viz., water, HQC 200mg/l, sucrose 50g/l + HQC 200 mg/l, citric acid 200mg/l,  $\alpha$ -lipoic acid 200mg/l, GA<sub>3</sub> 100 mg/l and BA 100mg/l to study the effect on flower quality and vase life of PP (polypropylene) packaged cold storage (at 2°C) rose cv.

<sup>++</sup> Assistant Professor;

<sup>#</sup> Professor and Head;

<sup>\*</sup>Corresponding author: E-mail: [drshimak@aaui.in](mailto:drshimak@aaui.in);

**Cite as:** Makwana, R. J., Alka Singh, and U. C. Chaudhary. 2024. "Polypropylene Packaged Ten Days Cold Stored Rose Cut Flowers Cv. Bordeaux Influenced by Different Pulsing Solution". *International Journal of Biochemistry Research & Review* 33 (6):315-21. <https://doi.org/10.9734/ijbcrr/2024/v33i6914>.

Bordeaux, for the period of 10 days. Pre-storage pulsing of GA<sub>3</sub> 100mg/l solution treatment showed highly promising results in extending flowers vase life along with maintained flower quality just after storage and at 4<sup>th</sup> DAS in vase life. Further, GA<sub>3</sub> 100mg/l solution treatment recorded significantly higher water up take (175.65 ml), change in fresh weight (10.50%) at 4<sup>th</sup> DAS, total dissolve solids (8.85 °brix, 8.05 °brix and 6.75 °brix) and membrane stability index (81.97 %, 65.97 % and 37.88 %) at 2<sup>nd</sup>, 4<sup>th</sup> and 6<sup>th</sup> DAS. Moreover, it was evident from the results that same treatment recoded significantly higher total soluble sugar (59.65 mg/g fresh weight), Estimated protein (18.79 mg/g fresh weight), catalase activity (20.42 μ mol H<sub>2</sub>O<sub>2</sub> min<sup>-1</sup> g<sup>-1</sup> protein), peroxidation activity (40.30 μ mol H<sub>2</sub>O<sub>2</sub> min<sup>-1</sup> g<sup>-1</sup> protein), pigment content (15.82 mg/g fresh weight) anthocyanin content in the petals at 4<sup>th</sup> DAS. Whereas, GA<sub>3</sub> 100 mg/l pulsing solution treated PP packaged cold stored rose cut flowers cv. Bordeaux recorded significantly higher vase life (6.88 days) which was at par with treatment P<sub>2</sub> (Sucrose 50 g/l + HQC 200 mg/l).

**Keywords:** Rose; polypropylene; low temperature storage; pulsing; GA<sub>3</sub>; sucrose.

## 1. INTRODUCTION

Roses are the most popular cut flowers around the world. Increasing popularity and demand in the world trade rose has become highly lucrative business amongst the flower growers. However, it is important to understand that the harvesting stages, postharvest handling and marketing techniques significantly affect the quality and the longevity of the flowers. It is very important to establish a well-planned post-harvest handling protocol to have best quality flowers in the market when it is high in demand. Makwana et al. [1] and Makwana et al. [2]. Observed that PP (Polypropylene) packaging along with cold storage at 2°C temperature provide better result comparing to different wet storage technique which is been used by most of the conventional flower growers and traders. Modified atmosphere packaging along with cold storage is known to create modified internal gaseous components passively Farber et al., [3] that minimizing metabolic activities during storage period and retain good flower quality after storage of 10 days [4]. Similar results with seal packaging of cut flowers with poly films at low temperature been found beneficial in retaining flower quality, improving opening ability, reducing water loss during post storage phase in stored flowers like rose [5,6] gladiolus [7,8]. Solidago Canadensis Zeltzer et al., [9] and in Lisianthus [10]. Makwana et al. [4] stated that higher CO<sub>2</sub> and lower O<sub>2</sub> during storage period decrease the production of ethylene and extended the life of fresh produce rose. Such beneficial conditions in orchid flowers showed promising results and maintained flower quality after completing of storage duration [11,12].

Plant growth regulators play an important role in quality of flower and extending storage life of

rose cut flowers [13]. Pulsing is a technique which is very important for extending post-harvest life of cut flowers. Pulsing treatment comprises of germicides and sugar are mostly used to improve flower diameter, flower size, shape, colour and life of cut flowers [8,14]. The 8-HQ has been known to possess strong anti-microbial properties that eliminate vascular blockage and enhance water up take in flowers [15,2]. Rose cut flowers kept in pulsing of GA<sub>3</sub> solution showed promising results from harvest to senescence [16,17]. Hence this experiment was laid out to evaluate best pulsing solution treatment to the 10 days cold stored rose cut flowers packed with PP material to get the better quality and post-harvest life of rose cut flowers cv. Bordeaux.

## 2. MATERIALS AND METHODS

Freshly harvest rose cut flowers of cv. Bordeaux were moved from greenhouse and brought to the Floriculture Laboratory, ASPEE College of Horticulture and Forestry, NAU Navsari at an ambient room temperature (18-21°C). The experiment was laid down in completely randomized block design. Total six Pulsing treatments and a control were taken under study along with each treatment replicated three times. Cut flowers of cv. Bordeaux having uniform bud size, shape and fresh weight (10±2 g) and stem length (50±5 cm) were selected and divided in to seven groups each having 30 flowers (10 flowers in each replication) and are provided with different pulsing solution treatment viz., P<sub>1</sub> - HQC 200 mg/l, P<sub>2</sub> - Sucrose 50 g/l + HQC 200 mg/l, P<sub>3</sub> - citric acid 200 mg/l, P<sub>4</sub> - α lipoic acid 200 mg/l, P<sub>5</sub> -GA<sub>3</sub> 100 mg/l, P<sub>6</sub> - BA 100 mg/l and P<sub>0</sub> - water for 3 hour before storage. After pulsing treatment all the bunches were packed with PP

packaging uniformly and moved to 2°C cold storage for the duration of 10 days. On completion of 10 day of storage duration, all the treated bunches were shifted from cold storage to laboratory and packaging were removed and stems were re cut 2 cm from the basal end and kept in distilled water at room temperature. The experiment was conducted for two times for two consecutive years represented as Year 1 and Year 2 for taking observations and recording data.

Data were collected regarding different postharvest parameters effecting quality of flowers at different intervals during vase life. Observations on post-harvest parameters like total water uptake (ml), change in fresh weight (%) at 4<sup>th</sup> DAS, total dissolve solids (TDS) at 2<sup>nd</sup>, 4<sup>th</sup> and 6<sup>th</sup> DAS, membrane stability index (MSI) at 2<sup>nd</sup>, 4<sup>th</sup> and 6<sup>th</sup> DAS, total soluble sugar (TSS), Estimated protein (mg/g fresh weight), catalase activity ( $\mu$  mol H<sub>2</sub>O<sub>2</sub> min<sup>-1</sup> g<sup>-1</sup> protein), peroxidaiton activity ( $\mu$  mol H<sub>2</sub>O<sub>2</sub> min<sup>-1</sup> g<sup>-1</sup> protein), pigment content (anthocyanin in the petals) were recorded 4<sup>th</sup> day after storage during vase life. TDS was measured with the help of digital refractometer in °Brix. Other parameters were recorded followed by methods listed below.

### 2.1 Membrane Stability Index (MSI) in petals

MSI was calculated on the basis of electrolyte leakage (ion leakage) of petals. The electrolyte leakage was measured by taking five petal discs (1cm<sup>2</sup>) of flower from the centre of the bud at 4<sup>th</sup> DAT. The petal discs were rinsed well in deionized water prior to incubation in 5 ml of deionized water for 3 h at room temperature. After incubation, conductivity of the bathing solution was measured with the conductivity meter (value A). The petal discs were boiled with bathing solution for 10 minutes to kill tissue. After cooling to room temperature, the conductivity of bathing solution was again measured (value B).

$$\text{Electrolyte Leakage (\%)} = (\text{value A} / \text{value B}) \times 100$$

$$\text{MSI (\%)} = [1 - (\text{value A} / \text{value B})] \times 100$$

### 2.2 Total Soluble Sugar (TSS) (mg/g fresh weight)

Petals (100 mg) were extracted with 5 ml of 80 per cent ethanol and centrifuged at 3000 rpm for

10 minutes. Extraction was repeated 4 times with 80 per cent ethanol and supernatants were collected into 25 ml volumetric flasks. Final volume of the extract was made to 25 ml with 80 per cent ethanol. The extract (0.3 ml) was pipetted from treatments into separate test tubes and the tubes were placed in a boiling water bath for 3 minutes to evaporate the ethanol. One ml of millipore water and 4 ml of 0.2 per cent anthrone reagent (200 mg in 100 ml H<sub>2</sub>SO<sub>4</sub>) was added in each test tube and placed in ice cold water. Reagent blank was prepared by adding 1 ml of distilled water and 4 ml of anthrone reagent. The intensity of colour was read at 600 nm on spectrophotometer. A standard curve was prepared using 10 mg glucose per 100 ml distilled water [18].

$$\text{Total soluble sugar (mg/g)} = \frac{\text{Sample O.D.} *}{\text{Standard O.D.} * \text{ Dilution factor}}$$

### 2.3 Estimation of Protein

Protein concentration of each enzyme extract and seed protein was estimated by method of Lowry et al. [19].

#### Reagents:

- (i) Solution A: 2% Na<sub>2</sub>CO<sub>3</sub> in 0.1 N NaOH
- (ii) Solution B : (a) 1% CuSO<sub>4</sub>.5H<sub>2</sub>O solution  
(b) 2% sodium potassium tartrate solution  
Working solution of B : Prepared fresh before use by mixing equal volume of solution B(a) and B(b).
- (iii) Solution C: Prepared fresh before use by mixing 50 ml of solution A and 1 ml of working solution of B.
- (iv) Solution D: Folin & Ciocalteu reagent (1N) reagent.

Enzymes extract/protein extract (25  $\mu$ l) were taken in test tube and volume was made up to 1 ml with millipore water. A tube with 1 ml of water served as blank. Five ml of solution C was mixed by vortexing and kept for 10 min. Then 0.5 ml of solution D (Folin & Ciocalteu reagent) was mixed with vortex and kept it room temperature for 30 min. Absorbance was read at 660 nm. A standard curve was prepared with bovine serum albumin in the range of 10-80  $\mu$ g.

## 2.4 Catalase Activity (CAT)

Total catalase (EC 1.11.1.6) activity was determined in the homogenates by measuring the decrease in absorption in 3ml mixture at 240nm as H<sub>2</sub>O<sub>2</sub> ( $\epsilon = 39.4 \text{ mM}^{-1} \text{ cm}^{-1}$ ) was consumed according the method of Aebi [20] and enzyme activity expressed as  $\mu\text{mol H}_2\text{O}_2$  oxidized  $\text{min}^{-1} \text{ g}^{-1}$  protein. The 3 ml mixture containing 50mM sodium phosphate buffer (pH 7.0), 10mM H<sub>2</sub>O<sub>2</sub> and 50 $\mu\text{l}$  enzyme extract.

## 2.5 Peroxidase (POD) Activity

POD (EC 1.11.1.7) activity was determined in the homogenates by measuring the increase in absorption at 470nm due to the formation of tetraguaiacol ( $\epsilon = 26.6 \text{ mM}^{-1} \text{ cm}^{-1}$ ) in a reaction mixture containing 50mM sodium phosphate buffer pH 7.0, 0.1mM EDTA, 0.1ml enzyme extract, 10mM guaiacol and 10mM H<sub>2</sub>O<sub>2</sub> [21].

**Enzyme unit of CAT and POD was calculated as:**

$$\text{mM/min/g protein} = \frac{\Delta \text{O.D.}}{\text{Enzyme Conc. (g)} * \epsilon}$$

Where,

$\epsilon$  = Extinction coefficient

Enzyme conc.(g) = Amount of enzyme in 3 ml reaction mixture

All the data recorded for various observations during the course of investigation were statistically analyzed (Panse and Sukhatme, 1978).

## 3. RESULTS AND DISCUSSION

Pulsing treatment significantly influence Physiological loss in weight, change in fresh weight, water uptake, bud diameter, per cent bud opening, MSI, TDS, dry weight and vase life of the flowers (Table 1). Among various pulsing solutions treated PP packed 10 days cold stored (2°C) rose cut spikes cv. Bordeaux under observations cut flowers treated with GA<sub>3</sub> 100 mg/l solution recorded significantly higher water up take (175.65 ml), change in fresh weight (10.50%) at 4<sup>th</sup> DAS, total dissolve solids (8.85 °brix, 8.05 °brix and 6.75 °brix) and membrane

stability index (81.97%, 65.97% and 37.88%) at 2<sup>nd</sup>, 4<sup>th</sup> and 6<sup>th</sup> DAS. Moreover, from Table 2, it was evident that same treatment recoded significantly higher total soluble sugar (59.65 mg/g fresh weight), Estimated protein (18.79 mg/g fresh weight), catalase activity (20.42  $\mu\text{mol H}_2\text{O}_2 \text{ min}^{-1} \text{ g}^{-1}$  protein), peroxidaiton activity (40.30  $\mu\text{mol H}_2\text{O}_2 \text{ min}^{-1} \text{ g}^{-1}$  protein), pigment content (15.82 mg/g fresh weight) anthocyanin content in the petals at 4<sup>th</sup> DAS. Whereas, GA<sub>3</sub> 100 mg/l pulsing solution treated PP packaged cold stored rose cut flowers cv. Bordeaux recorded significantly higher vase life (6.88 days) which was at par with treatment P<sub>2</sub> (Sucrose 50 g/l + HQC 200 mg/l).

GA<sub>3</sub> enhance the liposomal permeability of the cell membrane to glucose [22], hydrolyse starch, fructans and sucrose into glucose and fructose molecules [23]. This might have facilitated the higher intake of the sugar in the cell, which further enhanced water uptake due to osmotic pull [24]. It also known to be involved in mobilization of stored food Srivastava, [25] and further, to increase water uptake and retention of fresh weight [26]. Further antioxidant property of GA<sub>3</sub> contributed in stabilized cell membrane structure and thus reduced electrolyte leakage in the petal tissue as also reported in gladiolus [27], rose Sabehat and Zeislin, [28], in chrysanthemum [29]. Further, higher respiratory substrate and increased level of sugar in petal (TSS status, Table 2) facilitated higher rate of respiration necessary of cell division and cell enlargement (Ho and Nichols, 1977) leading to high TDS as well as high MSI [1]. The enhanced vase life of rose cut flowers pulsed with GA<sub>3</sub> 100 mg/l solution can be attributed to continued and increased water uptake, higher retention of fresh weight and also high maintained sugar level in petal tissue (high TSS), higher retained catalase and peroxidase activity and protein content resulted in delaying electrolytic leakage contributed to optimum continuation of the cell metabolism that facilitated cell growth and development, formation of cellular constituents and the liberation of energy for other cellular functions. Similar effects of enhanced postharvest vase life with GA<sub>3</sub> in rose Makwana et al. [1] and Makwana et al., [2] and gladiolus Dantuluri et al., [27] Singh et al., [30] and delayed leaf and petal abscission in rose Van Doorn et al., [31] and Ahmadi and Hassani [32] has been reported earlier [33].

**Table 1. Effect of different pulsing solution on total water uptake (ml), Change in fresh weight (%) at 4<sup>th</sup> DAS, Total dissolve solids (°brix) and membrane stability index (%) in PP packaged 10 days cold stored rose cut flowers cv. Bordeaux at 2<sup>nd</sup>, 4<sup>th</sup> and 6<sup>th</sup> DAS**

Treatment	Total water uptake (ml)	Change in fresh weight (%) at 4 <sup>th</sup> DAS	Total dissolve solids (°brix)			Membrane stability index (%)		
			2 <sup>nd</sup> DAS	4 <sup>th</sup> DAS	6 <sup>th</sup> DAS	2 <sup>nd</sup> DAS	4 <sup>th</sup> DAS	6 <sup>th</sup> DAS
P <sub>0</sub> (Water)	159.36	9.51	8.65	7.79	6.50	78.16	62.19	34.71
P <sub>1</sub> (HQC 200 mg/l)	161.45	9.63	8.68	7.88	6.58	78.63	62.60	34.58
P <sub>2</sub> (Sucrose 50 g/l + HQC 200 mg/l)	166.72	9.95	8.74	7.95	6.65	80.07	64.43	36.31
P <sub>3</sub> (Citric acid 200 mg/l)	162.62	9.76	8.69	7.89	6.59	78.34	62.28	34.23
P <sub>4</sub> (α-lipoic acid 200 mg/l)	165.02	9.75	8.70	7.90	6.60	78.37	62.31	34.31
P <sub>5</sub> (GA <sub>3</sub> 100 mg/l)	175.65	10.50	8.85	8.05	6.75	81.97	65.97	37.88
P <sub>6</sub> (BA 100 mg/l)	166.06	9.73	8.73	7.93	6.63	79.47	63.93	35.82
<b>CD at 5 %</b>	2.23	0.22	0.07	0.08	0.07	1.29	1.18	0.81

**Table 2. Effect of different pulsing solution on Total soluble sugar (mg/g fresh weight), estimated protein (mg/g fresh weight), Catalase activity (μ mol H<sub>2</sub>O<sub>2</sub> min<sup>-1</sup> g<sup>-1</sup> protein), Peroxidation activity (μ mol H<sub>2</sub>O<sub>2</sub> min<sup>-1</sup> g<sup>-1</sup> protein), Anthocyanin content (mg/g fresh weight) and vase life (days) in PP packaged 10 days cold stored rose cut flowers cv. Bordeaux at 4<sup>th</sup> DAS**

Treatment	Total soluble sugar (mg/g)	Estimated Protein (mg/g)	Catalase activity (μ mol H <sub>2</sub> O <sub>2</sub> min <sup>-1</sup> g <sup>-1</sup> protein)	Peroxidation activity (μ mol H <sub>2</sub> O <sub>2</sub> min <sup>-1</sup> g <sup>-1</sup> protein)	Anthocyanin content (mg/g)	Vase life (days)
P <sub>0</sub> (Water)	54.07	17.70	18.82	36.88	14.14	5.28
P <sub>1</sub> (HQC 200 mg/l)	55.20	17.56	18.85	37.09	14.39	6.30
P <sub>2</sub> (Sucrose 50 g/l + HQC 200 mg/l)	56.44	18.12	19.97	39.14	15.71	6.79
P <sub>3</sub> (Citric acid 200 mg/l)	54.85	17.64	19.18	36.37	14.87	6.21
P <sub>4</sub> (α-lipoic acid 200 mg/l)	55.14	17.71	17.05	37.07	14.71	6.23
P <sub>5</sub> (GA <sub>3</sub> 100 mg/l)	59.65	18.49	20.42	40.30	15.82	6.88
P <sub>6</sub> (BA 100 mg/l)	55.55	17.71	19.41	38.63	15.15	6.45
<b>CD at 5 %</b>	1.30	0.34	0.47	1.20	0.46	0.12

#### 4. CONCLUSION

In conclusion, GA<sub>3</sub> 100 mg/l solution as pulsing treatment had a significant effect on increasing vase life of PP packed rose cut flowers cv. Bordeaux cold stored (2°C) for 10 days and improving their quality of flowers by increase in retention of fresh weight, higher total water uptake, total soluble sugar, estimated protein, catalase activity and peroxidase activity and therefore improving flowers quality.

#### DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

#### ETHICAL APPROVAL

All experimental protocols including the involvement of human participants were approved and followed the guidelines of the Department of floriculture and landscaping of Aspee College of Horticulture of the Navsari Agricultural University, Navsari.

#### ACKNOWLEDGEMENT

I convey the depth of my feeling and gratitude to my major advisor Dr. B. K. Dhaduk and to my advisory committee members Dr. Alka Singh and other for giving me proper guidance throughout the course of study.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. Makwana RJ, Singh A, Dhaduk BK. Cold stored cut spikes influenced by different storage techniques in rose CVs. Bordeaux, poison and avalanche. International Journal of Advanced Biochemistry Research. 2024a;SP-8(1):402-405.
2. Makwana RJ, Singh A, Neelima P. Effect of cold storage techniques on flower quality and vase life of rose var. 'Sunking'. The Bioscan. 2015a;10(1):1-4.
3. Farber JN, Harris LJ, Parish ME, Beuchat LR, Suslow TV, Gorney JR, Garrett EH, Busta FF. Microbiological safety of controlled and modified atmosphere packaging of fresh and fresh-cut produce. Comprehensive Review in Food Sci. and Food Safety. 2003;2:142-158.
4. Makwana RJ, Singh A, Dhaduk BK. Effect of Low Temperature Storage Techniques on Post Storage Quality of Rose cvs. Bordeaux, Poison and Avalanche. Biological Forum – An International Journal. 2024b;16(1):22-25.
5. Singh A, Makwana RJ, Mangave BD. Influence of low temperature storage techniques on post-storage quality of rose cut flowers var. Passion. International Journal of Innovative Horticulture. 2012;1(2):151-157, 2012
6. Makwana RJ, Singh A, Ahlawat TR, Neelima P. Standardization of low temperature storage technology with novel packaging techniques in rose cut flower cv. Passion. HortFlora Research Spectrum. 2015b;4(1): 44-47.
7. Grover JK, Singh K, Gupta AK, Ashok K. Effect of pre-storage pulsing treatment on storage of gladiolus spikes under modified atmosphere. Indian J. Plant Physiol. 2005; 10(3):248-253.
8. Singh A, Dhaduk BK, Desai JR. Post harvest Technology and Value Addition in Cut Flowers. Floriculture Today. 2007;12(4):40-46.
9. Zeltzer S, Meir S, Mayak S. Modified atmosphere packaging (map) for long-term shipment of cut flowers. Acta Hort. 2001; 553:631-634
10. Akbudak B, Erisand A, Kucukahmetler O. Normal and modified atmosphere packaging storage of lisianthus (*Lisianthus grandiflorum*) grown in saline conditions. New Zealand J. Crop and Hort. Sci. 2005; 33:185-191.
11. Poonsri W. Storage of Dendrobium orchid in modified atmosphere storage system of agriculture products for retailer. Agric. Sci. 2017;46(3/1):243-246.
12. Poonsri W. Effects of active and passive modified atmosphere packaging on biochemical properties of cut Dendrobium orchid flowers. Heliyon. 2021;7(6):1-10.
13. Sharma KA. review of plant growth regulators and its effect on rose flower. Int. J Adv. Biochem. Res. 2023;SP- 7(2):366-370.
14. Vidhya S., Bhattacharjee SK. Floriculture Research Trend in India. Ind Soc. Ornamental Hort. New Delhi, 2002;83-86.

15. Burdett AN. The cause of bent neck in cut roses. J. American. Soc. Hort. Sci. 1970; 95:427-431.
16. Bhattacharjee SK. Postharvest life of "Eiffel Tower" cut roses and biochemical constituents of petal tissues as influenced by growth regulating chemicals in the holding solution. Haryana J. Hort. Sci. 2000;29(1/2):66-68
17. Makwana RJ, Dhaduk BK, Singh A. Effect of pulsing solution on quality and post-harvest life in PP packaged cold stored rose cut flowers cv. Bordeaux. International Journal of Creative Research. 2024c;12(1):178-181
18. Franscistt W, David FB, Robert MD. The estimation of the total soluble carbohydrate in cauliflower tissue. Experiment in plant physiology, Van, Nostrand. Reinhold Camp., New York. 1971;16.
19. Lowry OH, Rosbrough NJ, Farr AL, Randall RJ. Protein measurement with the folin phenol reagent. J. Biol. Chem. 1951; 193:265-275.
20. Aebi H. Catalase *in vitro*. *Meth. Enzymol.*, 1984; 105 : 121-126.
21. Costa H, Gallego SM, Tomaro ML. Effect of UV-B radiation on antioxidant defense system in sunflower cotyledons. Plant Science. 2002;62:939-945.
22. Wood A, Paleg LG. The influence of GA3 on the permeability of model membrane systems. Plant Physiol. 1972; 50, 103–108.
23. Salisbury FB, Ross CW. Plant Physiology, Hormones and Plant Regulators: Auxins and Gibberellins. 4th Edition, Wadsworth Publishing, Belmont. 1992; 357-381.
24. Ho LC, Nichols R. Translocation of <sup>14</sup>C sucrose in relation to change in carbohydrate content in rose corollas cut at different stages of development. Annals of Botany. 1997; 41:227-242.
25. Srivastava R, Kandpal K, Jauhari S. Effect of pulsing solution, packaging material and storage duration on post harvest life of gladiolus. J. Ornam. Hort. 2005;8(2):115-118.
26. Van Doorn WG. Is petal Senescence due to sugar starvation. Plant Physiology. 2004;134:35-42.
27. Dantuluri VSR, Misra RL, Singh VP. Effect of polyamines on post-harvest life of gladiolus spikes. Journal of Ornamental Horticulture. 2008;11(1):66-68.
28. Sabehat A, Zeislin N. GA<sub>3</sub> effects on post harvest alteration in cell membranes of rose (Rosa x Hybrida) petals. Journal of Plant Physiology. 1994;144: 513–7.
29. Elanchezhian R, Srivastava GC. Effect of growth regulators on senescence of chrysanthemum flowers. Indian J Plant Physiol. 2001;6:233-243
30. Singh A, Kumar J, Kumar P. Effects of plant growth regulators and sucrose on post harvest physiology, membrane stability and vase life of cut spikes of gladiolus. Plant Growth Regul. 2008;55:221-229.
31. Van Doorn WG, K. D'Hont. Interaction between the effects of bacteria and dry storage on the opening and water relations of cut rose flowers. J. App. Bacteriology. 1994;77(6):644-649.
32. Ahmadi Z, Hassani RN. Effect of Gibberellic Acid Pulsing and Sucrose Continuous Treatment on Some Qualitative Characteristics of Cut Rose Flower cv. Velvet. Journal of Ornamental Plants. 2015;5(3):189-195.
33. Panse VG, Sukhatme PV. Statistical Methods for Agricultural Workers. Publication and Information Division, ICAR, New Delhi. 1967.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/124707>