



Ameliorative Effects of N-Acetyl Cysteine and Zinc Sulfate on Reproductive Dysfunction Induced by Short-Term Crude Oil Exposure in Male Wistar Rats

Tarela Melish Elias Daubry ^a,
Olaghaguo Macstephen Adienbo ^b,
Blessing Zeinab Ovili-Odili ^a
and Bruno Chukwuemeka Chinko ^{b*}

^a Department of Physiology, Faculty of Basic Medical Sciences, Delta State University Abraka, Delta State, Nigeria.

^b Department of Human Physiology, Faculty of Basic Medical Sciences, University of Port Harcourt, Port Harcourt, Rivers State, Nigeria.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/ajmah/2024/v22i111111>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/125280>

Original Research Article

Received: 11/08/2024
Accepted: 15/10/2024
Published: 19/10/2024

*Corresponding author: E-mail: bruno.chinko@uniport.edu.ng;

Cite as: Daubry, Tarela Melish Elias, Olaghaguo Macstephen Adienbo, Blessing Zeinab Ovili-Odili, and Bruno Chukwuemeka Chinko. 2024. "Ameliorative Effects of N-Acetyl Cysteine and Zinc Sulfate on Reproductive Dysfunction Induced by Short-Term Crude Oil Exposure in Male Wistar Rats". *Asian Journal of Medicine and Health* 22 (11):1-11. <https://doi.org/10.9734/ajmah/2024/v22i111111>.

ABSTRACT

Background: Crude oil contamination by accidental or voluntary ingestion is prevalent in the Niger Delta region of Nigeria. One of the possible impacts of crude oil contamination is reproductive toxicity leading to infertility. The present study evaluated the potential protective effects of N-Acetyl Cysteine (NAC) and Zinc Sulfate ($ZnSO_4$) in mitigating reproductive dysfunction caused by short-term Bonny Light Crude oil (BLCO) exposure in male Wistar rats.

Materials and Methods: Fifty (50) male Wistar (180 – 200g) were used for the study. They were divided into ten (10) groups of five (5) animals each. Groups I and II served as the control and negative control and received distilled water and BLCO (600mg/kg) respectively while groups III to X served as the experimental groups and received NAC (100 and 200mg/kg) and $ZnSO_4$ (0.5mg/kg and 1mg/kg) orally for three (3) weeks. Animals were euthanized, and blood and semen were collected for biochemical and seminal analysis.

Results: The results of this current study show that BLCO exposure caused a disruption in reproductive functions in rats by decreasing reproductive hormones and semen quality parameters in Wistar rats. NAC and $ZnSO_4$ administration significantly improved semen quality parameters by improving sperm count, sperm motility and morphology in experimental rats.

Conclusion: Evidence from the present study suggests that NAC and $ZnSO_4$ protected the testes, preventing reproductive alterations linked to BLCO exposure.

Keywords: Crude Oil; bonny crude light oil; N-Acetyl cysteine; zinc sulphate; reproductive dysfunction.

1. INTRODUCTION

Petroleum pollution is a prevalent issue that can occur both accidentally and during routine operations wherever oil is produced, transported, stored, processed, or utilized, whether on land or at sea. On land, petroleum products often represent a significant portion of the contaminants found at polluted sites (Orisakwe et al. 2004, Alabi et al. 2022). Despite the economic importance of crude oil in Nigeria, its exploration has brought about many pollution problems, caused by oil spillage which is a common occurrence in Nigeria (Adeola et al. 2022, Ite et al. 2013). Interestingly, the traditional practice of consuming crude oil for its purported medicinal benefits is becoming more widespread (Ajonuma et al. 2023). They are directly used as curative agents for anti-poisoning (snake venom antidotes), anti-convulsion, treatment of skin infection or indirectly by eating marine animals found in surrounding coastal waters as a source of protein (Dede et al. 2002). Bonny light crude oil (BLCO) is used in combination with olive oil in folklore medicine in some parts of the Niger Delta region of Nigeria to treat burns, gastrointestinal disorders, ulcers, witchcraft attacks and poisoning (Dede et al. 2002, Raji et al. 2012, Orisakwe et al. 2000). The pattern of crude oil toxicity involves multiple mechanisms that cause damage to living organisms: Polycyclic aromatic hydrocarbons (PAHs) in crude oil generate reactive oxygen species (ROS), causing oxidative stress that damages cells and DNA, leading to mutations and cancer risks (Han et al.

2014, Adedara et al. 2012). These PAHs are also known to disrupt heart development and function. Similarly, volatile organic compounds (VOCs) like benzene impair nervous system and respiratory system function, leading to cognitive, behavioural issues, and long-term lung damage (Noh et al. 2023, Khan et al. 2023, Ritchie et al. 2001). Studies have equally demonstrated significant immunotoxicity due to crude oil exposure, leading to a weakened immune system and increased susceptibility to diseases (Weatherly et al. 2022, McLoone et al. 2021). Several chemicals contained in BLCO have been identified as endocrine disruptors, resulting in unwanted reproductive malfunction and developmental disorders (Raji et al. 2012, Indarto and Izawa 2001, Adesanya et al. 2009, Farombi et al. 2010, Naiho et al. 2014).

Infertility is one of the major health problems in different societies (Daubry et al. 2022), affecting between fifty and eighty million couples at some point in their reproductive lives globally (Boivin et al. 2007). Approximately 8% of men seek medical help for fertility-associated problems (Friedler et al. 2002). An estimated 3-4 million Nigerian couples have fertility-associated problems (Sule et al. 2008). Infertility affects about 20-30% of couples in Nigeria and around the world, irrespective of their race or ethnicity (Bablok et al. 2001). Studies have shown that BLCO significantly impairs reproductive functions by causing reduced sperm count, sperm motility and normal morphology within seven days of administration (Orisakwe et al. 2004, Adesanya et

al. 2009, Farombi et al. 2010). The mechanism behind the reproductive damage caused by BLCO is associated with the generation of free radicals, reactive oxygen and reactive nitrogen species which overwhelm cellular antioxidant systems and induce cellular damage (Saulsbury et al. 2009). Oxidative stress occurs when the balance between the production of reactive species and the antioxidant defence system is disrupted (Zahra et al. 2021, Bano et al. 2021, Chinko et al. 2020). Cellular injury is primarily due to the inability of the antioxidants to neutralize the effects of the oxygen radicals, which have ultimately led to infertility (Saulsbury et al. 2009). Antioxidants are molecules that prevent oxidative damage by scavenging reactive oxygen species (ROS) or inhibiting their production, thereby limiting or preventing the oxidation of other molecules (Gulcin 2020, Neha et al. 2019, Bano et al. 2021, Awajiomowa et al. 2022). Antioxidant compounds such as N-acetylcysteine (NAC) play an important role in the protection of cell constituents from oxidative stress (Moschou et al. 2008) by acting as a cysteine supplier, maintaining and increasing the intracellular levels of glutathione (Atkuri et al. 2007, Sadowska et al. 2007). Zinc (Zn) is another important trace element found in small amounts in a variety of cells and tissues of organisms, it is a cofactor of more than 300 enzymes (Tapiero and Tew 2003). It is reported to be necessary for signal transduction, DNA replication, RNA polymerases, protein synthesis, growth processes and various metabolic processes (Cousins et al. 2006). Various studies have shown that zinc sulfate may act as an antioxidant (Alvarez and Storey 1983), capacitating agent (Meizel 1985, Meizel et al. 1980), membrane-stabilizing factor (Mrsny and Meizel 1985) and sperm motility factor (Boatman et al. 1990, Egwurugwu et al. 2013). Research is scarce on the potential ameliorative effects of these key antioxidant compounds on reproductive dysfunction caused by crude oil exposure. This study aims to address this gap by evaluating the potential protective role of N-Acetyl Cysteine and Zinc Sulfate in mitigating reproductive dysfunction induced by short-term crude oil exposure in male Wistar rats.

2. MATERIALS AND METHODS

2.1 Source of Crude, Drugs and Research Animals

Bonny light crude oil was obtained from the Nigerian National Petroleum Corporation (NNPC) Warri, Nigeria. N-acetyl cysteine was purchased

from Sigma Aldrich, USA (A7250-10G) while Zinc Sulfate was sourced locally (Bactolac Pharmaceutical Inc, US). Fifty (50) male Wistar (180 – 200g) were sourced from the Animal house of the Department of Physiology, Delta State University, Abraka and used for the study. They were allowed three (3) weeks of acclimatization under standard animal husbandry conditions. The animals were housed in clean, well-ventilated wooden cages under optimal conditions, including a 12-hour light/dark cycle, a temperature range of 28–31°C, and humidity levels of 45–50%. They had unrestricted access to standard rat pellets and tap water. The animals were divided into ten (10) groups of five (5) animals each and received BLCO (Raji and Hart 2012, Orisakwe et al. 2000), NaC and ZnSO₄ (Al-Ani et al. 2015, Al-Hamdany and Al-Hubaity 2014) orally via orogastric cannula between 8 am and 10 am daily for three (3) weeks according to the following protocol:

Group I	Distilled water only (control)
Group II	BLCO (600mg/kg) only (negative control)
Group III	NAC (100mg/kg) only
Group IV	NAC (200mg/kg) only
Group V	NAC (100mg/kg) + BLCO (600mg/kg)
Group VI	NAC (200mg/kg) + BLCO (600mg/kg)
Group VII	ZnSO ₄ (0.5 mg/kg) only
Group VIII	ZnSO ₄ (1mg/kg) only
Group IX	ZnSO ₄ (0.5mg/kg) + BLCO (600mg/kg)
Group X	ZnSO ₄ (1mg/kg) + BLCO (600mg/kg)

2.2 Sample Collection and Biochemical Laboratory Assay

At the end of the experimental period, the animals were fasted overnight after which they were euthanized by cervical dislocation and cardiac puncture was used to collect blood samples for hormonal and biochemical analysis. The enzyme-linked immunosorbent assay technique was used for follicle-stimulating hormone (FSH), luteinizing hormone (LH), testosterone, estrogen and prolactin estimation using standard ELIZA kits (Cal biotech Inc. California). The procedure for the estimation of the serum concentration of each hormone was carried out according to the kits' manual as earlier described by Khourdaji et al. (2018).

2.3 Semen Analysis

For sperm count, the caudal epididymis was homogenized in formal-saline and the sperm count was carried out using the improved Neubauer counting chamber (LABART, Germany). The sperm count was examined under the light microscope at a magnification of

x40 while evaluating different fields and was calculated using the formula described by Omirinde et al. (2019). Sperm motility was determined by conventional method as described by Khatun et al. (2018). After the sperm had been squeezed on the pre-warmed slide, two drops of warm 2.9% sodium citrate were added to it, then covered with a cover slip and examined under the microscope using an X40 objective with reduced light. Sperm morphology was done by collecting a thin smear of the sperm sample on a clean slide, fixed with 95% ethanol and allowed to air dry. The fixed slide was then sequentially immersed into different concentrations of ethanol and appropriately stains namely Harris haematoxylin, G-6 orange stain and EA-50 green stain for one minute. The slide was then examined microscopically at a magnification of x40, 200 sperm were assessed and the sperm abnormalities were expressed in percentages (Omirinde et al. 2019, Yu et al. 2018). Sperm viability (live/death ratio) was done by observing the percentage of spermatozoa in a unidirectional progressive movement over a field on a slide under the light microscope fitted with a camera using the Eosin/Nigrosin stain (Omirinde et al. 2019).

2.4 Statistical Analysis

Data obtained from this study was analyzed using Graph pad Prism 8 Biostatistics software (Graph pad Software, Inc., Lajolla, USA version 8.0). All data was presented as Mean±SEM. Further analysis was done by one analysis of

variance (ANOVA) and followed by a post hoc test (Bonferroni) for multiple comparisons. The level of significance for all tests was set at p < 0.05.

3. RESULTS

Table 1 illustrates the effects of NAC and ZnSO₄ administration on testosterone, estrogen, follicle-stimulating hormone (FSH), luteinizing hormone (LH), and prolactin levels in BLCO-exposed Wistar rats. The data reveal that BLCO exposure in Wistar rats significantly reduced levels of testosterone, estrogen, FSH, LH, and prolactin compared to the control group. Co-treatment with NAC and ZnSO₄ led to various improvements in these hormone levels. NAC and ZnSO₄ treatments significantly increased testosterone levels, with dose-dependent effects, but testosterone was still lower when co-treated with NAC_a + BLCO and ZnSO₄ + BLCO compared to their respective single treatments. BLCO exposure decreased estrogen levels, but NAC_b + BLCO and ZnSO₄ + BLCO treatments led to significant increases. However, NAC_a + BLCO still showed reduced estrogen levels. Both FSH and LH were significantly reduced by BLCO exposure. While NAC and ZnSO₄ co-treatment did not improve FSH levels, they significantly increased LH levels. Also, BLCO exposure decreased prolactin levels, but co-treatment with NAC and ZnSO₄ significantly restored prolactin levels.

Table 1. Role of NAC and ZnSO₄ administration on Testosterone, Estrogen, Follicle stimulating hormone, Luteinizing hormone and Prolactin level of BLCO-exposed Wistar rats

Groups/Doses/(kg bw)	Testosterone (ng/ml)	Estrogen (pg/ml)	FSH (miU/ml)	LH (miU/ml)	Prolactin (µg/L)
Control	78.03±1.28	0.61±0.01	1.68±0.01	3.92±0.04	6.23±0.13
BLCO (600mg/kg)	42.43±1.37 [*]	0.39±0.02 [*]	0.23±0.02 [*]	1.34±0.03 [*]	4.52±0.27 [*]
NAC (100mg/kg)	75.27±0.87 [#]	0.57±0.01 [#]	1.09±0.03 ^{*#}	2.69±0.08 [#]	6.05±0.10 [#]
NAC (200mg/kg)	72.04±2.17 [#]	0.41±0.01 ^{*#}	1.24±0.01 ^{*#}	2.76±0.07 [#]	5.09±0.20 ^{*#}
NAC(100mg/kg)+ BLCO(600mg/kg)	69.04±0.62 ^{*#β}	0.47±0.01 ^{*α}	0.19±0.01 ^{*αβ}	2.14±0.01 ^{*#β}	5.88±0.13 [#]
NAC(200mg/kg) + BLCO(600mg/kg)	74.24±0.47 [#]	0.57±0.02 ^{#β}	0.22±0.01 ^{*αβ}	2.32±0.07 ^{*α}	5.68±0.34 [#]
ZnSO ₄ (0.5mg/kg)	60.48±0.79 [#]	0.46±0.01 [*]	1.02±0.02 [#]	2.65±0.05 [#]	5.53±0.15
ZnSO ₄ (1mg/kg)	73.43±0.41 [#]	0.54±0.02 [#]	0.87±0.01 ^{*#}	3.43±0.05 [#]	5.83±0.26 [#]
ZnSO ₄ (0.5mg/kg) + BLCO(600mg/kg)	36.22±1.49 ^{*#ab}	0.52±0.01 ^{*#}	0.23±0.01 ^{*ab}	2.81±0.14 ^{*#ab}	5.31±0.26
ZnSO ₄ (1mg/kg) + BLCO(600mg/kg)	53.60±0.80 ^{*#abc}	0.54±0.01 [#]	0.13±0.02 ^{*#ab}	2.84±0.24 ^{*#ab}	5.52±0.12

Values are expressed as Mean±SEM (n = 5) (ANOVA followed by Tukey's test).
 BLCO. (BLCO=Bony Light Crude Oil. NAC = N-acetylcysteine. ZnSO₄ = Zinc Sulfate).
^{*}P<0.05: significantly different when compared with the control group.
[#]P<0.05: Significantly different when compared with BLCO.
^αP<0.05: significantly different when compared with NAC_a (100mg/kg).
^βP<0.05: significantly different when compared with NAC_b (200mg/kg).
^φP<0.05: Dose-dependent significant difference.
^aP<0.05: significantly different when compared with ZnSO_{4a}(0.5mg/kg).
^bP<0.05: significantly different when compared with ZnSO_{4a}(1mg/kg).
^cP<0.05: significantly different when compared with ZnSO_{4a}(0.5mg/kg)

Table 2. Role of NAC and ZnSO₄ administration on sperm count, motility, morphology, and viability of BLCO-exposed Wistar rats

Groups/Doses/(kg bw)	Sperm Count (x10 ⁻⁶ cells/mm ³)	Sperm Motility (%)	Sperm Morphology (%)	Sperm Viability (%)
Control	110.20±1.77	76.24±1.22	72.80±0.58	81.60±1.03
BLCO (600mg/kg)	76.48±3.86*	42.38±0.95*	43.40±0.97*	46.20±1.35*
NAC (100mg/kg)	94.40±1.69*#	58.82±0.68*#	56.60±0.81*#	72.20±0.66*#
NAC (200mg/kg)	109.80±2.63# ^φ	63.40±2.15*#	68.20±1.15*# ^φ	69.40±1.93*#
NAC(100mg/kg)+ BLCO(600mg/kg)	97.40±2.61*#	46.40±0.24* ^β	57.20±0.73*# ^β	57.80±1.28*# ^{αβ}
NAC(200mg/kg) + BLCO(600mg/kg)	106.70±2.79#	51.80±0.58*# ^β	67.20±1.35*# ^α	61.00±1.51*# ^{αβ}
ZnSO ₄ (0.5mg/kg)	79.70±2.80*	47.37±1.07*	60.40±0.81*# ^φ	57.80±1.28*#
ZnSO ₄ (1mg/kg)	89.13±2.58*	60.32±0.03*# ^φ	66.60±0.92*#	73.60±1.96*#
ZnSO ₄ (0.5mg/kg) + BLCO(600mg/kg)	105.30±2.24* ^a	59.66±2.52*# ^a	62.00±1.37*#	55.60±1.12*# ^b
ZnSO ₄ (1mg/kg) + BLCO(600mg/kg)	84.11±1.80* ^{bc}	67.55±1.23*# ^{abc}	66.60±1.12*# ^a	73.60±0.74*# ^a

Values are expressed as Mean±SEM (n = 5) (ANOVA followed by Tukey's test).

BLCO. (BLCO=Bony Light Crude Oil. NAC = N-acetylcysteine. ZnSO₄ = Zinc Sulfate).

*P<0.05: significantly different when compared with the control group.

#P<0.05: Significantly different when compared with BLCO.

^aP<0.05: significantly different when compared with NAC_a (100mg/kg).

^βP<0.05: significantly different when compared with NAC_b (200mg/kg).

^φP<0.05: Dose-dependent significant difference.

^aP<0.05: significantly different when compared with ZnSO_{4a}(0.5mg/kg).

^bP<0.05: significantly different when compared with ZnSO_{4a}(1mg/kg).

^cP<0.05: significantly different when compared with ZnSO_{4a}(0.5mg/kg)

Table 2 shows the effect of NAC and ZnSO₄ administration on sperm count, motility, morphology, and viability of BLCO-exposed Wistar rats. The results show that BLCO exposure in Wistar rats significantly impaired sperm health, including reductions in sperm count, motility, morphology, and viability. Co-treatment with NAC and ZnSO₄ helped mitigate these effects. BLCO exposure significantly reduced sperm count, but co-treatment with NAC and ZnSO₄ (especially at higher doses) increased sperm count in a dose-dependent manner. BLCO significantly decreased sperm motility. Co-treatment with NAC_b and ZnSO₄ restored motility, although NAC + BLCO treatment was less effective compared to NAC alone, while ZnSO₄ + BLCO improved motility over ZnSO₄ alone. BLCO exposure led to a significant decrease in normal sperm morphology. Co-treatment with NAC and ZnSO₄ significantly improved morphology in a dose-dependent manner, though the co-treated groups still showed lower improvements compared to NAC or ZnSO₄ alone. Finally, BLCO reduced sperm viability, but co-treatment with NAC and ZnSO₄ increased viable sperm cells. However, sperm viability remained lower in co-treated

groups compared to NAC or ZnSO₄ alone, with ZnSO₄ treatment showing a dose-dependent increase.

4. DISCUSSION

Crude oil pollution can occur accidentally during transportation or as a routine consequence of production and processing. While accidental ingestion may result from contamination, there are instances where crude oil is voluntarily consumed by humans due to its purported medicinal benefits. Both accidental and voluntary ingestion are common in Nigeria's Niger Delta region. This study assessed the potential protective effects of N-Acetyl Cysteine and Zinc Sulfate in mitigating reproductive dysfunction caused by short-term crude oil exposure in male Wistar rats.

Effect of NAC and ZnSO₄ on reproductive hormones of BLCO-exposed Wistar rats: The result from the present study shows that BLCO significantly reduced reproductive hormones in Wistar rats after three weeks when compared with the control group (p<0.05, Table 1). This shows that BLCO contamination and

consumption could negatively regulate releasing of gonadotropins through the pituitary gland. FSH and LH are the main factors needed for the development of testicles and for proper stimulation of releasing of reproduction hormones like testosterone, estrogen and progesterone. The decrease in the level of these bio-substances in rats treated with BLCO could mean a negative effect on the anterior pituitary gland which influences normal control and regulation of processes involved in secreting these reproduction-related hormones. Findings from the present study are consistent with studies done by (Raji and Hart 2012, Farombi et al. 2010, Naiho et al. 2014, Nwafor et al. 2013, Georgewill and Nwankwoala 2006). The reduced level of testosterone in BLCO-administered rats could be an indicator of the chemical toxic level in the male reproductive organ, which is crucial and required to sustain spermatogenesis and also to maintain their function and structure as regards male sex glands (Oduwole et al. 2021). Our study observed that co-administration of BLCO with NAC and ZnSO₄ substantially increased all reproductive hormone levels. This finding is consistent with the study by Nashwa et al. (2011) where a notable increase in testosterone, LH and FSH levels was observed due to NACs. NACs are precursors of L-cysteine which leads to glutathione elevation synthesis. They are antioxidant agents and common nutrition supplements, applied anti-oxidant *in-vivo* and *in-vitro*. It acts directly as a scavenger of free radicals, especially oxygen. It is equally recommended as a possible remedial alternative to several illnesses that arise from the formation of oxygenated radicals (Mokhtari et al. 2016). NACs could prevent the negative impact of different toxins like arsenic, lead and cadmium on male reproduction (Reddy et al. 2011, Kumar et al. 2016). A similar result also shows that co-administration of ZnSO₄ with BLCO significantly increased all reproductive hormones (Table 1). This corroborates findings from previous studies of Egwurugwu et al. (2013) on the effect of ZnSO₄ on male reproductive hormones. Zinc supplementation activates the secretion and action of testosterone and can lead to increased efficiency of spermatogenic machinery and increased number of germ cells in the seminiferous tubules (Abdella et al. 2011). The reduction in serum levels of FSH and non-significant effect on Luteinizing hormone (LH) following the increased levels of testosterone may be attributed to the negative feedback effect of testosterone on the hypothalamus which in turn causes a decrease in the

secretion of FSH and LH by the anterior pituitary gland. This observed improvement in the values of testosterone in the ZnSO₄-supplemented group might be due to the stimulating effect of ZnSO₄ on testicular steroidogenesis as ZnSO₄ has been shown to affect testicular functions by activating the adenyl cyclase system, which stimulates testosterone synthesis (El-Masry et al. 1994). Overall, NAC and ZnSO₄ provided some protective effects against BLCO-induced hormone disruptions, particularly for testosterone, estrogen, LH, and prolactin.

Effect of NAC and ZnSO₄ on sperm parameters of BLCO-exposed Wistar rats:

Results from the present study show that BLCO in a dose-dependent manner significantly decreased sperm quality parameters (sperm count, motility, morphology and viability) when compared with control ($p < 0.05$, Table 2). The findings from the present study are consistent with the result of Farombi et al. (2010) where it was demonstrated that BLCO caused a reduction in epididymal sperm number (ESN), daily spermatozoa production (DSP), and sperm motility. Also, previous studies have observed a reduction in sperm count, motility and abnormal morphology in BLCO-exposed rats (Adesanya et al. 2009, Ita et al. 2020, Fischer et al. 2016). The study also demonstrated that NAC co-administration significantly improved sperm count, motility, viability and sperm morphology) when compared with the administered group ($p < 0.05$, Table 2). Several other studies have highlighted the ability of NAC to improve sperm count, motility and morphology after paracetamol, arsenic and aluminium sulfate-induced oxidative stress and reproductive toxicity in male albino rats (Reddy et al. 2011, El-Maddawy and El-Sayed 2018, Da Silva et al. 2016, Rawi et al. 2015, Oeda et al. 1997). Similarly, the co-administration of BLCO and ZnSO₄, showed significantly improved sperm count, motility, viability and sperm morphology when compared with the BLCO-administered group ($p < 0.05$). Zinc supplementation has been demonstrated to promote the transformation of sperm nuclear protein (i.e., from lysine to arginine) and inhibit the premature depolymerization of the sperm nucleus, improve sperm motility and semen quality in infertile patients without obvious side effects (Als Salman et al. 2018, Rathnayake et al. 2016). It has also been reported that high ZnSO₄ concentrations are correlated with enhanced sperm parameters, including sperm count, motility, and normal morphology (Chia et al. 2000, Eggert-Kruse et al.

2002, Mankad et al. 2016) while lower content of ZnSO₄ has been observed in the seminal plasma of infertile subjects with poor sperm production and poor sperm motility (Zhao et al. 2016). There are several mechanisms by which ZnSO₄ might interfere with sperm function. First, ZnSO₄ is a cofactor for several hundred metalloenzymes, particularly the enzymes responsible for protein synthesis (Stefanidou et al. 2006, Vallee and Falchuk 1993). It influences phospholipases, thus modulating the stability of biological membranes. It has been suggested that the removal of ZnSO₄ from the sperm cell surface destabilizes the plasma membrane, playing an important role in preparation for the completion of capacitation and the acrosome reaction. Some studies have reported that ZnSO₄ supplementation can also improve the synthesis of metallothioneins which protect sperm against damage (Di Leo et al. 2001). It has also been suggested that ZnSO₄ in seminal plasma is involved in maintaining the stability of sperm chromatin (Björndahl and Kvist 2011) and may also exert an *in vitro* effect on oxidative changes in human semen as it is considered a scavenger of excessive O₂ production by defective spermatozoa and/or leukocytes after ejaculation (Gavella et al. 1999). Overall, NAC and ZnSO₄ partially protected against BLCO-induced damage to sperm parameters, with dose-dependent improvements in most cases.

5. CONCLUSION

This study assessed the reproductive function in rats exposed to BLCO and treated with NAC and ZnSO₄. It demonstrated the ability of NAC and ZnSO₄, administered at different doses, to improve and restore reproductive function in these BLCO-exposed rats. The results indicate that NAC and ZnSO₄ are powerful antioxidants that can counteract the harmful effects of oxidative damage caused by ROS generated during BLCO exposure. Additionally, the study showed that NAC and ZnSO₄ protected the testes, preventing reproductive alterations linked to BLCO exposure.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative Artificial intelligence (AI) technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during the writing or editing of this manuscript.

CONSENT

It is not applicable.

ETHICAL APPROVAL

The animals were treated in accordance with the highest ethical standards for animal experimentation. The research protocol and study design were approved by the University of Port Harcourt Research Ethics Committee (UPH/CEREMAD/REC/MM100/056).

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Abdella, A. M., Elabed, B. H., Bakhiet, A. O., Gadir, W. S. A., & Adam, S. E. I. (2011). In vivo study on lead, cadmium and zinc supplementations on spermatogenesis in albino rats.
- Adedara, I. A., Teberen, R., Ebokaiwe, A. P., Ehwerhemuepha, T., & Farombi, E. O. (2012). Induction of oxidative stress in liver and kidney of rats exposed to Nigerian Bonny light crude oil. *Environmental Toxicology*, 27(6), 372-379.
- Adeola, A. O., Akingboye, A. S., Ore, O. T., Oluwajana, O. A., Adewole, A. H., Olawade, D. B., & Ogunyele, A. C. (2022). Crude oil exploration in Africa: Socio-economic implications, environmental impacts, and mitigation strategies. *Environment Systems and Decisions*, 42(1), 26-50.
- Adesanya, O. A., Shittu, L. A. J., Omonigbehin, E. A., & Tayo, A. O. (2009). Spermatotoxic impact of Bonny light crude oil (BLCO) ingestion on adult male Swiss albino mice. *International Journal of Physical Sciences*, 4(5), 349-353.
- Ajonuma, L. C. C., Fapohunda, D. O., Bamiro, S. A., Afolabi, D., Abokede, N., Giwa, K. T., & Oyetunji, S. (2023). Prolonged ingestion of crude oil spill contaminated water affects male reproductive system, hormones, and accessory organs in Wistar rats. *Fertility and Sterility*, 120(4), e130.
- Alabi, O. A., Olukunle, O. F., Ojo, O. F., Oke, J. B., & Adebo, T. C. (2022). Comparative study of the reproductive toxicity and modulation of enzyme activities by crude

- oil-contaminated soil before and after bioremediation. *Chemosphere*, 299, 134352.
- Al-Ani, N. K., Al-Kawaz, U., & Saeed, B. T. (2015). Protective influence of zinc on reproductive parameters in male rat treated with cadmium. *American Journal of Medicine and Medical Sciences*, 5(2), 73-81.
- Al-Hamdany, M. Z., & Al-Hubaity, A. Y. (2014). Protective effects of N-acetylcysteine against 5-fluorouracil-induced pulmonary toxicity in albino rats. *Iraqi Journal of Medical Sciences*, 12(2).
- Alsaman, A. R. S., Almashhedy, L. A., & Hadwan, M. H. (2018). Effect of oral zinc supplementation on the thiol oxidoreductive index and thiol-related enzymes in seminal plasma and spermatozoa of Iraqi asthenospermic patients. *Biological Trace Element Research*, 184, 340-349.
- Alvarez, J. G., & Storey, B. T. (1983). Taurine, hypotaurine, epinephrine and albumin inhibit lipid peroxidation in rabbit spermatozoa and protect against loss of motility. *Biology of Reproduction*, 29(3), 548-555.
- Atkuri, K. R., Mantovani, J. J., Herzenberg, L. A., & Herzenberg, L. A. (2007). N-acetylcysteine—a safe antidote for cysteine/glutathione deficiency. *Current Opinion in Pharmacology*, 7(4), 355-359.
- Awajiomowa, J., Chinko, B. C., & Green, K. I. (2022). Haematological parameters and oxidative stress changes in apparently healthy pregnant women in Bori, Nigeria. *International Journal of Research and Reports in Hematology*, 5(4), 30-39.
- Bablok, L., Dziadecki, W., Szymusik, I., Wołczyński, S., Kurzawa, R. S., Pawelczyk, L., et al. (2011). Patterns of infertility in Poland: Multicenter study. *Neuroendocrinology Letters*, 32(6).
- Bano, A., Gupta, A., Rai, S., Fatima, T., Sharma, S., & Pathak, N. (2021). Mechanistic role of reactive oxygen species and its regulation via the antioxidant system under environmental stress. *Journal Title*.
- Bano, A., Gupta, A., Rai, S., Fatima, T., Sharma, S., & Pathak, N. (2021). Mechanistic role of reactive oxygen species and its regulation via the antioxidant system under environmental stress. In M. Hasanuzzaman & M. K. Nahar (Eds.), *Plant stress physiology - Perspectives in agriculture*. IntechOpen. <https://doi.org/10.5772/intechopen.101045>
- Björndahl, L., & Kvist, U. (2011). A model for the importance of zinc in the dynamics of human sperm chromatin stabilization after ejaculation in relation to sperm DNA vulnerability. *Systems Biology in Reproductive Medicine*, 57(1-2), 86-92.
- Boatman, D. E., Bavister, B. D., & Cruz, E. (1990). Addition of hypotaurine can reactivate immotile golden hamster spermatozoa. *Journal of Andrology*, 11(1), 66-72.
- Boivin, J., Bunting, L., Collins, J. A., & Nygren, K. G. (2007). International estimates of infertility prevalence and treatment-seeking: Potential need and demand for infertility medical care. *Human Reproduction*, 22(6), 1506-1512.
- Chia, S. E., Ong, C. N., Chua, L. L., Ho, L. M., & Tay, S. K. (2000). Comparison of zinc concentrations in blood and seminal plasma and the various sperm parameters between fertile and infertile men. *Journal of Andrology*, 21(1), 53-57.
- Chinko, B. C., & Umeh, O. U. (2023). Alterations in lipid profile and oxidative stress markers following heat stress on Wistar rats: Ameliorating role of vitamin C. *Biomedical Sciences*, 9(1), 12-17.
- Cousins, R. J., Liuzzi, J. P., & Lichten, L. A. (2006). Mammalian zinc transport, trafficking, and signals. *Journal of Biological Chemistry*, 281(34), 24085-24089.
- Da Silva, R. F., Borges, C. d. S., Villela e Silva, P., Missassi, G., Kiguti, L. R. A., Pupo, A. S., et al. (2016). The coadministration of N-acetylcysteine ameliorates the effects of arsenic trioxide on the male mouse genital system. *Oxidative Medicine and Cellular Longevity*, 2016, 4257498.
- Daubry, T. M. E., Chukwuebuka, N. B., Yabrade, T. P., Ehizokhale, E. S., Richard, O. N., Temitope, O. G., & others. (2022). Prenatal restraint stress-induced maternal exposure and Clomifene citrate administration modulates reproductive programming in female Wistar rats. *Pakistan Journal of Biological Sciences*, 25(12), 1066-1076.
- Dede, E. B., & Kagbo, H. D. (2002). A study on the acute toxicological effects of commercial diesel fuel in Nigeria in rats (*Rattus rattus*) using hematological parameters. *Journal of Applied Sciences and Environmental Management*, 6(1), 84-86.
- Di Leo, V., D'inca, R., Barollo, M., Tropea, A., Fries, W., Mazzon, E., et al. (2001). Effect

- of zinc supplementation on trace elements and intestinal metallothionein concentrations in experimental colitis in the rat. *Digestive and Liver Disease*, 33(2), 135-139.
- Eggert-Kruse, W., Zwick, E.-M., Batschulat, K., Rohr, G., Armbruster, F. P., Petzoldt, D., & Strowitzki, T. (2002). Are zinc levels in seminal plasma associated with seminal leukocytes and other determinants of semen quality? *Fertility and Sterility*, 77(2), 260-269.
- Egwurugwu, J. N., Ifedi, C. U., Uchefuna, R. C., Ezeokafor, E. N., & Alagwu, E. A. (2013). Effects of zinc on male sex hormones and semen quality in rats. *Nigerian Journal of Physiological Sciences*, 28(1), 17-22.
- El-Maddawy, Z. K., & El-Sayed, Y. S. (2018). Comparative analysis of the protective effects of curcumin and N-acetyl cysteine against paracetamol-induced hepatic, renal, and testicular toxicity in Wistar rats. *Environmental Science and Pollution Research*, 25, 3468-3479.
- El-Masry, K. A., Nasr, A. S., & Kamal, T. H. (1994). Influences of season and dietary supplementation with selenium and vitamin E or zinc on some blood constituents and semen quality of New Zealand White rabbit males. *World Rabbit Science*, 2(3).
- Farombi, E. O., Adedara, I. A., Ebokaiwe, A. P., Teberen, R., & Ehwerhemuepha, T. (2010). Nigerian Bonny light crude oil disrupts antioxidant systems in testes and sperm of rats. *Archives of Environmental Contamination and Toxicology*, 59, 166-174.
- Fischer, V. A., Fischer, C. E., Akpaso, M., & Igbigbi, P. S. (2016). Effects of ingestion of Bonny Light crude oil on sperm motility and morphology of male Wistar rats. *International Journal of Science and Research*, 5(10), 1230-1233.
- Friedler, S., Razieli, A., Strassburger, D., Schachter, M., Soffer, Y., & Ron-El, R. (2002). Factors influencing the outcome of ICSI in patients with obstructive and non-obstructive azoospermia: A comparative study. *Human Reproduction*, 17(12), 3114-3121.
- Gavella, M., Lipovac, M., Vučić, V., & Šverko, S. (1999). In vitro inhibition of superoxide anion production and superoxide dismutase activity by zinc in human spermatozoa. *International Journal of Andrology*, 22(4), 266-274.
- Georgewill, O. A., & Nwankwoala, R. N. P. (2006). The effects of prolonged exposure to Nigerian crude oil on reproductive hormones of guinea pigs. *African Journal of Applied Zoology and Environmental Biology*, 8, 30-33.
- Gülçin, İ. (2020). Antioxidants and antioxidant methods: An updated overview. *Archives of Toxicology*, 94(3), 651-715.
- Han, J., Won, E.-J., Hwang, D.-S., Shin, K.-H., Lee, Y. S., Leung, K. M.-Y., & others. (2014). Crude oil exposure results in oxidative stress-mediated dysfunctional development and reproduction in the copepod *Tigriopus japonicus* and modulates expression of cytochrome P450 (CYP) genes. *Aquatic Toxicology*, 152, 308-317.
- Indarto, D., & Izawa, M. (2001). Steroid hormones and endocrine disruptors: Recent advances in receptor-mediated actions. *Yonago Acta Medica*, 44(1), 1-6.
- Ita, S. O., Effiong, F. E., Udokang, N. E., Uzoma, J. C., Etim, E. E., & Francis, U. E. (2020). Mitigating toxic effects of Nigerian Bonnylight crude oil on sperm motility and morphology of Wistar rats with ethanol leaves extract of *Ageratum conyzoides* supplementation. *Biomedical Journal of Scientific & Technical Research*, 29(1), 22156-2265.
- Ite, A. E., Ibok, U. J., Ite, M. U., & Petters, S. W. (2013). Petroleum exploration and production: Past and present environmental issues in Nigeria's Niger Delta. *American Journal of Environmental Protection*, 1(4), 78-90.
- Khan, A., Kanwal, H., Bibi, S., Mushtaq, S., Khan, A., Khan, Y. H., & Mallhi, T. H. (2021). Volatile organic compounds and neurological disorders: From exposure to preventive interventions. In *Environmental Contaminants and Neurological Disorders* (pp. 201-230). Springer.
- Khatun, A., Rahman, M. S., & Pang, M.-G. (2018). Clinical assessment of male fertility. *Obstetrics & Gynecology Science*, 61(2), 179-191.
- Khourdaji, I., Zillioux, J., Eisenfrats, K., Foley, D., & Smith, R. (2018). The future of male contraception: A fertile ground. *Translational Andrology and Urology*, 7(Suppl 2), S220.
- Kumar, J., Barhydt, T., Awasthi, A., Lithgow, G. J., Killilea, D. W., & Kapahi, P. (2016). Zinc levels modulate lifespan through multiple

- longevity pathways in *Caenorhabditis elegans*. *PLoS ONE*, 11(4), e0153513.
- Mankad, M., Sathawara, N. G., Doshi, H., Saiyed, H. N., & Kumar, S. (2006). Seminal plasma zinc concentration and α -glucosidase activity with respect to semen quality. *Biological Trace Element Research*, 110, 97-106.
- McLoone, P., Dyussupov, O., Nurtlessov, Z., Kenessariyev, U., & Kenessary, D. (2021). The effect of exposure to crude oil on the immune system: Health implications for people living near oil exploration activities. *International Journal of Environmental Health Research*, 31(7), 762-787.
- Meizel, S. (1985). Molecules that initiate or help stimulate the acrosome reaction by their interaction with the mammalian sperm surface. *American Journal of Anatomy*, 174(3), 285-302.
- Meizel, S., Lui, C. W., Working, P. K., & Mrsny, R. J. (1980). Taurine and hypotaurine: Their effects on motility, capacitation and the acrosome reaction of hamster sperm in vitro and their presence in sperm and reproductive tract fluids of several mammals. *Development, Growth & Differentiation*, 22(3), 483-494.
- Mokhtari, V., Afsharian, P., Shahhoseini, M., Kalantar, S. M., & Moini, A. (2016). A review on various uses of N-acetyl cysteine. *Cell Journal*, 19, 11-17.
- Moschou, M., Kosmidis, E. K., Kaloyianni, M., Geronikaki, A., Dabarakis, N., & Theophilidis, G. (2008). In vitro assessment of the neurotoxic and neuroprotective effects of N-acetyl-L-cysteine (NAC) on the rat sciatic nerve fibers. *Toxicology in Vitro*, 22(1), 267-274.
- Mrsny, R. J., & Meizel, S. (1985). Inhibition of hamster sperm Na^+ , K^+ -ATPase activity by taurine and hypotaurine. *Life Sciences*, 36(3), 271-275.
- Naiho, A. O., Chris-Ozoko, L. E., & Aloamaka, C. P. (2014). The impact of crude oil on reproduction in Wistar rats. *OSR Journal of Pharmacy and Biological Sciences*, 9(3), 1-5.
- Nashwa, A. A. A., Kawkab, A. A., & Mouneir, S. M. (2011). The protective effect of ginger and N-acetyl cysteine on ciprofloxacin-induced reproductive toxicity in male rats.
- Neha, K., Haider, M. R., Pathak, A., & Yar, M. S. (2019). Medicinal prospects of antioxidants: A review. *European Journal of Medicinal Chemistry*, 178, 687-704.
- Noh, S. R., Kim, J.-A., Cheong, H.-K., Ha, M., Jee, Y.-K., Park, M.-S., & others. (2022). Exposure to crude oil-related volatile organic compounds associated with lung function decline in a longitudinal panel of children. *International Journal of Environmental Research and Public Health*, 19(23), 15599.
- Nwafor, A., Asiegbu, L. N., Adienbo, M. O., & Mmom, F. C. (2013). Anti-fertility activity of ingestion of Nigerian Bonny Light crude oil contaminated feed in male rat reproduction: A possible hypothalamo-pituitary axis mechanism.
- Oduwole, O. O., Huhtaniemi, I. T., & Misrahi, M. (2021). The roles of luteinizing hormone, follicle-stimulating hormone and testosterone in spermatogenesis and folliculogenesis revisited. *International Journal of Molecular Sciences*, 22(23), 12735.
- Oeda, T., Henkel, R., Ohmori, H., & Schill, W. B. (1997). Scavenging effect of N-acetyl-L-cysteine against reactive oxygen species in human semen: A possible therapeutic modality for male factor infertility? *Andrologia*, 29(3), 125-131.
- Omirinde, J. O., Olukole, S. G., & Oke, B. O. (2019). Age-related changes in the testicular and epididymal sperm parameters in the African greater cane rat (*Thryonomys swinderianus*, Temminck, 1827). *Animal Research International*, 16(1), 3255-3264.
- Orisakwe, O. E., Akumka, D. D., Afonne, O. J., & Gamanniel, K. S. (2000). Investigation into the pharmacological basis for some of the folkloric uses of Bonny light crude oil in Nigeria. *Indian Journal of Pharmacology*, 32(3), 231-234.
- Orisakwe, O. E., Akumka, D. D., Njan, A. A., & Afonne, O. J. (2004). Testicular toxicity of Nigerian Bonny light crude oil in male albino rats. *Reproductive Toxicology*, 18(3), 439-442.
- Raji, Y., & Hart, V. O. (2012). Influence of prolonged exposure to Nigerian Bonny light crude oil on fertility indices in rats. *Nigerian Journal of Physiological Sciences*, 27(1), 55-63.
- Rathnayake, K. M., Silva, K. D. R. R., & Jayawardena, R. (2016). Effects of zinc supplementation on obesity: Study protocol for a randomized controlled clinical trial. *Trials*, 17, 1-5.
- Rawi, S. M., & Seif Al Nassr, F. M. (2015). Zinc sulphate and vitamin E alleviate

- reproductive toxicity caused by aluminium sulphate in male albino rats. *Toxicology and Industrial Health*, 31(3), 221-234.
- Reddy, P. S., Rani, G. P., Sainath, S. B., Meena, R., & Supriya, C. H. (2011). Protective effects of N-acetylcysteine against arsenic-induced oxidative stress and reprotoxicity in male mice. *Journal of Trace Elements in Medicine and Biology*, 25(4), 247-253.
- Ritchie, G. D., Still, K. R., Alexander, W. K., Nordholm, A. F., Wilson, C. L., Rossi III, J., & Mattie, D. R. (2001). A review of the neurotoxicity risk of selected hydrocarbon fuels. *Journal of Toxicology and Environmental Health, Part B: Critical Reviews*, 4(3), 223-312.
- Sadowska, A. M., Manuel-Y-Keenoy, B., & De Backer, W. (2007). Antioxidant and anti-inflammatory efficacy of NAC in the treatment of COPD: Discordant in vitro and in vivo dose-effects: A review. *Pulmonary Pharmacology & Therapeutics*, 20(1), 9-22.
- Saulsbury, M. D., Heyliger, S. O., Wang, K., & Johnson, D. J. (2009). Chlorpyrifos induces oxidative stress in oligodendrocyte progenitor cells. *Toxicology*, 259(1-2), 1-9.
- Stefanidou, M., Maravelias, C., Dona, A., & Spiliopoulou, C. (2006). Zinc: A multipurpose trace element. *Archives of Toxicology*, 80, 1-9.
- Sule, J. O., Erigbali, P., & Eruom, L. (2008). Prevalence of infertility in women in a southwestern Nigerian community. *African Journal of Biomedical Research*, 11(2).
- Tapiero, H., & Tew, K. D. (2003). Trace elements in human physiology and pathology: Zinc and metallothioneins. *Biomedicine & Pharmacotherapy*, 57(9), 399-411.
- Vallee, B. L., & Falchuk, K. H. (1993). The biochemical basis of zinc physiology. *Physiological Reviews*, 73(1), 79-118.
- Weatherly, L. M., Shane, H. L., Baur, R., Lukomska, E., Roberts, J. R., Fedan, J. S., & Anderson, S. E. (2022). Biological effects of inhaled crude oil. VI. Immunotoxicity. *Toxicology and Applied Pharmacology*, 449, 116100.
- Yu, S., Rubin, M., Geevarughese, S., Pino, J. S., Rodriguez, H. F., & Asghar, W. (2018). Emerging technologies for home-based semen analysis. *Andrology*, 6(1), 10-19.
- Zahra, K. F., Lefter, R., Ali, A., Abdellah, E.-C., Trus, C., Ciobica, A., & Timofte, D. (2021). The involvement of the oxidative stress status in cancer pathology: A double view on the role of the antioxidants. *Oxidative Medicine and Cellular Longevity*, 2021.
- Zhao, J., Dong, X., Hu, X., Long, Z., Wang, L., Liu, Q., et al. (2016). Zinc levels in seminal plasma and their correlation with male infertility: A systematic review and meta-analysis. *Scientific Reports*, 6(1), 22386.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/125280>