



# Effect of Land Configuration and Rainfall on *Pennisetum glaucum* (L.) R. Br. under Saline Irrigation

Muli Naga Surekha <sup>a++</sup>, V. Vasuki <sup>b#\*</sup> and S. Kavitha <sup>c†</sup>

<sup>a</sup> Department of Agronomy, TNAU, Coimbatore – 641 003, Tamil Nadu, India.

<sup>b</sup> Department of Millets, Directorate of Plant Breeding and Genetics, TNAU, Coimbatore – 641 003, Tamil Nadu, India.

<sup>c</sup> Seed Centre, TNAU, Coimbatore – 641003, Tamil Nadu, India.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: <https://doi.org/10.9734/ijecc/2024/v14i114560>

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/125792>

**Original Research Article**

**Received: 24/08/2024**

**Accepted: 27/10/2024**

**Published: 04/11/2024**

## ABSTRACT

Field experiment was conducted in Eastern Block Farm of Tamil Nadu Agricultural University, Coimbatore to evaluate the effect of land configuration and rainfall on growth and yield of pearl millet under saline water irrigation in summer, 2024. The study was carried out in split plot design with three main plots (flatbed (M<sub>1</sub>), ridges and furrows (M<sub>2</sub>) and raised bed paired row planting (M<sub>3</sub>)) along with four subplot treatments (direct sowing (S<sub>1</sub>), 10 Day Old Seedlings (DOS) (S<sub>2</sub>), 15 DOS (modified dapog) (S<sub>3</sub>) and 20 DOS (S<sub>4</sub>)) which was replicated thrice. Water availability is a critical factor in influencing the length of the growing period while water quality significantly influences crop

<sup>++</sup>PG Scholar;

<sup>#</sup>Associate Professor (Agronomy);

<sup>†</sup>Associate Professor (SST);

<sup>\*</sup>Corresponding author: E-mail: [vasuki.v@tnau.ac.in](mailto:vasuki.v@tnau.ac.in);

**Cite as:** Surekha, Muli Naga, V. Vasuki, and S. Kavitha. 2024. "Effect of Land Configuration and Rainfall on *Pennisetum Glaucum* (L.) R. Br. under Saline Irrigation". *International Journal of Environment and Climate Change* 14 (11):466-72. <https://doi.org/10.9734/ijecc/2024/v14i114560>.

growth and yield. Consequently, rainfall received during the growing period affects plant height, Leaf Area Index (LAI), effective tillers and yield. The cumulative rainfall received 208 mm in 22 days and with 10 rainy days during the crop growth period. Rainfall between 40 to 60 Days After Sowing/Planting (DAS/P) corresponded with Crop Growth Rate (CGR). Among main plot treatments, ridges and furrows (M<sub>2</sub>) recorded higher plant height (173 cm) and effective tillers (2.6) at harvest. Grain and stover yields (t ha<sup>-1</sup>) were also higher in ridges and furrows (M<sub>2</sub>) (3.31 and 6.12, respectively) which was statistically on par with raised bed paired row planting (M<sub>3</sub>) (3.31 and 6.12, respectively).

**Keywords:** Crop growth rate; rainfall; raised bed sowing; ridges and furrows; salinity.

## 1. INTRODUCTION

Pearl millet (*Pennisetum glaucum*) is the most extensively cultivated millet. Its resilience to harsh growing conditions, such as drought, poor soil fertility, and high temperature, allows it to thrive in regions where other cereal crops like maize (*Zea mays*) or wheat (*Triticum aestivum*) would fail. The majority of pearl millet production is concentrated in developing countries, which account for over 95% of global production and acreage (Basavaraj et al., 2010). India stands as one of the top millet producers globally, with pearl millet contributing to approximately two-thirds of the nation's total millet output (Singh et al, 2022). The area, production and productivity of India, in 2022-2023 accounts for about 6.84 million hectares, 9.7 million tonnes, 1430 kg ha<sup>-1</sup>. Rajasthan accounts for 44 % of pearl millet produced in India (India stat, 2023).

Crop growth and development are influenced by numerous factors such as the type of soil, irrigation water properties, seeds, and climatic factors like temperature, relative humidity etc. In those factors, irrigation is essential in maintaining yields, mainly in arid regions (Nascimento, 2020). Water is essential for plant growth and development by transporting vitamins, nutrients and other necessary elements from the soil to the plant (Patel et al., 2022). Irrigation management as well as irrigation water quality are equally important for successful crop production (Malash et al., 2005). The main parameters to assess water quality include pH, Electrical Conductivity (EC) and Total Dissolved Salts (TDS). High-quality water can improve seedling growth and yields, whereas poor water quality can slow the growth rates and reduce yields (Chin, 1976).

India is facing a serious problem of shortage of natural assets particularly, water (Gorde, 2013).

Due to this, the use of saline water for irrigation is increasing (Prashanthi et al., 2020) where salts are dissolved in water (Bauder et al., 2007). Guy (2013), stated that irrigation water with excess salinity than the threshold of the crop then yield reduction occurs. As salinity increases the osmotic potential of root zone increases, reducing the water availability to plants (Stewart, 2020). To improve the growth and yield of the crop and reduce the effect of salinity on crop, there is a need to improve agronomic practices in the field (Prashanthi et al., 2020).

Hence, a study was formulated to study the influence of rainfall and land configuration on pearl millet grown under irrigated conditions with saline water.

## 2. MATERIALS AND METHODS

The experiment was conducted at Eastern Block Farm of Department of Agronomy during summer, 2024 at Tamil Nadu Agricultural University, Coimbatore. Pearl millet hybrid COH 10 was used. The experimental site is located in the Western Agro-Climatic Zone of Tamil Nadu, with the co-ordinates of 11°01'00.9" N Latitude, 76° 56'03.9" E Longitude and at an altitude of 426.8 meters above Mean Sea Level. The mean maximum and minimum mean temperature recorded were 34.8°C and 24.8°C, respectively. Wind speed was 5.65 km hr<sup>-1</sup>. A total rainfall of 208 mm was received during crop growing period in 22 days in 10 rainy days. Relative humidity of 81% at 07.22 hr and 49% at 14.22 hr were documented. Mean sunshine hours and solar radiation were 7.0 hr. and 351.6 Cal/cm<sup>2</sup>/day, respectively. The mean evaporation recorded was 6.5 mm. The data on weather parameters that prevailed during the cropping period (April to June) are presented in Fig. 1. The pH and EC of irrigation water were 8.03 (slightly alkaline) and 5.2 (moderately saline), respectively.




Fig. 1. Weather parameters prevailed during the cropping period of Pearl Millet Hybrid COH 10 during Chithirai pattam, Summer (April- June), 2024

In this field study, the split-plot design was replicated thrice. Main plot with three land configurations ( $M_1$ - flatbed,  $M_2$ - ridges and furrow and  $M_3$ - raised bed paired row planting), and subplots with four levels of sowing ( $S_1$ -direct sowing,  $S_2$ -10 DOS,  $S_3$ -15 days old dapog seedlings and  $S_4$ -20 DOS) was followed. The dapog nursery bed was prepared by spreading a polythene sheet of 50-60 gauge on top of the raised bed. Perforations of 1-2 mm diameter were made to drain the excess water from the seedbed. Over the plastic sheet, a mixture of soil (35 kg), well-decomposed FYM (10 kg) and vermicompost (5 kg) (70:20:10) was thoroughly mixed and was spread to a height of 5 cm and properly levelled for uniform distribution of water. Pre-germinated seeds were broadcasted gently without damaging the radicle. For different establishment methods, line sowing of seeds was done along with transplanting 10, 15 and 20 DOS. Normal spacing of 45 cm  $\times$  15 cm and paired row spacing of 60/30 cm  $\times$  15 cm were adopted.

Fertilisation, weeding, irrigation and pest management were done uniformly in all the treatment plots to avoid local control. The five plants in each plot were selected randomly. Selected plants were tagged and subsequently used for recording growth and yield parameters by adopting standard procedures. For calculating grain, stover and biological yield, crop from net plots was selected. Data obtained were statistically analysed in R software (CD at  $P=0.05$ ) to determine the significant difference between the average data of treatments.

### 3. RESULTS AND DISCUSSION

The establishment parameters of pearl millet in various land configurations were significantly different. Establishment percentage and initial plant population lakh  $ha^{-1}$  were observed higher in ridges and furrows ( $M_2$ ) (87.1% and 1.29 lakh  $ha^{-1}$ , respectively) which was statistically on par with paired row raised bed planting ( $M_3$ ) (81.5% and 1.20 lakh  $ha^{-1}$ , respectively). The lower establishment percentage and initial plant population  $ha^{-1}$  was recorded in flatbed ( $M_1$ ) (77.6% and 1.15 lakh  $ha^{-1}$ , respectively). The results were in conformity with Thakare et al., 2019. Higher establishment and population might be due to the raised sowing zone in ridges and furrows ( $M_2$ ) and raised bed paired row planting ( $M_3$ ) which reduced the effect of salinity in irrigation water (Jat et al., 2012). The treatment combination of ridges and furrows and 15 DOS showed better performance.


Plant height and chlorophyll content at harvest was not significantly influenced by the land configurations. Higher plant height (cm) (at harvest) was registered in ridges and furrows ( $M_2$ ) (173) while the lowest among treatments was observed in flatbed ( $M_1$ ) (160). Chlorophyll content at harvest was higher in raised bed paired row planting ( $M_3$ ) (74.0) and lower was recorded in flatbed ( $M_1$ ) (72.2). The effect of different land configurations was significant on LAI. LAI at harvest was higher in raised bed paired row planting ( $M_3$ ) (9.89) which was comparable to ridges and furrows ( $M_2$ ) (9.40). The lowest LAI was recorded in flatbed ( $M_1$ ) (8.52). This might be due to high salt stress in flatbed and a reduced effect of salt was observed in ridges and furrows and raised bed paired row planting (Kang et al., 2010) (Table 1). There was no significant of treatment combinations on growth parameters (plant height, chlorophyll content and LAI).

Crop Growth Rate (CGR) was significantly affected by land configuration and rainfall indicating a correlation between the two factors. From 0-20 DAS/P, higher CGR was recorded in ridges and furrows ( $M_2$ ) (0.004) while the lowest was observed in flatbed ( $M_1$ ) (0.003). From 20-40 DAS/P, higher CGR was registered in ridges and furrows ( $M_2$ ) (14.7) and the next best treatment was raised bed paired row planting ( $M_3$ ) (11). Lowest CGR for this period was found in flatbed ( $M_1$ ) (8.6). During 40-60 DAS/P, ridges and furrows ( $M_2$ ) (17.1) recorded higher CGR whereas lowest CGR was observed in flatbed ( $M_1$ ) (11.1) (Fig. 2). Higher CGR from 40 - 60 DAS/P, might be attributed to rainfall received in that period. Similar results were obtained by Okorogbona et al., 2018, showing that crop irrigated with rainwater performed better than those with groundwater. Treatment combinations did not show significant effect on CGR.

Yield attributes and yield were significantly affected by land configurations. Effective tillers were registered higher in ridges and furrows ( $M_2$ ) and raised bed paired row planting ( $M_3$ ) (2.6) and lowest in flatbed ( $M_1$ ) (2.3). Higher grain yield (t  $ha^{-1}$ ) was observed in ridges and furrows ( $M_2$ ) (3.31) and raised bed paired row planting ( $M_3$ ) (3.31) and poor yield was obtained from flatbed ( $M_1$ ) (2.75). Stover yield (t  $ha^{-1}$ ) was recorded high in raised bed paired row planting ( $M_3$ ) (6.12) which was comparable to ridges and furrows ( $M_2$ ) (6.12) and the lowest was observed in flatbed ( $M_1$ ) (5.49) (Table 1). Enhanced performance of ridges and furrows and

**Table 1. Effect of saline irrigation water on establishment percentage and initial plant population (no.) at 10 DAS/P and other growth and yield parameters (at harvest)**

Treatment	Establishment percentage (%)	Initial plant population lakh ha <sup>-1</sup> (no.)	Plant height (cm) (at harvest)	Chlorophyll content (at harvest)	LAI (at harvest)	Effective tillers (at harvest)	Grain yield (t ha <sup>-1</sup> )	Stover yield (t ha <sup>-1</sup> )
<b>M<sub>1</sub></b>	77.6	1.15	160.8	72.2	8.52	2.3	3.00	5.49
<b>M<sub>2</sub></b>	87.1	1.29	173.4	72.7	9.40	2.6	3.31	6.12
<b>M<sub>3</sub></b>	81.5	1.20	172.7	74.0	9.89	2.6	3.31	6.12
<b>SE(d)±</b>	<b>2.2</b>	<b>0.03</b>	<b>5.3</b>	<b>2.3</b>	<b>0.29</b>	<b>0.1</b>	<b>0.07</b>	<b>0.19</b>
<b>CD</b>	<b>6.0</b>	<b>0.09</b>	<b>NS</b>	<b>NS</b>	<b>0.82</b>	<b>0.2</b>	<b>0.20</b>	<b>0.53</b>
<b>S<sub>1</sub></b>	81.9	1.21	191.8	70.1	9.58	2.5	3.43	6.23
<b>S<sub>2</sub></b>	73.1	1.08	172.7	68.9	9.73	2.4	3.17	5.89
<b>S<sub>3</sub></b>	86.6	1.28	155.2	75.1	8.77	2.5	3.09	5.72
<b>S<sub>4</sub></b>	86.1	1.27	156.2	77.9	9.02	2.6	3.15	5.81
<b>SE(d)±</b>	<b>1.4</b>	<b>0.02</b>	<b>5.1</b>	<b>2.4</b>	<b>0.25</b>	<b>0.1</b>	<b>0.12</b>	<b>0.27</b>
<b>CD</b>	<b>2.9</b>	<b>0.04</b>	<b>10.7</b>	<b>5.1</b>	<b>0.54</b>	<b>NS</b>	<b>0.25</b>	<b>NS</b>
<b>Interaction</b>								
<b>M × S</b>								
<b>SE(d)±</b>	<b>3.0</b>	<b>0.04</b>	<b>9.3</b>	<b>4.3</b>	<b>0.48</b>	<b>0.1</b>	<b>0.19</b>	<b>0.44</b>
<b>CD</b>	<b>6.3</b>	<b>0.09</b>	<b>NS</b>	<b>NS</b>	<b>NS</b>	<b>NS</b>	<b>NS</b>	<b>NS</b>
<b>S × M</b>								
<b>SE(d)±</b>	<b>2.4</b>	<b>0.03</b>	<b>8.8</b>	<b>4.2</b>	<b>0.44</b>	<b>0.1</b>	<b>0.21</b>	<b>0.46</b>
<b>CD</b>	<b>5.0</b>	<b>0.07</b>	<b>NS</b>	<b>NS</b>	<b>NS</b>	<b>NS</b>	<b>NS</b>	<b>NS</b>



**Fig. 2. Correlation between rainfall and CGR in different land configurations (M<sub>1</sub>- Flatbed, M<sub>2</sub>- Ridges and furrows, M<sub>3</sub>- Raised bed paired row planting)**

raised bed paired row planting might be linked to better water management, reduced salt stress and improved soil structure which led to higher plant growth and production (Ming chen et al., 2009). The combination effect of land configuration and methods of establishment on grain and stover yield was insignificant.

Initial establishment was higher in 15 and 20 DOS (S<sub>4</sub>). Among subplots, higher plant height, LAI, grain yield and stover yield were recorded in direct sowing (S<sub>1</sub>) than in transplanted plots. This might be due to the uninterrupted growth cycle and extended vegetative growth. Similar results were recorded by Gudadhe et al., 2020.

#### 4. CONCLUSION

It can be concluded that sowing in raised beds such as ridges and furrows and raised bed paired row planting effectively mitigate the salinity effects of irrigation water. These land configurations resulted in higher plant height, and LAI and DMP compared to flatbed. However, increased rainfall during the crop growing period further enhanced CGR. Yield parameters such as effective tillers were significantly higher in

sowing in raised beds than flatbed as the salts accumulated on top of the raised side of the beds. Notably, the grain and stover yield was increased up to 10% in ridges and furrows and raised bed paired row planting over flatbed. These findings show the importance of a raised sowing zone and optimal rainfall in enhancing crop growth and yield under saline water irrigation conditions.

#### DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

Babu, M. (2020). Impact of saline water on growth, yield, quality, nutrient uptake in

- various crops: A review. *International Journal Conservation Science*, 8, 2344-2347.
- Basavaraj, G., Rao, P. P., Bhagavatula, S., & Ahmed, W. (2010). Availability and utilization of pearl millet in India. *Sorghum and Millet ejournal*, 8.
- Bauder T.A., Waskom R.M., & Davis J.G. (2007). Irrigation water quality criteria. Colorado State University, US Department of Agriculture. *Research Report*, 7(03).
- Chen, M., Kang, Y., Wan, S., & Liu, S. P. (2009). Drip irrigation with saline water for oleic sunflower (*Helianthus annuus* L.). *Agricultural water management*, 96(12), 1766-1772.
- Chin, H.F. (1976). Influence of seed quality on plant growth and development.
- Gorde, S. P., & Jadhav, M. V. (2013). Assessment of water quality parameters: a review. *Journal of Engineering Research and Applications*, 3(6), 2029-2035.
- Gudadhe, Nitin N., Thanki, J. D., Pankhaniya, R. M., & Usdadia, V. P. (2020). "Feasibility of Late Transplanted Summer Pearl Millet for Prolonged Rabi Season with Integrated Nitrogen Management under Indian Coastal Region."
- Guy S. 2013. Know your irrigation water quality. *Irrigation Australia*, 29,12–13.
- Jackson, E. D., Casolaro, C., Nebeker, R. S., Scott, E. R., Dukes, J. S., Griffin, et al. (2023). Current and legacy effects of precipitation treatments on growth and nutrition in contrasting crops. *Agriculture, Ecosystems & Environment*, 352, 108513.
- Kang, Y., Chen, M., & Wan, S. Effects of drip irrigation with saline water on waxy maize (*Zea mays* L. var. ceratina kulesh) in North China Plain (2010). *Agricultural Water Management*, 97(9),1303-1309.
- Malash, N., Flowers, T. J., & Ragab, R. (2005). Effect of irrigation systems and water management practices using saline and non-saline water on tomato production. *Agricultural Water Management*, 78(1-2), 25-38.
- Okorogbona, A. O., Denner, F. D., Managa, L. R., Khosa, T. B., Maduwa, K., Adebola, et al. (2018). Water quality impacts on agricultural productivity and environment. *Sustainable Agriculture Reviews*, 27, 1-35.
- Nascimento, D. M. (2020). A importância da qualidade da água para seu uso na irrigação. *Boletim do Tempo Presente*, 9(1), 70-92.
- Patel, N., Oza, K., & Maitreya, B. (2022). Impact of the different irrigation sources on the plants development: a review. *International Association of Biologicals and Computational Digest*. DOI: 1. 81-84. 10.56588/iabcd.v1i2.48.
- Prashanthi, B., Suresh, K., Ramulu, V., & Sridevi, S. (2020). Growth and Yield attributes of various rice varieties under saline irrigation water as affected by agronomic management practices. *International Journal of Agricultural Statistical Sciences*, 16, 599-606.
- Singh, Y., Tomar, S. S., Yadav, R., Gurjar, R., & Singh, S. (2022). Innovative ridge furrow technology impact on area, productivity and economics of pearl millet: A case study. *Indian Farming*. 72, 67-69.
- Stewart, B. (2020). Irrigation: Saline Water. *Managing Water Resources and Hydrological Systems*, 171-173.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/125792>