



# Review on Modern Breeding and Biotechnological Techniques for Enhancing Breeding Potential of Legume Crops

Padminee Das <sup>a++</sup>, Amit Kumar <sup>b#\*</sup>, Losa Lajeo <sup>ct</sup>,  
Shriram Mahadev Mhaske <sup>dt</sup>, Dhruvkumar C Savaliya <sup>et</sup>,  
Santosh Araganji <sup>ft</sup>, Rajat Singh <sup>g++</sup>  
and Priyanka Solanki <sup>h++</sup>

<sup>a</sup> Department of Plant Breeding and Genetics, SCS College of Agriculture, Assam Agricultural University Rangamati, Chapar, Dhubri- 783371, India.

<sup>b</sup> Department of Genetics and Plant Breeding, G. B. Pant University of Agriculture and Technology, Pantnagar (Uttarakhand), India.

<sup>c</sup> Department of Horticulture (Vegetable Science), School of Agricultural Sciences, Nagaland University, Medziphema Campus (Nagaland)

<sup>d</sup> Vegetable Science, MPKV, Rahuri, India.

<sup>e</sup> Department of Horticulture, (Floriculture and Landscape Architecture), ASPEE College of Horticulture, Navsari Agricultural University, Navsari, India.

<sup>f</sup> Department of Agronomy, UAS, Bangalore – 560065, India.

<sup>g</sup> School of Agriculture, Uttarakhand University, Dehradun -248007, Uttarakhand, India.

<sup>h</sup> Department of Social Science, College of Horticulture and Forestry, Jhalawar, AU, Kota, India.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: <https://doi.org/10.9734/ijpss/2024/v36i115172>

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/126488>

<sup>++</sup> Assistant Professor;

<sup>#</sup> Ph. D. Research Scholar;

<sup>†</sup> Research Scholar;

<sup>‡</sup> M.Sc. Scholar;

\*Corresponding author: E-mail: [singhamitrlbcu001@gmail.com](mailto:singhamitrlbcu001@gmail.com);

**Cite as:** Das, Padminee, Amit Kumar, Losa Lajeo, Shriram Mahadev Mhaske, Dhruvkumar C Savaliya, Santosh Araganji, Rajat Singh, and Priyanka Solanki. 2024. "Review on Modern Breeding and Biotechnological Techniques for Enhancing Breeding Potential of Legume Crops". *International Journal of Plant & Soil Science* 36 (11):587-605. <https://doi.org/10.9734/ijpss/2024/v36i115172>.

## ABSTRACT

Legume crops are critical to global food security and sustainable agriculture, providing essential plant-based proteins and amino acids, while also contributing to soil fertility through symbiotic nitrogen fixation. Despite their nutritional and ecological significance, legume crops production faces numerous challenges, including low yields, susceptibility to biotic and abiotic stresses, and the impacts of climate change on water and land resources. Addressing these issues requires innovative solutions that combine traditional breeding with cutting-edge biotechnological approaches. The recent advancements in legume crops improvement through modern breeding and genome-editing technologies, such as CRISPR/Cas9, TALEN, and ZFN, which enable precise modifications to enhance the suitability and genetic potential of agronomic traits. CRISPR/Cas9, in particular, has emerged as a powerful tool in legume breeding, facilitating targeted mutations, gene knockouts, and gene expression regulation. The review discusses its application across various legume species, including soybeans, cowpeas, chickpeas, and peanuts, to improve traits for example, CRISPR/Cas9 has been used to increase oleic acid content in peanuts and improve photoperiodic flowering in soybeans. Advances in transformation protocols, such as Agrobacterium-mediated methods and biolistic technologies, as well as improvements in tissue culture and phenotyping techniques, are helping to overcome these challenges. Despite significant progress, challenges in legume transformation and regeneration remain, but recent improvements in tissue culture protocols and high-throughput phenotyping have enhanced the efficiency of these genome-editing techniques. It also explores the potential of integrating genome-editing technologies with traditional breeding programs to accelerate genetic gains and develop bio-fortified, climate-resilient legume varieties. By leveraging the vast genetic diversity in legumes and employing advanced genomics tools, researchers can create crops that are not only high-yielding but also nutritionally superior and environmentally sustainable. Integrating genome-editing technologies with traditional breeding paves the way for developing high-yielding, nutritionally enhanced, and climate-resilient legume varieties.

**Keywords:** Legumes; biotechnological; genome editing; CRISPR/Cas9; agrobacterium-mediated transformation.

## 1. INTRODUCTION

“The legume family ranked as the third largest among flowering plants, encompassing over 19,500 recognized species distributed across 751 genera” (Lewis, 2005). “This family included numerous food crops that were vital providers of plant-based proteins and essential amino acids. Leguminous crops were crucial for sustainable agriculture, as they enhanced soil health through symbiotic nitrogen fixation and contributed high-quality organic matter to the soil. Despite their health benefits and ecological importance, legume production faced challenges due to low yields. The reasons of low yield in legume are very from region to region e.g. low productivity is due to weed infestation, poor irrigation management in Chickpea, Urdbean and Mungbean in Bundelkhand region in Uttar Pradesh and Madhya Pradesh while highly

productive land is one of the major cause for poor yield in Punjab, Haryana and Western Uttar Pradesh” (Kumar 2020 and Kumar et al., 2023).

“Addressing these yield issues was complex but could be improved by implementing innovative strategies like genomics-assisted selection (including marker-assisted or genomic selection) and precision breeding” (Bhowmik et al., 2021). Ensuring food security with a focus on high nutrition and yield was anticipated to be a major challenge for researchers and the farming community by 2050.

“Grain legumes were rich in proteins, vitamins, minerals, iron, calcium, zinc, magnesium, and essential omega-3 fatty acids. Soybeans had the highest protein content, ranging from 33% to 45%, followed by common beans (21%–39%), wing beans (30%–37%), cowpeas (21%–35%),

groundnuts (24%–34%), and peas (21%–33%). Other protein-rich legumes included moth beans and urd beans (21%–31%), lentils (20%–31%), grass peas (23%–30%), chickpeas (15%–30%), horse grams (19%–29%), and rice beans (18%–27%) (Pandey et al., 2016). Increasing legume production to desired levels was hindered by biotic and abiotic limitations. Furthermore, the reduction in available land and water resources due to climate change was expected to intensify these adverse conditions in the coming years, placing protein-rich crops like legumes at risk. Eliminating anti-nutritional factors, such as tannins” (Mikic et al., 2009), was a key focus in legume crop improvement programs.

“Efforts were aimed at accelerating genetic gains related to yield, stress tolerance, and nutritional quality. Over the past fifty years, most legume genetic improvements had been achieved through pedigree and performance-based selection. To boost genetic gains more rapidly, innovative genomics techniques and high-throughput phenomics were widely applied, leading to improved legume varieties with valuable agronomic traits” (Varshney et al., 2018). “With growing access to information on genes and haplotypes associated with agronomically important traits, genome editing now enabled the modification of multiple SNPs without altering the original characteristics of popular cultivars. Genetic barriers, such as differences in ploidy levels, often prevented natural genetic exchange between legume species. As a result, the vast genetic diversity found in their wild relatives remained largely untapped” (Varshney et al., 2018).

“The availability of complete genome sequences for various organisms played a crucial role in advancing next-generation genome editing research. Among legumes, there was a greater number of recent genome editing studies focused on *Lotus japonicus*” (Sato et al., 2008), *Glycine max* (Schmutz et al., 2010), and *Medicago truncatula* (Young et al., 2011), all of which had been fully sequenced. This underscored the significance of having complete genome data for facilitating such studies. This article discussed the mechanisms behind new-generation genome-editing technologies, including TALEN (transcription activator-like effector nucleases), ZFN (zinc finger nucleases), and the CRISPR/Cas9 system (clustered regularly interspaced short palindromic repeat/CRISPR-associated protein 9 nuclease), along with detailed examples of their applications

in legume species. Additionally, it provided an in-depth examination of the regulatory framework and the future prospects of genome-editing technologies in these crops. This study served as a comprehensive resource on genome-editing research in plants within the legume family.

A growing number of fields, including plant science and crop breeding, characterization of gene function, and crop enhancement are turning to genetic modification as a cutting-edge and precise strategy of genetic modification. This includes editing to change a specific gene inside an organism’s genome (Ahmad M., 2022, Menz et al., 2020).

## 2. EDITING TECHNOLOGIES

### 1. CRISPR/Cas9

“Recently, CRISPR/Cas9 technology has gained significant popularity for genome editing across a range of organisms, including plants. This advancement has broadened agricultural research by enabling the development of new plant varieties that either eliminate undesirable traits or introduce beneficial characteristics. CRISPR is a fast-evolving method applicable for various genetic modifications, such as creating gene knockouts, making accurate edits, facilitating multiplex genome engineering, and regulating gene expression through activation and repression” (Arora and Narula, 2017). The CRISPR/Cas9 system is composed of two main elements: the Cas9 protein and the guide RNA (gRNA). The Cas9 protein functions as an RNA-guided DNA endonuclease, forming a complex with the gRNA. The gRNA comprises 20 nucleotides complementary to the target DNA sequence and functions as a recruitment signal for Cas9. Unlike other genome-editing technologies, which rely on DNA–protein interactions, CRISPR/Cas9 utilizes RNA–DNA interactions to identify the target DNA sequence. In contrast, ZFNs and TALENs require two distinct DNA-binding domains for each target site due to their use of DNA–protein interactions, making the process more complicated. With CRISPR/Cas9, an 18–20 base pair sequence is all that’s required, since Cas9 and the guide RNA (gRNA) need to bind to a specific protospacer adjacent motif (PAM) located at the 3’ end of the target sequence. The most commonly used PAM sequence for Cas9 from *Streptococcus pyogenes* is 5’-NGG-3’. Cas9 induces DNA repair by creating double-stranded breaks (DSBs) in the target DNA. This repair can occur

through non-homologous end joining (NHEJ) to produce gene knockouts or homology-directed repair (HDR) to facilitate gene modifications and insertions (Huang and Puchta, 2019). “A frameshift mutation takes place when Non-Homologous End Joining (NHEJ) randomly inserts or removes DNA strands within a coding region, leading to a gene knockout. This process does not require a homologous repair template, making NHEJ the most widely used and optimized DNA repair method in plants, including legumes. In contrast, the Homology-Directed Repair (HDR) technique allows for the precise insertion of specific sequences from a donor DNA template. However, HDR has limited application in plants due to its low editing efficiency” (Atkins and Voytas, 2020). CRISPR technology offers various applications, such as gene expression analysis, and gain- or loss-of-function studies. CRISPR has also been applied in legume agriculture, and this genome-editing tool holds potential for creating sustainable, high-quality agricultural products, including legumes.

### **A. gRNA Design and Validation**

The guide RNA (gRNA) is a critical element of the CRISPR/Cas9 system, determining the specificity and efficiency of gene editing. In plants, the presence of large gene families with high sequence similarity increases the risk of off-target binding by the gRNA, which can lead to unintended effects. Therefore, it is essential to design gRNAs that offer high on-target accuracy while minimizing off-target interactions. For species with low transformation rates and extended tissue culture processes, it is advantageous to select highly active and multiple gRNAs. Conversely, in species that are more easily transformable, approximately two gRNAs per gene are generally sufficient to achieve a mutant.

### **B. gRNA delivery**

#### **i) Transformation of legume crops**

“For successful transformation, including CRISPR/Cas9 gene editing, it is essential to deliver DNA/RNA components and regenerate a whole plant from transformed tissues. Legumes are notably challenging in this regard, as they tend to resist both DNA uptake and integration” (Yadav et al., 2017) and are generally difficult to regenerate. Complicating matters further, the tissues that are amenable to transformation and those capable of regeneration do not always

overlap in legumes. This highlights the need for a transformation protocol that integrates the transformation vector from the start, rather than merely focusing on developing a regeneration protocol.

Large-seeded legumes primarily utilized *Agrobacterium*-mediated transformation, though biolistics technologies were also applied. Successful transformations generally involved explants from seed tissues, including whole or partially developing or mature embryos, as well as cotyledonary nodes, sometimes including deembryonated cotyledons. A Common treatments for explants involved wounding, often paired with acetosyringone to activate *Agrobacterium* virulence genes. Wounding methods included puncturing, cutting embryos in half, or making longitudinal slices. Additional techniques to enhance DNA uptake involved sonication and/or vacuum infiltration, along with other compounds such as lipoic acid.

Transformation protocols typically consisted of several phases. For organogenesis in species like pea and chickpea, these phases included a co-cultivation period, shoot initiation, shoot elongation, and rooting, each requiring distinct media formulations. Protocol development often utilized a reporter gene like *gus* ( $\beta$ -glucuronidase) to monitor progress. Although using a reporter gene required sacrificing some tissue, it was often easier to detect than fluorescence in green, photosynthetic tissues. Regardless of the reporter method, molecular analysis techniques such as PCR or ddPCR were essential to confirm the successful regeneration of a transgenic plant.

Protocols utilizing embryogenesis typically included stages such as embryogenesis, proliferation of embryonic tissue for bombardment, maturation, desiccation, and germination. Media formulations, incubation conditions, and durations varied significantly among different species and protocols.

#### **ii) CRISPR/Cas9 mutagenesis in doubled haploids**

A functional microspore-based gRNA delivery system presents several advantages for rapidly achieving homozygosity and shortening the generation cycle. It allows for the quick isolation of a large number of genetically identical and physiologically uniform microspores, facilitating functional gene validation, genetic

characterization, and the enhancement of valuable traits. Additionally, fluorescent reporter systems enable the high-throughput screening of multiple gRNAs simultaneously using microspores. One of the initial experiments involving microspore mutagenesis with the CRISPR/Cas9 system was conducted by Bhowmik et al. (2018), who utilized a Neon electroporation system to deliver gRNAs into isolated wheat microspores. Furthermore, Bilichak et al. (2020) demonstrated that a complex of cell-penetrating peptides and zinc-finger nucleases (CPP-ZFN) could effectively introduce edits in haploid embryo-like structures regenerated from wheat microspores.

Microspores from legume species held potential for gRNA delivery and CRISPR/Cas9-mediated gene editing. However, the recalcitrant nature of these legumes significantly hindered the production of double haploid (DH) plants. Various research groups attempted to regenerate DH plants (Bobkov, 2010; Gupta, 1975; Lulsdorf et al., 2011; Ochatt et al., 2009; Ribalta et al., 2012), but the frequency of successfully regenerating complete haploid plants remained very low (Bobkov, 2014; Ochatt et al., 2009). To date, successful regeneration of DH plants had only been achieved in chickpea through anther culture and androgenesis.

Recognizing the importance of double haploid (DH) technology in pulse breeding, several laboratories, including the National Research Council of Canada, have recently initiated efforts to re-evaluate and enhance strategies for producing DH plants. These advancements aim to overcome previous challenges and improve the efficiency and success rates of DH plant production in legumes.

### 3. DETECTION OF GENETIC MODIFICATIONS

Identifying and monitoring modifications caused by genome editing were crucial at various stages, including the optimization of experimental conditions, characterization, selection, and tracking of edited lines. The methods available for detecting these modifications were diverse and continually evolving, with no single method meeting all genotyping requirements.

Key challenges in reliably detecting modifications stemmed from the biological characteristics of gene-editing proteins, which could induce a variety of possible mutations, as well as the

biological traits of the target organism, which might involve multiple homologs and varying efficiencies in creating edited lines. Given the difficulties associated with regenerating legumes, it was often necessary to screen a large number of lines or heterogeneous tissues, emphasizing the need for methods that were both cost-effective and sensitive.

Although most economically important legumes were not complicated by recent polyploidization, they exhibited a higher propensity for gene duplications (Young et al., 2011). High-quality genome assemblies played a crucial role in the reliable interpretation of detection experiments and were especially important for assessing off-target effects.

Therefore, developing and utilizing detection methods that ensured high on-target accuracy while minimizing off-target interactions was essential, particularly in species with low transformation rates and lengthy tissue culture processes. In this section, we discuss some of the most widely used and promising methods for detecting edited sequences in legumes. For a more comprehensive comparison of established methodologies, we refer the reader to a recent review dedicated entirely to this topic.

#### 3.1 Mismatch Detection

One of the original and most popular methods for screening mutations involves detecting mismatches formed during the denaturation and annealing of heterogeneous DNA populations. Several endonucleases, such as *CelI*, that can cleave at mismatches have been employed for this purpose. While *CelI* may excel in single nucleotide polymorphism (SNP) detection compared to *T7EI*, the latter is recommended for CRISPR/Cas9 experiments, as it tends to perform better on insertions and deletions (indels) and generally produces less background signal, which can be a limiting factor for this approach. Although mismatch detection is relatively easy to implement and requires only standard molecular biology equipment, it is limited by low sensitivity and high rates of both false positives and negatives.

#### 3.2 Electrophoretic Separation

Electrophoretic mobility was used to directly provide a simpler method for identifying mutants. Large insertions or deletions were easily monitored through PCR followed by agarose gel

electrophoresis, while PCR coupled with nondenaturing polyacrylamide gel electrophoresis (PAGE) was able to detect indels as small as 1 bp. Although these methods are quick, they have limited sensitivity for detecting low-abundance edits, making them more suitable for systems with high editing efficiency or for screening later generations to identify homozygous lines.

When targeting a restriction enzyme recognition site is feasible, restriction enzyme-suppressed PCR (RE-PCR) offers a more sensitive alternative (Xie & Yang, 2013). This method involves conducting a restriction digest before PCR amplification, allowing only the mutated sites to be amplified.

Annealing at critical temperature PCR (ACT-PCR) is a specialized PCR strategy designed to detect homozygous mutant lines by exploiting the annealing characteristics of primers in relation to the presence of mutations. Key following points:

**1. Principle-** ACT-PCR relies on the concept that primers will have different annealing efficiencies based on the presence or absence of mutations in the template DNA. The strategy involves finding the highest possible annealing temperature at which a primer can anneal to the wild-type template but fails to anneal when a mutation is present.

**2. Detection of Mutations-** By adjusting the annealing temperature, it is possible to ensure that the primer overlaps the target site, thereby creating conditions where it will successfully anneal to the wild-type sequence while failing on the edited (mutated) sequence. This results in no amplification of the target from the mutant template.

**3. Internal Positive Control-** To enhance the reliability of the assay, an additional competing flanking primer can be included, which serves as an internal positive control. This primer should amplify a region that is not affected by the mutation, ensuring that the PCR reaction is working properly.

**4. Equipment Requirements-** While the method has minimal equipment requirements, utilizing a gradient thermocycler is recommended. This allows for precise identification of the critical annealing temperature, enabling researchers to fine-tune the PCR conditions for optimal results.

**5. Applications-** ACT-PCR can be a powerful tool in genetic studies, particularly for identifying

homozygous mutations in plant breeding, genetic engineering, and crop improvement programs.

This method represents a valuable approach in molecular biology, particularly for detecting specific genetic changes with high sensitivity and specificity.

Incorporating fluorescence-based detection methods can significantly enhance the sensitivity of PCR techniques. The application of fluorophore-labeled primers has been shown to produce labeled amplicons that can be detected using DNA capillary electrophoresis. This approach allows for the detection of insertions and deletions (indels) with a limit of detection (LOD) of approximately 0.1%, and it can differentiate mutation populations with a resolution of 1 to 2 base pairs. Additionally, this method capitalizes on the automation capabilities of capillary electrophoresis equipment. Although not yet demonstrated, it has the potential to simultaneously monitor multiple targets by using different fluorophores for each amplicon, similar to the high-throughput analysis of simple sequence repeats (SSRs). The method's combination of low cost, high sensitivity, and scalability makes it an appealing choice for screening in legumes, especially in settings where quick turnaround times are feasible due to available instrumentation.

## 4. FLUOROPHORE BINDING-BASED DETECTION

Various highly sensitive techniques for detecting edits were employed, including high-resolution melt analysis (Thomas et al., 2014), quantitative PCR (qPCR) (Peng et al., 2018), and droplet digital PCR (ddPCR) (Miyaoaka et al., 2018), which utilized fluorescent probes or dyes for binding. Although these methods were both sensitive and cost-effective, they required access to specific equipment.

### 4.1 Sanger Sequencing

Sanger sequencing is an affordable and reliable technique for genotyping that can verify modifications. Traditionally, Sanger sequencing data analysis required clonal samples, but newer methods like TIDE (Tracking of Indels by Decomposition; Brinkman & van Steensel, 2019) and ICE (Inference of CRISPR Edits; Hsiao et al., 2018) now allow for the interpretation of Sanger sequencing data from mixed populations of DNA fragments. This capability, combined with

its low cost and speed, has made Sanger sequencing popular in plant research. However, it has limitations, such as a moderate limit of detection (LOD) and low tolerance for off-target amplification. While the LOD can be as sensitive as 1–2% (Brinkman et al., 2014) depending on the length of indels, in practical applications, the LOD for these Sanger-based methods is typically around 10%.

## 4.2 Next-Generation Sequencing

Massively parallel sequencing technologies enabled the simultaneous examination of numerous sites across multiple samples. Next-generation sequencing (NGS) techniques were highly sensitive (with a limit of detection, LOD, of less than 1%, as noted by Kim et al., 2019) and could characterize the range of edits within a cell population. Sample preparation for NGS involved attaching sequencing adapters to the target sequences. For Illumina platforms, a common method for preparing amplicon libraries was to add a partial adapter sequence as a tail on the target-specific primers. A second PCR then completed the adapter sequence and added sample-specific barcodes. This method allowed the use of universal indexing primers, facilitating the pooling of multiple amplicons from a single sample after the initial PCR step. Generic indices became commercially available with up to 384 unique dual-index combinations, enabling greater scalability in multiplexing. Additionally, it was possible to multiplex the initial target-specific PCR step, further simplifying the preparation of numerous amplicons per sample. Various tools, like Oligo (Hending et al., 2018), supported multiplex PCR design, and methods such as rhAmp-seq from IDT could handle multiplexing for over 100 targets.

While there are currently no established methods for utilizing Oxford Nanopore Technologies for detecting edits, the smaller MinION or Flongle platforms are appealing for their quick turnaround capabilities. Despite the high error rates associated with this platform, both the technology and its application strategies are advancing swiftly. One approach to manage these higher error rates involves adding unique molecular identifiers (UMIs) to amplicons (Karst et al., 2021). Although this adds an extra step to the process, it significantly enhances the reliability and sensitivity in detecting mutations.

Although per-experiment costs remain relatively high compared to other strategies mentioned here, the extensive data output and multiplexing

capability of NGS methods offer the potential for reduced costs per sample. NGS of genomic libraries is also currently the only method that can provide complete genome coverage, allowing for a thorough examination of genomic alterations (Tang et al., 2018). Despite its powerful capabilities, NGS approaches often have longer turnaround times and require batching to fully capitalize on their advantages. NGS methods yield information for individual library fragments, which offers some tolerance for off-target amplification; however, off-target sequences and discrepancies between the experimental line and the reference genome are primary contributors to false positives. As a result, incorporating negative controls is essential for accurately interpreting variant calling data. We anticipate that NGS screening methods will continue to gain popularity as editing experiments advance towards higher multiplexity, and as the efficiency of NGS experiments continues to improve.

While there were no established methods for utilizing Oxford Nanopore Technologies to detect edits, the smaller MinION or Flongle platforms were appealing for their quick turnaround capabilities. Despite the high error rates associated with this platform, both the technology and its application strategies were advancing swiftly. One approach to manage these higher error rates involved adding unique molecular identifiers (UMIs) to amplicons (Karst et al., 2021). Although this added an extra step to the process, it significantly enhanced reliability and sensitivity in detecting mutations.

Although per-experiment costs remained relatively high compared to other strategies mentioned, the extensive data output and multiplexing capability of NGS methods offered the potential for reduced costs per sample. NGS of genomic libraries was also the only method that could provide complete genome coverage, allowing for a thorough examination of genomic alterations (Tang et al., 2018). Despite its powerful capabilities, NGS approaches often had longer turnaround times and required batching to fully capitalize on their advantages.

NGS methods yielded information for individual library fragments, offering some tolerance for off-target amplification; however, off-target sequences and discrepancies between the experimental line and the reference genome were primary contributors to false positives. As a result, incorporating negative controls was

essential for accurately interpreting variant calling data. It was anticipated that NGS screening methods would continue to gain popularity as editing experiments advanced towards higher multiplexity, and as the efficiency of NGS experiments continued to improve.

## 5. APPLICATIONS IN LEGUMES

### 1. *M. truncatula* (Alfalfa)

*Medicago truncatula* is widely used as a model organism for legume crops because of its short life cycle, ease of genetic transformation, self-fertility, diploid genome, and relatively small genome size. These features make it a preferred option for conducting molecular and physiological research on legumes. In a study by Michno et al. (2015), mutations were induced in the soybean genes for glutamine synthase (GS1) and chalcone-flavanone isomerase (CHI20) in *Glycine max*, as well as in the  $\beta$ -glucuronidase (GUS) gene in *M. truncatula* using hairy root transformation.

Bottero et al. (2021) generated two transgenic alfalfa lines, designated as 3-1 and 5-1, using the CRISPR/Cas9 system with a pBI121 binary vector that included the GUS gene. This resulted in an average GUS inactivation of 55%. In the scientific literature, phytoene desaturase (PDS) genes were often chosen for study due to the ease of observing phenotypic changes, which made it simpler to evaluate the effectiveness of CRISPR/Cas9 gene editing. For example, Meng et al. (2017) established an effective CRISPR/Cas9 system to induce mutations in the MtPDS gene in *M. truncatula*, finding that 32 of 309 T0 transgenic plants displayed an albino phenotype. Sequencing 16 randomly selected albino plants confirmed that all had mutations at the targeted MtPDS gene site. Additionally, Wolabu et al. (2020) discovered that the UBQ10 promoter-driven Cas9 resulted in high mutation efficiency (95% in *Arabidopsis* and 70% in *M. truncatula*). Similarly, Zhang et al. (2020) targeted MtPDS genes with CRISPR/Cas9, and all homozygous/biallelic MtPDS mutant seedlings showed albino traits.

The CRISPR/Cas9 gene-editing tool was also applied in studies focusing on flowering and secondary metabolite production. Rodas et al. (2021) used CRISPR/Cas9 to edit the SUPERMAN (MtSUP) gene in *M. truncatula*, which disrupted MtSUP function, leading to defects in floral development and inflorescence structure in plants carrying the mtsup mutant

allele. Similarly, Confalonieri et al. (2021) utilized CRISPR/Cas9 to knock out two cytochrome P450 genes, CYP93E2 and CYP72A61, which were involved in the production of soyasapogenol B in *Medicago* species. Their findings revealed that 51 putative CYP93E2 mutant plant lines, with an 84% editing efficiency, did not produce soyasapogenols in leaves, stems, or roots. Instead, this alteration shifted the metabolic pathway towards producing valuable hemolytic sapogenins.

### 2. *Japonicus* (Lotus)

*L. japonicus* served as a model organism for legume crops and shared similarities with *M. truncatula*. However, unlike *M. truncatula*, *L. japonicus* formed determinate nodules, similar to soybean (*G. max*) and cowpea (*Vigna unguiculata*). Wang et al. (2016) demonstrated that gene mutations related to symbiotic nitrogen fixation (SNF) were achievable in *L. japonicus* using CRISPR/Cas9 through hairy root transformation. In 2018, Cai et al. edited the cytokinin receptor Lotus histidine kinase I-interacting protein (LjCZF1) to investigate cytokinin signaling involved in rhizobia-legume symbiosis. They found that knock-out mutants had significantly fewer infection threads and nodules, indicating that LjCZF1 positively regulated symbiotic nodulation. Subsequently, Wang et al. (2019) used CRISPR/Cas9 to explore the role of leghemoglobin (Lbs) in *L. japonicus*, discovering that the absence of Lbs led to early nodule senescence. In another study, a loss-of-function analysis of the CYP716A51 gene, involved in triterpenoid C-28 oxidation, showed that cyp716a51 mutant *L. japonicus* hairy roots were unable to produce C-28 oxidized triterpenoids.

### 3. *G. max* (Soybean)

The growing demand for soybean products, due to their high economic value as a protein and oil source for both animal and human nutrition, has highlighted the need to accelerate genetic advancements using gene-editing technologies. Bao et al. (2020) emphasized that these advancements are essential to meet rising demands and address environmental challenges. Curtin et al. (2011) initially optimized a gene-editing approach in soybean by targeting the green fluorescent protein (GFP) coding region with a zinc finger nuclease (ZFN) array, developed using context-sensitive selection strategies. This approach resulted in deletions of up to 71 base pairs at the target



site. With further optimization, they subsequently targeted two distinct RNA-dependent polymerases in soybean.

The study's most intriguing result was the design of two independent ZFN pairs, both of which successfully recognized their specific targets and caused two base pair differences in both genes. This demonstrated the precise targeting and effectiveness of the ZFN approach in gene editing. In another study by Curtin et al. (2015), researchers investigated the disruption of miRNA maturation and the regulation of miRNA gene expression. They developed two distinct ZFN pairs aimed at the Dicer-like 1a (DCL1a) and Dicer-like 1b (DCL1b) genes in soybean. Although single mutants of the DCL genes did not yield significant results, the double mutants of the DCL genes exhibited notable morphological changes. Additionally, these double mutants showed impaired processing efficiency of miRNA precursor transcripts and deregulated expression of miRNA target genes. In 2018, Curtin et al. employed TALENs to target the *G. max* Dicer-like2 gene and used whole-genome sequencing to reveal multiple transgene insertion events, producing a range of combinatorial mutant plants. With CRISPR-Cas, they targeted two endogenous soybean genes (GmFEI2 and GmSHR), designing sgRNAs against a transgene (*bar*) and creating targeted DNA mutations in hairy roots using six sgRNAs. Li et al. (2015) successfully utilized CRISPR/Cas9 to induce mutations at two genomic sites, DD20 and DD43, on chromosome 4 with success rates of 59% and 76%, respectively. Sun et al. (2015) further developed two vectors with *Arabidopsis* U6-26 and soybean U6-10 promoters to target Glyma06g14180, Glyma08g02290, and Glyma12g37050 in protoplasts. Biallelic mutations were observed in Glyma06g14180 and Glyma08g02290 within transgenic hairy roots.

Additionally, a comparative analysis of CRISPR/Cas9 and TALENs for editing GmPDS11 and GmPDS18 genes revealed albino and dwarf phenotypes (PDS knock-outs) following cotyledon node transformations. TALENs demonstrated a slightly higher mutation efficiency than CRISPR/Cas9 with the AtU6-26 promoter, though it was lower when using the GmU6-16g-1 promoter in hairy roots. The study concluded that both technologies effectively achieved gene targeting in soybean. Curtin et al. (2018) also used CRISPR/Cas9 and TALENs in parallel on *G. max* and *M. truncatula*, creating a bi-allelic double mutant for the soybean

GmDrb2a and GmDrb2b genes and a mutation in the *M. truncatula* Hua enhancer1 (MthEn1) gene. Soybean flowering time is crucial for accelerating breeding processes to enhance yield and quality. Consequently, various studies have focused on editing genes associated with flowering time. Cai Y. et al. (2018) developed an efficient dual-sgRNA/Cas9 system to induce deletions in the GmFT2a and GmFT5a genes. They achieved deletion frequencies of 15.6% and 15.8% in GmFT2a and GmFT5a, respectively, with deletions over 4.5 kb occurring in 12.1% of GmFT2a samples. These deletions were inheritable in T2 transgene-free homozygous *ft2a* mutants, which displayed a late-flowering phenotype. In a separate study, Han et al. (2019) edited the soybean maturity gene E1, responsible for flowering regulation, producing 11 bp and 40 bp deletions that resulted in premature stop codons and truncated E1 proteins. Additionally, CRISPR/Cas9 was used to generate knock-out and overexpression mutations in the GmPRR37 gene, which encodes the flowering time regulator qFT12-2, demonstrating its role in controlling photoperiodic flowering in soybean. Another study examined GmFT2a and GmFT5a genes using CRISPR/Cas9 and showed that these genes jointly regulate flowering time by analyzing *ft2a*, *ft5a*, and *ft2a/ft5a* mutants under short-day and long-day conditions.

The CRISPR/Cas9 strategy has also been used to target three GmLox genes (GmLox1, GmLox2, and GmLox3), which encode lipoxygenases (LOX1, LOX2, and LOX3) that contribute to a beany flavor limiting soybean's appeal for human consumption. In this study, 60 T0 positive transgenic plants were generated with combinations of sgRNAs, resulting in various mutations, including two triple mutants and one double mutant, all of which lost the corresponding lipoxygenase activities. Similarly, Li et al. (2019) applied CRISPR/Cas9 to edit the conglycinin (7S) and glycinin (11S) storage protein genes, achieving editing efficiencies of 5.8%, 3.8%, and 43.7% for Glyma.20g148400, Glyma.03g163500, and Glyma.19g164900, respectively. Furthermore, CRISPR/Cas9 has been used to modify plant architecture in soybean by targeting squamosa promoter binding protein-like genes (GmSPL9a, GmSPL9b, GmSPL9c, and GmSPL9). T2 double homozygous *spl9a/spl9b* mutants displayed a shorter plastochron length and, in T4 mutant plants, increased main stem node numbers and branch numbers were observed.

**Table 1. Gene-editing technology in different legume crops**

<b>Legume</b>	<b>Technique</b>	<b>Target</b>	<b>Legume</b>	<b>Technique</b>
<i>Medicago truncatula</i>	CRISPR/Cas 9	<i>MtSUP</i> (regulates the floral organ number)	<i>MtSUP</i> was found to be orthologous of <i>AtSUP</i>	Rodas et al. 2021
	CRISPR/Cas 9	CYP93E2 and CYP72A61 (soyasapogenol B biosynthesis)	51 CYP93E2 mutant plant lines	Confalonieri et al.
	CRISPR/Cas 9	GmGS1, GmCHI20, MtGUS	Mutated genes	Michno et al.
	CRISPR/Cas 9	MtHen1 (Hua enhancer1 gene)	Efficient mutation	Curtin et al. (2018)
<i>Lotus japonicus</i>	CRISPR/Cas 9	Lbs genes (nodule senescence)	Early nodule senescence	Wang et al. (2019)
	CRISPR/Cas 9	SNF (symbiotic nitrogen fixation) genes	CRISPR/Cas9 system can effectively induce mutations in SNF related genes	Wang et al. (2016)
	CRISPR/Cas 9	LjCZF1 and LjCZF2 (root nodule symbiosis)	Decrease in nodule formation	Cai, K., (2018). et al.
Glycine max (Soybean)	CRISPR/Cas 9	GmPRR37 (photoperiodic flowering)	Changes in flowering time	Wang, L., et al.(2020)
	CRISPR/Cas 9	GmLox1, GmLox2, GmLox3 (encoding lipoxygenases)	Loss of lipoxygenase activity	Wang, J., et al.
	CRISPR/Cas 9	GmAGO7a and GmAGO7b (controlling leaf pattern)	Inherited mutation until T2 lines	Zheng et al. (2020)
	TALEN	<i>FAD2-1A/B</i> (seed content improvement)	Increased oleic acid content, reduced linolenic acid content	Haun et al. (2014)
Cowpea ( <i>Vigna unguiculata</i> )	CRISPR/Cas 9	VuSPO11-1 (cowpea meiosis gene)	Mutations	Che et al. (2021)
	CRISPR/Cas 9	SNF (symbiotic nitrogen fixation) genes	Blocked nodule formation	Ji et al. (2019)
Chickpea ( <i>Cicer arietinum</i> )	CRISPR/Cas 9	<i>4CL</i> (4-coumarate ligase) <i>RVE7</i> ( <i>Reveille 7</i> ) (drought tolerance)	High efficiency in editing	Badhan et al.
Peanut ( <i>Arachis hypogaea</i> )	CRISPR/Cas 9	<i>AhNFR1</i> and <i>AhNFR5</i> (nodulation)	Successfully edited genes	Shu et al. (2020)
	CRISPR/Cas 9	<i>AhFAD2</i> (seed content improvement)	G448A, 441_442insA, G451T mutations	Yuan et al. (2019)
	CRISPR/Cas 9	<i>AhFAD2</i> (seed content improvement)	G448A, 441_442insA, G451T mutations	Yuan et al. (2019)
	TALEN	<i>AhFAD2</i> (seed content improvement)	Increase in the oleic acid content	Wen et al. (2018)

Improving the oleic acid content in soybeans has also become a significant breeding objective. Gene-editing technologies have played a crucial role in this area. While TALENs have been less frequently used in legumes, successful applications have targeted the Fatty Acid Desaturase 2 (FAD2) genes, which convert oleic acid to linoleic acid, to increase oleic acid levels. Haun et al. (2014) targeted the FAD2-1A and FAD2-1B genes, using four TALEN pairs; two pairs (FAD2\_T01 and FAD2\_T04) were expressed, with FAD2\_T04 achieving a 7.2% mutation rate. As a result, they observed a decrease in linoleic acid (down to 4%) and an increase in oleic acid content (up to 80%). Demorest et al. (2016) conducted a similar study, targeting the FAD2-1A, FAD2-1B, and FAD3A genes, continuing the effort to enhance oleic acid content through targeted mutagenesis.

TALENs were designed to target the FAD3A gene in lines already mutated at FAD2-1A and FAD2-1B, achieving mutation rates of 11.2%, 16.0%, and 4.9%, respectively. These mutations led to an increase in oleic acid content of over 80% and a reduction in linoleic acid to 2%. Do et al. (2019) also targeted the GmFAD2-1A and GmFAD2-1B genes, generating T0 transgenic plants that displayed an 80% increase in oleic acid content and a 1.3%–1.7% decrease in linoleic acid in T1 seeds homozygous for both genes. Similarly, Al Amin et al. (2019) used CRISPR-Cas9 to mutate the FAD2-2 gene, resulting in significant changes in the oleic acid/linoleic acid ratios due to frequent deletions and insertions.

In 2020, Wu and colleagues applied CRISPR/Cas9 to GmFAD2-1A and GmFAD2-2A genes, producing single and double knock-out mutants. They found editing efficiencies of 95% for GmFAD2-1A, 55.56% for GmFAD2-2A, and 66.67% for double mutants. These mutants exhibited an increase in oleic acid content up to 73.5% and a decrease in linoleic acid down to 12.23% in the T2 generation, with similar trends continuing into the T3 generation. Di et al. (2019) enhanced CRISPR/Cas9 efficiency by utilizing highly active U6 promoters to target Glyma03g36470, Glyma14g04180, and Glyma06g136900, which resulted in insertion, deletion, and substitution mutations. In the following year, they constructed 70 CRISPR-Cas9 vectors to target 102 candidate genes, creating 407 T0 mutant lines with a mutagenesis frequency of 59.2%, of which 35.6% had

multiplex mutations. These mutants showed increased nodule numbers in gmric1/gmric2 double mutants, while gmrnd1-1/1-2/1-3 triple mutants exhibited reduced nodulation.

#### 4. *V. unguiculata* (Cowpea)

Cowpea (*Vigna unguiculata* (L.) Walp.) is recognized for its high nutritional value and health benefits. It possesses efficient symbiotic nitrogen fixation (SNF) capabilities, tolerance to low rainfall, and minimal fertilization needs, making it one of the most significant legumes globally. Consequently, there has been growing interest in employing gene-editing techniques in cowpea.

In 2019, Ji and colleagues demonstrated CRISPR/Cas9-mediated genome editing in cowpea by targeting SNF genes, leading to non-inheritable mutations in hairy roots. They found that nodule formation was entirely blocked in mutants with disruptions in both alleles. Following this, Juranic et al. (2020) identified three cowpea meiosis-related genes: SPO11-1 (which encodes the SPO11 protein responsible for initiating meiotic double-stranded breaks), REC8 (which encodes a meiotic recombination protein), and OSD1 (which encodes Ophiostoma scytalone dehydratase, promoting meiotic progression). They used CRISPR/Cas9 to induce asexual seed formation, resulting in biallelic mutations in exons 1 and 3 of the SPO11-1 gene, causing complete male and female sterility in T0 plants. More recently, Che et al. (2021) knocked out the cowpea meiosis gene VuSPO11-1 using CRISPR/Cas9 and confirmed mutations at the target site.

This work underscored the potential of CRISPR/Cas9 technology to advance genetic research in cowpea, addressing both fundamental biological questions and practical agricultural challenges.

#### 5. *C. arietinum* (Chickpea)

Chickpea (*Cicer arietinum*) is an important commercial crop globally, and gene-editing tools offer potential solutions to challenges in its production. In a study by Badhan et al. (2021), the researchers targeted drought tolerance-related genes, specifically 4-coumarate ligase (4CL) and Reveille 7 (RVE7), using CRISPR/Cas9 editing in chickpea protoplasts. The knockout of the RVE7 gene resulted in highly efficient editing in vivo. These findings

demonstrated that DNA-free CRISPR/Cas9 gene editing can be effectively applied to drought tolerance genes in chickpea protoplasts. To date, this study represents the first and only application of CRISPR/Cas9 gene editing in chickpea.

### 6. A. *hypogaea* (Peanut)

Peanut (*Arachis hypogaea*) is a significant legume crop known for its high oleic acid content. A spontaneous mutant line with high oleate levels, designated F435, has been identified through a peanut germplasm screening project, which contains 80% oleic acid (Norden et al., 1987). In this line, two mutations were reported: a “G” to “A” substitution at 448 bp after the start codon (G448A) in the ahFAD2A gene and an “A” insertion between bp 441 and 442 (441\_442insA) in the ahFAD2B gene. The study also identified a new mutation, G451T, in the ahFAD2B gene. These findings indicate that the mutations induced in ahFAD2B using CRISPR/Cas9 may be beneficial for developing high oleate peanut lines. Additionally, TALENs were employed to target the ahFAD2 genes to enhance oleic acid content (Wen et al., 2018). Two TALEN pairs were constructed; one pair inoculated 216 regenerated roots, while the other inoculated 105. The observed mutation frequencies were 8.33% and 12.38%, respectively, with most mutations consisting of small deletions ranging from 1 to 10 base pairs. In the resulting mutant lines, the oleic acid content of the seeds reached 80.45%, indicating a two-fold increase compared to wild-type plants, while linoleic acid content decreased to 3%, with no change in the total fatty acid content. Furthermore, Shu et al. (2020) edited nod factor receptors (NFRs) with CRISPR/Cas9 to investigate their roles in peanut nodulation. Mutants with edited AhNFR5 genes displayed a Nod- phenotype, while mutants with two selected AhNFR1 genes were still able to form nodules after inoculation.

Peanut (*Arachis hypogaea*) was a significant legume crop known for its high oleic acid content. A spontaneous mutant line with high oleate levels, designated F435, was identified through a peanut germplasm screening project, which contained 80% oleic acid (Norden et al., 1987). In this line, two mutations were reported: a “G” to “A” substitution at 448 bp after the start codon (G448A) in the ahFAD2A gene and an “A” insertion between bp 441 and 442 (441\_442insA) in the ahFAD2B gene. The study also identified a new mutation, G451T, in the ahFAD2B gene.

These findings indicated that the mutations induced in ahFAD2B using CRISPR/Cas9 could be beneficial for developing high oleate peanut lines.

Additionally, TALENs were employed to target the ahFAD2 genes to enhance oleic acid content (Wen et al., 2018). Two TALEN pairs were constructed; one pair inoculated 216 regenerated roots, while the other inoculated 105. The observed mutation frequencies were 8.33% and 12.38%, respectively, with most mutations consisting of small deletions ranging from 1 to 10 base pairs. In the resulting mutant lines, the oleic acid content of the seeds reached 80.45%, indicating a two-fold increase compared to wild-type plants, while linoleic acid content decreased to 3%, with no change in the total fatty acid content.

Furthermore, Shu et al. (2020) edited nod factor receptors (NFRs) with CRISPR/Cas9 to investigate their roles in peanut nodulation. Mutants with edited AhNFR5 genes displayed a Nod- phenotype, while mutants with two selected AhNFR1 genes were still able to form nodules after inoculation.

### 6. NON-EDITED LEGUME SPECIES

Lentil (*Lens culinaris*) is a diploid, self-pollinating plant belong from the Fabaceae family, known for its high protein, mineral, fiber, and carbohydrate content. In developing countries and among populations that cannot afford costly animal proteins, lentils can play a significant role in combating malnutrition and micronutrient deficiencies (Kumar et al., 2015). Additionally, lentils enhance soil quality by fixing nitrogen. The availability of draft genome sequences allows for the targeted identification of candidate genes associated with nutritional value, responses to abiotic and biotic stresses, and herbicide resistance. Genomic tools and technologies have the potential to advance lentil breeding efforts. Some gene transformation studies have been conducted on lentils. A notable example involves the successful transfer of the dehydration-responsive element binding gene (DREB1A), which helps plants respond to abiotic stresses, into lentils via *Agrobacterium*, producing transgenic plants that exhibit drought and salinity resistance (Khatib et al., 2011). In vitro regeneration post-transformation was crucial, with findings suggesting that using decapitated embryos was more effective for shoot generation than other tissues (Sarker et al., 2003). These

initiatives provided valuable resources for future genome-editing research. Several candidate genes related to abiotic and biotic stress factors, as well as agronomic traits, were identified in lentils (Kumar et al.). However, to date, there had been no genome-editing research on these candidate genes in lentils using ZFNs, TALENs, or CRISPR/Cas9 technologies. Genome-editing technologies could offer a cost-effective and straightforward method for elucidating the functions of these candidate genes, leading to the development of cultivars with desirable characteristics, such as stress tolerance and improved agronomic traits (Bhowmik et al., 2021).

Pea (*Pisum sativum*) was a significant legume crop worldwide, second only to the common bean (*Phaseolus vulgaris*), and was rich in dietary proteins, mineral nutrients, complex carbohydrates, and fiber (Bastianelli et al., 1998). Its ability to fix nitrogen symbiotically enhanced soil fertility, making it a valuable addition to agricultural systems (Mabrouk et al., 2018). Additionally, peas were historically important as a model organism used by Mendel to establish the principles of inheritance (Ellis et al., 2011). The pea genome was approximately 4.45Gb in size, with a reference genome published in 2019 that offered insights into the evolution of legume genomes (Kreplak et al., 2019). Genomic approaches were crucial for identifying genes associated with important traits and developing tools for crop improvement. Despite significant advances in pea cultivation, efforts to enhance crop yield and quality had to continue to meet the needs of the growing global population.

Peas were susceptible to various threats, including parasitic weeds, viruses, bacteria, fungi, and abiotic stressors such as drought, salinity, heat, and cold, all of which could lead to significant yield and growth losses. A successful stable transformation study was conducted, transferring the cry1Ac gene (which encodes a protoxin) from *Bacillus thuringiensis* (Negawo et al., 2013) and the alpha-amylase inhibitor gene from *Phaseolus vulgaris* to enhance insect tolerance using *Agrobacterium*-mediated transformation. Another attempt involved introducing antifungal genes against *Fusarium* spp. into peas; however, this resulted in unstable expression over three years of field trials. Recent advancements improved *Agrobacterium*-mediated transformation efficiency and regeneration frequency through extended

infection times and the addition of zeatin to the selection medium. The availability of genome sequence information, successful transformation methods, and regeneration techniques were essential for utilizing genome-editing tools in vegetable crops (Cardi et al., 2017). To date, there had been no studies employing new genome-editing technologies in peas, likely due to insufficient regeneration capabilities (Pandey et al., 2021). The advancement of new genome-editing methods could create opportunities in breeding for higher yields and the development of nutritionally enhanced plants.

Faba beans possessed several advantages over other legumes in cold climates, making them suitable for sustainable farming practices (Temesgen et al., 2015). Like other legumes, they contributed significantly to soil fertility. Faba bean breeding was essential to meet the growing demand for food and feed, as they were a rich source of protein, fiber, and other nutrients (Khazaei et al., 2021). However, publicly available genome sequence data for faba beans were lacking, likely due to the challenges associated with assembling their large genome (Cooper et al., 2017). Transformation and regeneration of faba beans remained difficult, although there had been attempts at *Agrobacterium*-mediated transformation with limited success. The first successful transformation was reported by Böttinger et al. (2001), who used in vitro development of internode stem segments invaded by *Agrobacterium*, and later by Hanafy et al. (2005), who infiltrated cut embryo axes with *Agrobacterium* to obtain stable transgenic lines. Additionally, abiotic stress-tolerant lines were generated by Hanafy et al. (2013) by transforming the potato PR10a gene into faba beans using the same method. However, various biotic and abiotic stress factors, including heat, insects, viruses, and parasitic weeds, continued to reduce faba bean yields.

The availability of whole-genome data for commonly cultivated non-edited legume species such as lentil, pea, common bean, and mung bean—which were rich in nutrients for human diets—presented opportunities for developing genome-editing strategies. Identifying and demonstrating the functions of genes associated with abiotic and biotic stress tolerance, yield, and quality could facilitate the application of new genome-editing tools to enhance these traits in key legume crops essential for human nutrition. Additionally, legumes played a crucial role in

maintaining soil fertility; thus, targeting symbiotic nitrogen-fixing pathways could be beneficial in genome-editing initiatives aimed at improving soil health. Despite some promising efforts, low transformation efficiency and challenges in regeneration remained significant obstacles, which may explain the limited application of new genome-editing techniques. To fully realize the potential for substantial improvements in these legumes, it was critical to develop innovative methods that enhanced transformation and regeneration efficiency, thereby supporting the nutritional needs of a large portion of the global population.

## 7. CONCLUSION AND FUTURE PERSPECTIVE

The assessment of genetic resources in legume crops has revealed significant variation in protein content and micronutrient levels. This review highlights ongoing efforts to screen germplasm, map quantitative trait loci (QTLs) associated with seed protein and micronutrient content, and identify future focus areas for enhancing the nutritional quality of key pulses. The concentration of micronutrients in seeds is a complex trait influenced by multiple genes and QTLs, with its levels significantly impacted by various soil and environmental factors.

Currently, an integrated breeding strategy that combines QTL mapping, genome-wide association studies, and differential gene expression profiling is considered the most effective approach for the genetic analysis of complex traits such as nutritional quality. There is considerable potential to improve the nutritional quality of pulse crops and develop biofortified varieties. The variability in iron (Fe) and zinc (Zn) content, along with other micronutrients in food legumes, can be utilized to enhance their levels and bioavailability.

Additionally, reducing anti-nutritional factors like phytates, trypsin inhibitors, and chymotrypsin inhibitors can further increase the bioavailability and consumption of legumes. A key objective in legume improvement programs is to identify genotypes with low anti-nutrient content, whether from existing germplasm or through methods like induced mutagenesis and genome editing, to achieve nutritional security. Furthermore, crop wild relatives of various legumes, which naturally possess lower levels of these anti-nutrients, can be integrated into pre-breeding programs (Singh and Jauhar, 2005).

## DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that generative AI technologies such as Large Language Models, etc have been used during writing or editing of this manuscript. This explanation will include the name, version, model, and source of the generative AI technology and as well as all input prompts provided to the generative AI technology.

Details of the AI usage are given below:

1. Chatgpt
2. copilot

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

- Abdollahi, M. R., & Rashidi, S. (2018). Production and conversion of haploid embryos in chickpea (*Cicer arietinum* L.) anther cultures using high 2,4-D and silver nitrate containing media. *Plant Cell, Tissue and Organ Culture (PCTOC)*, 133, 39–49.
- Ahmad, M. (2022). Genomics and transcriptomics to protect rice (*Oryza sativa* L.) from abiotic stressors: pathways to achieving zero hunger. *Frontiers in Plant Science*, 13, 1002596.
- Al Amin, N., Ahmad, N., Wu, N., Pu, X., Ma, T., Du, Y., et al. (2019). CRISPR-Cas9 Mediated Targeted Disruption of FAD2-2 Microsomal Omega-6 Desaturase in Soybean (*Glycine max*L.). *BMC Biotechnol.* 19, 9.
- Arora, L., and Narula, A. (2017). Gene Editing and Crop Improvement Using CRISPR-Cas9 System. *Front. Plant Sci.* 8, 1932. doi:10.3389/fpls.2017.01932
- Arora, L., and Narula, A. (2017). Gene Editing and Crop Improvement Using CRISPR-Cas9 System. *Front. Plant Sci.* 8, 1932.
- Atkins, P. A., and Voytas, D. F. (2020). Overcoming Bottlenecks in Plant Gene Editing. *Curr. Opin. Plant Biol.* 54, 79–84.
- Atkins, P. A., and Voytas, D. F. (2020). Overcoming Bottlenecks in Plant Gene Editing. *Curr. Opin. Plant Biol.* 54, 79–84.
- Badhan, S., Ball, A. S., and Mantri, N. (2021). First Report of CRISPR/Cas9 Mediated DNA-free Editing of *4CL* and *RVE7* Genes

- in Chickpea Protoplasts. *Int. J. Mol. Sci.* 22, 396.
- Badhan, S., Ball, A. S., and Mantri, N. (2021). First Report of CRISPR/Cas9 Mediated DNA-free Editing of *4CL* and *RVE7* Genes in Chickpea Protoplasts. *Int. J. Mol. Sci.* 22, 396.
- Bao, A., Zhang, C., Huang, Y., Chen, H., Zhou, X., and Cao, D. (2020). Genome Editing Technology and Application in Soybean Improvement. *Oil Crop Sci.* 5 (1), 31–40. ISSN 2096-2428.
- Bastianelli, D., Grosjean, F., Peyronnet, C., Duparque, M., and Régnier, J. M. (1998). Feeding Value of Pea (*Pisum Sativum*, L.) 1. Chemical Composition of Different Categories of Pea. *Anim. Sci.* 67 (3), 609–619.
- Bhowmik, P., Ellison, E., Polley, B., Bollina, V., Kulkarni, M., Ghanbarnia, K., Song, H., Gao, C., Voytas, D. F., & Kagale, S. (2018). Targeted mutagenesis in wheat microspores using CRISPR/Cas9. *Scientific Reports*, 8, 6502.
- Bhowmik, P., Konkin, D., Polowick, P., Hodgins, C. L., Subedi, M., Xiang, D. (2021). CRISPR/Cas9 Gene Editing in Legume Crops: Opportunities and Challenges. *Legume Sci.* 3(3).
- Bilichak, A., Sastry-Dent, L., Sriram, S., Simpson, M., Samuel, P., Webb, S., Jiang, F., & Eudes, F. (2020). Genome editing in wheat microspores and haploid embryos mediated by delivery of ZFN proteins and cell-penetrating peptide complexes. *Plant Biotechnology Journal*, 18, 1307–1316.
- Bobkov, S. (2014). Obtaining callus and regenerated plants in anther cultures of pea. *Czech Journal of Genetics and Plant Breeding*, 50: 123–129.
- Bottero, E., Massa, G., González, M., Stritzler, M., Tajima, H., Gómez, C., et al. (2021). Efficient CRISPR/Cas9 Genome Editing in Alfalfa Using a Public Germplasm. *Front. Agron.* 3: 661526.
- Böttinger, P., Steinmetz, A., Schieder, O., and Pickardt, T. (2001). Agrobacterium Mediated Transformation of *Vicia faba*. *Mol. Breed.* 8: 243–254.
- Brinkman, E. K., & van Steensel, B. (2019). Rapid quantitative evaluation of CRISPR genome editing by TIDE and TIDER. *Methods in Molecular Biology*, 1961, 29–44.
- Brinkman, E. K., Chen, T., Amendola, M., & van Steensel, B. (2014). Easy quantitative assessment of genome editing by sequence trace decomposition. *Nucleic Acids Research*. 42: 168–e168.
- Cai K, K., Yin, J., Chao, H., Ren, Y., Jin, L., Cao, Y., et al. (2018). A C3HC4-type RING Finger Protein Regulates Rhizobial Infection and Nodule Organogenesis in *Lotus Japonicus*. *J. Integr. Plant Biol.* 60(9): 878–896.
- Cardi, T., D'Agostino, N., and Tripodi, P. (2017). Genetic Transformation and Genomic Resources for Next-Generation Precise Genome Engineering in Vegetable Crops. *Front. Plant Sci.* 8, 241.
- Che, P., Chang, S., Simon, M. K., Zhang, Z., Shaharyar, A., Ourada, J., et al. (2021). Developing a Rapid and Highly Efficient Cowpea Regeneration, Transformation and Genome Editing System Using Embryonic axis Explants. *Plant J.* 106 (3), 817–830.
- Che, P., Chang, S., Simon, M. K., Zhang, Z., Shaharyar, A., Ourada, J., et al. (2021). Developing a Rapid and Highly Efficient Cowpea Regeneration, Transformation and Genome Editing System Using Embryonic axis Explants. *Plant J.* 106 (3), 817–830.
- Chopra, R., & Saini, R. (2012). Use of sonication and vacuum infiltration for *Agrobacterium*-mediated transformation of an Indian lentil (*Lens culinaris* Medik.) cultivar. *Scientia Horticulturae*, 143, 127–134.
- Confalonieri, M., Carelli, M., Gianoglio, S., Moglia, A., Biazzi, E., and Tava, A. (2021). CRISPR/Cas9-Mediated Targeted Mutagenesis of *CYP93E2* Modulates the Triterpene Saponin Biosynthesis in *Medicago Truncatula*. *Front. Plant Sci.* 12, 690231.
- Cooper, J. W., Wilson, M. H., Derks, M. F. L., Smit, S., Kunert, K. J., Cullis, C., et al. (2017). Enhancing Faba Bean (*Vicia faba* L.) Genome Resources. *J. Exp. Bot.* 68 (8), 1941–1953.
- Curtin, S. J., Xiong, Y., Michno, J.-M., Campbell, B. W., Stec, A. O., Čermák, T., et al. (2018). CRISPR/Cas9 and TALENs Generate Heritable Mutations for Genes Involved in Small RNA Processing of Glycine max and *Medicago Truncatula*. *Plant Biotechnol. J.* 16, 1125–1137.
- Curtin, S. J., Zhang, F., Sander, J. D., Haun, W. J., Starker, C., Baltes, N. J., et al. (2011). Targeted Mutagenesis of Duplicated Genes in Soybean with Zinc-Finger

- Nucleases. *Plant physiol.* 156 (2), 466–473.
- Dan, Y., Armstrong, C. L., Dong, J., Feng, X., Fry, J. E., Keithly, G. E., Martinell, B. J., Roberts, G. A., Smith, L. A., & Tan, L. J. (2009). Lipoic acid—An unique plant transformation enhancer. *In Vitro Cellular & Developmental Biology-Plant*, 45, 630.
- Demorest, Z. L., Coffman, A., Baltes, N. J., Stoddard, T. J., Clasen, B. M., Luo, S., et al. (2016). Direct Stacking of Sequence-specific Nuclease-Induced Mutations to Produce High Oleic and Low Linolenic Soybean Oil. *BMC Plant Biol.* 16, 1–8.
- Di, Y.-H., Sun, X.-J., Hu, Z., Jiang, Q.-Y., Song, G.-H., Zhang, B., et al. (2019). Enhancing the CRISPR/Cas9 System Based on Multiple GmU6 Promoters in Soybean. *Biochem. Biophysical Res. Commun.* 519 (4), 819–823.
- Do, P. T., Nguyen, C. X., Bui, H. T., Tran, L. T. N., Stacey, G., Gillman, J. D., et al. (2019). Demonstration of Highly Efficient Dual gRNA CRISPR/Cas9 Editing of the Homeologous GmFAD2-1A and GmFAD2-1B Genes to Yield a High Oleic, Low Linoleic and  $\alpha$ -linolenic Acid Phenotype in Soybean. *BMC Plant Biol.* 19, 311.
- Ellis, T. H. N., Hofer, J. M. I., Timmerman-Vaughan, G. M., Coyne, C. J., and Hellens, R. P. (2011). Mendel, 150 Years on. *Trends Plant Sci.* 16, 590–596.
- Gnanasambandam, A., Paull, J., Torres, A., Kaur, S., Leonforte, T., Li, H., Materne, M., 2012. Impact of molecular technologies on faba bean (*Vicia faba* L.) breeding strategies. *Agronomy* 2 (3), 132–166.
- Han, J., Guo, B., Guo, Y., Zhang, B., Wang, X., and Qiu, L.-J. (2019). Creation of Early Flowering Germplasm of Soybean by CRISPR/Cas9 Technology. *Front. Plant Sci.* 10, 1446.
- Hanafy, M. S., El-Banna, A., Schumacher, H. M., Jacobsen, H.-J., and Hassan, F. S. (2013). Enhanced Tolerance to Drought and Salt Stresses in Transgenic Faba Bean (*Vicia faba* L.) Plants by Heterologous Expression of the PR10a Gene from Potato. *Plant Cell Rep.* 32, 663–674.
- Hanafy, M., Pickardt, T., Kiesecker, H., and Jacobsen, H.-J. (2005). Agrobacterium-mediated Transformation of Faba Bean (*Vicia faba* L.) Using Embryo Axes. *Euphytica* 142, 227–236.
- Haun, W., Coffman, A., Clasen, B. M., Demorest, Z. L., Lowy, A., Ray, E., et al. (2014). Improved Soybean Oil Quality by Targeted Mutagenesis of the Fatty Acid Desaturase 2 Gene Family. *Plant Biotechnol. J.* 12, 934–940.
- Hendling, M., Pabinger, S., Peters, K., Wolff, N., Conzemius, R., & Barišić, I. (2018). Oli2go: An automated multiplex oligonucleotide design tool. *Nucleic Acids Research*, 46, W252–W256.
- Hsiau, T., Maures, T., Waite, K., Yang, J., Kelso, R., Holden, K., & Stoner, R. (2018). Inference of CRISPR edits from Sanger trace data. *bioRxiv*, 251082.
- Huang, T.-K., and Puchta, H. (2019). CRISPR/Cas-mediated Gene Targeting in Plants: Finally a Turn for the Better for Homologous Recombination. *Plant Cell Rep.* 38, 443–453.
- Ji, J., Zhang, C., Sun, Z., Wang, L., Duanmu, D., and Fan, Q. (2019). Genome Editing in Cowpea *Vigna Unguiculata* Using CRISPR-Cas9. *Int. J. Mol. Sci.* 20 (10), 2471.
- Juranić, M., Nagahatenna, D. S. K., Salinas-Gamboa, R., Hand, M. L., Sánchez-León, N., Leong, W. H., et al. (2020). A Detached Leaf Assay for Testing Transient Gene Expression and Gene Editing in Cowpea (*Vigna Unguiculata* [L.] Walp.). *Plant Methods* 16, 88.
- Karst, S. M., Ziels, R. M., Kirkegaard, R. H., Sørensen, E. A., McDonald, D., Zhu, Q., Knight, R., & Albertsen, M. (2021). High-accuracy long-read amplicon sequences using unique molecular identifiers with Nanopore or PacBio sequencing. *Nature Methods*, 18, 165–169.
- Khatib, F., Makris, A., Yamaguchi-Shinozaki, K., Kumar, S., Sarker, A., Erskine, W., et al. (2011). Expression of the *DREB1A* Gene in Lentil (*Lens Culinaris* Medik. Subsp. *Culinaris*) Transformed with the Agrobacterium System. *Crop Pasture Sci.* 62 (6), 488–495.
- Khazaei, H., O'Sullivan, D. M., Stoddard, F. L., Adhikari, K. N., Paull, J. G., Schulman, A. H., et al. (2021). Recent Advances in Faba Bean Genetic and Genomic Tools for Crop Improvement. *Legume Sci.* 3 (3), 75.
- Kim, J., Kim, D., Lim, J. S., Maeng, J. H., Son, H., Kang, H.-C., Nam, H., Lee, J. H., & Kim, S. (2019). The use of technical replication for detection of low-level somatic mutations in next-generation sequencing. *Nature Communications*, 10, 1047.



- Kreplak, J., Madoui, M.-A., Cápál, P., Novák, P., Labadie, K., Aubert, G., et al. (2019). A Reference Genome for Pea Provides Insight into Legume Genome Evolution. *Nat. Genet.* 51 (9), 1411–1422.
- Kumar, A., Singh, A., Arya, M., Bhanu, A. N., Jaiswal, A., Kumar, S., & Chaturvedi, S. K. (2023). Screening of chickpea (*Cicer arietinum* L.) germplasm accessions for tolerance against post-emergence herbicide imazethapyr. *Journal of Food Legumes*, 36(2&3), 136-142. 10.59797/jfl.v36.i2.143
- Kumar, Amit. (2020). Evaluation and screening of chickpea (*Cicer arietinum* L.) germplasm accessions for tolerance against post-emergence herbicide. Master's thesis. Rani Lakshmi Bai Central Agricultural University Jhansi – 284 003 (Uttar Pradesh), <https://krishikosh.egranth.ac.in/handle/1/5810179129>
- Kumar, S., Rajendran, K., Kumar, J., Hamwiah, A., and Baum, M. (2015). Current Knowledge in Lentil Genomics and its Application for Crop Improvement. *Front. Plant Sci.* 6, 78.
- Lai, E. M., Shih, H. W., Wen, S. R., Cheng, M. W., Hwang, H. H., & Chiu, S. H. (2006). Proteomic analysis of *Agrobacterium tumefaciens* response to the *vir* gene inducer acetosyringone. *Proteomics*, 6, 4130–4136.
- Lewis, G. P. (2005). Legumes of the world. Richmond, UK: Royal Botanic Gardens, Kew.
- Lewis, G. P., Schrire, B., Mackinder, B., and Lock, M. (2005). Legumes of the World.
- Li, C., Zong, Y., Wang, Y., Jin, S., Zhang, D., Song, Q., et al. (2018). Expanded Base Editing in Rice and Wheat Using a Cas9-Adenosine Deaminase Fusion. *Genome Biol.* 19, 59.
- Liu, Z., Park, B. -J., Kanno, A., & Kameya, T. (2005). The novel use of a combination of sonication and vacuum infiltration in *Agrobacterium*-mediated transformation of kidney bean (*Phaseolus vulgaris* L.) with lea gene. *Molecular Breeding*, 16, 189.
- Lulsdorf, M., Croser, J., & Ochatt, S. (2011). 11 Androgenesis and doubled- haploid production in food legumes. *Biology and Breeding of Food Legumes*, 159, 159–177.
- Mahmoudian, M., Yücel, M. & Öktem, H. A. (2002). Transformation of lentil (*Lens culinaris* M.) cotyledonary nodes by vacuum infiltration of *Agrobacterium tumefaciens*. *Plant Molecular Biology Reporter*, 20: 251–257.
- Meng, Y., Hou, Y., Wang, H., Ji, R., Liu, B., Wen, J., et al. (2017). Targeted Mutagenesis by CRISPR/Cas9 System in the Model Legume *Medicago Truncatula*. *Plant Cell Rep.* 36, 371–374.
- Menz, J., Modrzejewski, D., Hartung, F., Wilhelm, R., Sprink, T. (2020). Genome edited crops touch the market: a view on the global development and regulatory environment. *Front. Plant Sci.* 11, 586027.
- Michno, J., Wang, X., Liu, J., Curtin, S., Kono, T., & Stupar, R. (2015). CRISPR/Cas mutagenesis of soybean and *Medicago truncatula* using a new web-tool and a modified Cas9 enzyme. *GM Crops Food*, 6, 243–252.
- Mikic, A., Peric´, V., Đorđević´, V., Srebric´, M., Mihailovic´, V., 2009. Anti-nutritional factors in some grain legumes. *Biotechnol. Anim. Husbandry* 25 (5-6-2), 1181–1188.
- Miyaoka, Y., Mayerl, S. J., Chan, A. H., & Conklin, B. R. (2018). Detection and Quantification of HDR and NHEJ Induced by Genome Editing at Endogenous Gene Loci Using Droplet Digital PCR. *Methods in Molecular Biology*, 1768, 349–362.
- Negawo, A. T., Aftabi, M., Jacobsen, H.-J., Altosaar, I., and Hassan, F. S. (2013). Insect Resistant Transgenic Pea Expressing *cry1Ac* Gene Product from *Bacillus Thuringiensis*. *Biol. Control* 67, 293–300.
- Norden, A. J., Gorbet, D. W., Knauff, D. A., and Young, C. T. (1987). Variability in Oil Quality Among Peanut Genotypes in the Florida Breeding Program1. *Peanut Sci.* 14, 7–11.
- Ochatt, S., Conreux, C., Moussa Mcolo, R., Despierre, G., Magnin- Robert, J.-B., & Raffiot, B. (2018). Phytosulfokine-alpha, an enhancer of in vitro regeneration competence in recalcitrant legumes. *Plant Cell, Tissue and Organ Culture (PCTOC)*, 135, 189–201.
- Ochatt, S., Pech, C., Grewal, R., Conreux, C., Lulsdorf, M., & Jacas, L. (2009). Abiotic stress enhances androgenesis from isolated micro- spores of some legume species (Fabaceae). *Journal of Plant Physiology*, 166, 1314–1328.
- Olhoft, P., Lin, K., Galbraith, J., Nielsen, N., & Somers, D. (2001). The role of thiol compounds in increasing *Agrobacterium*-mediated transformation of soybean

- cotyledonary-node cells. *Plant Cell Reports*, 20, 731–737.
- Pandey, A. K., Rubiales, D., Wang, Y., Fang, P., Sun, T., Liu, N., et al. (2021). Omics Resources and Omics-Enabled Approaches for Achieving High Productivity and Improved Quality in Pea (*Pisum Sativum* L.). *Theor. Appl. Genet.* 134, 755–776.
- Pandey, M.K., Roorkiwal, M., Singh, V.K., Ramalingam, A., Kudapa, H., Thudi, M., Varshney, R.K., 2016. Emerging genomic tools for legume breeding: current status and future prospects. *Front. Plant Sci.* 7, 455.
- Peng, C., Wang, H., Xu, X., Wang, X., Chen, X., Wei, W., Lai, Y., Liu, G., Godwin, I. D., & Li, J. (2018). High-throughput detection and screening of plants modified by gene editing using quantitative real-time polymerase chain reaction. *The Plant Journal*, 95, 557–567.
- Polowick, P., Quandt, J. and Mahon, J. (2000). The ability of pea transformation technology to transfer genes into peas adapted to western Canadian growing conditions. *Plant Science*, 153, 161–170.
- Ribalta, F., Croser, J. S., & Ochatt, S. (2012). Flow cytometry enables identification of sporophytic eliciting stress treatments in gametic cells. *Journal of Plant Physiology*, 169, 104–110.
- Rodas, A. L., Roque, E., Hamza, R., Gómez-Mena, C., Minguet, E. G., Wen, J., et al. (2021). MtSUPERMAN Plays a Key Role in Compound Inflorescence and Flower Development in *Medicago Truncatula*. *Plant J.* 105, 816–830.
- Sarker, R. H., Biswas, A., Mustafa, B. M., Mahbub, S., and Hoque, M. I. (2003). Agrobacterium-mediated Transformation of Lentil (*Lens Culinaris* Medik.). *Plant Tissue Cult.* 13 (1), 1–12.
- Sato, S., Nakamura, Y., Kaneko, T., Asamizu, E., Kato, T., Nakao, M., et al. (2008). Genome Structure of the Legume, *Lotus Japonicus*. *DNA Res.* 15 (4), 227–239.
- Schmutz, J., Cannon, S. B., Schlueter, J., Ma, J., Mitros, T., Nelson, W., et al. (2010). Genome Sequence of the Palaeopolyploid Soybean. *Nature* 463, 178–183
- Schroeder, H. E., Gollasch, S., Moore, A., Tabe, L. M., Craig, S., Hardie, D. C., et al. (1993). Bean [alpha]-Amylase Inhibitor Confers Resistance to the Pea Weevil (*Bruchus Pisorum*) in Transgenic Peas (*Pisum Sativum* L.). *Plant Physiol.* 107, 1233–1239.
- Shu, H., Luo, Z., Peng, Z., and Wang, J. (2020). The Application of CRISPR/ Cas9 in Hairy Roots to Explore the Functions of *AhNFR1* and *AhNFR5* Genes during Peanut Nodulation. *BMC Plant Biol.* 20, 417.
- Simmonds, D., & Donaldson, P. (2000). Genotype screening for proliferative embryogenesis and biolistic transformation of short-season soybean genotypes. *Plant Cell Reports*, 19, 485–490.
- Tang, X., Liu, G., Zhou, J., Ren, Q., You, Q., Tian, L., Xin, X., Zhong, Z., Liu, B., Zheng, X., Zhang, D., Malzahn, A., Gong, Z., Qi, Y., Zhang, T., & Zhang, Y. (2018). A large-scale whole-genome sequencing analysis reveals highly specific genome editing by both Cas9 and Cpf1 (Cas12a) nucleases in rice. *Genome Biology*, 19, 84.
- Temesgen, T., Kenei, G., Sefera, T., and Jarso, M. (2015). Yield Stability and Relationships Among Stability Parameters in Faba Bean (*Vicia faba* L.) Genotypes. *Crop J.* 3 (3), 258–268.
- Thomas, H. R., Percival, S. M., Yoder, B. K., & Parant, J. M. (2014). High-throughput genome editing and phenotyping facilitated by high resolution melting curve analysis. *PLoS One*, 9, e114632.
- Trick, H. N., & Finer, J. J. (1997). SAAT: Sonication-assisted Agrobacterium-mediated transformation. *Transgenic Research*, 6, 329–336.
- Varshney, R. K., Thudi, M., Pandey, M. K., Tardieu, F., Ojiewo, C., Vadez, V., et al. (2018). Accelerating Genetic Gains in Legumes for the Development of Prosperous Smallholder Agriculture: Integrating Genomics, Phenotyping, Systems Modelling and Agronomy. *J. Exp. Bot.* 69 (13), 3293–3312.
- Wang J, J., Kuang, H., Zhang, Z., Yang, Y., Yan, L., Zhang, M., et al. (2020). Generation of Seed Lipxygenase-free Soybean Using CRISPR-Cas9. *Crop J.* 8 (3), 432–439. ISSN 2214-5141.
- Wang, L., Rubio, M. C., Xin, X., Zhang, B., Fan, Q., Wang, Q., et al. (2019). CRISPR/Cas9 Knockout of Leghemoglobin Genes in Lotus Japonicus uncovers Their Synergistic Roles in Symbiotic Nitrogen Fixation. *New Phytol.* 224, 818–832.
- Wang, L., Wang, L., Tan, Q., Fan, Q., Zhu, H., Hong, Z., et al. (2016). Efficient Inactivation of Symbiotic Nitrogen Fixation Related Genes in *Lotus Japonicus*

- Using CRISPR-Cas9. *Front. Plant Sci.* 7, 1333.
- Wen, S., Liu, H., Li, X., Chen, X., Hong, Y., Li, H., et al. (2018). TALEN-mediated Targeted Mutagenesis of *Fatty Acid Desaturase 2 (FAD2)* in Peanut (*Arachis hypogaea* L.) Promotes the Accumulation of Oleic Acid. *Plant Mol. Biol.* 97, 177–185.
- Wolabu, T. W., Park, J. J., Chen, M., Cong, L., Ge, Y., Jiang, Q., et al. (2020). Improving the Genome Editing Efficiency of CRISPR/Cas9 in Arabidopsis and *Medicago Truncatula*. *Planta* 252, 15.
- Wu, N., Lu, Q., Wang, P., Zhang, Q., Zhang, J., Qu, J., et al. (2020). Construction and Analysis of *GmFAD2-1A* and *GmFAD2-2A* Soybean Fatty Acid Desaturase Mutants Based on CRISPR/Cas9 Technology. *Int. J. Mol. Sci.* 21 (3), 1104.
- Xie, K., Minkenberg, B., and Yang, Y. (2013). Boosting CRISPR/Cas9 Multiplex Editing Capability with the Endogenous tRNA-Processing System. *Proc. Natl. Acad. Sci. U.S.A.* 112, 3570–3575.
- Yadav, R., Mehrotra, M., Singh, A. K., Niranjana, A., Singh, R., Sanyal, I., Lehri, A., Pande, V., & Amla, D. (2017). Improvement in *Agrobacterium*-mediated transformation of chickpea (*Cicer arietinum* L.) by the inhibition of polyphenolics released during wounding of cotyledonary node explants. *Protoplasma*, 254, 253–269.
- Yadav, S. K., Katikala, S., Yellisetty, V., Kannepalle, A., Narayana, J. L., Maddi, V., et al. (2012). Optimization of *Agrobacterium* Mediated Genetic Transformation of Cotyledonary Node Explants of *Vigna Radiata*. *SpringerPlus* 1 (1), 1–8.
- Young, N. D., Debelle, F., Oldroyd, G. E., Geurts, R., Cannon, S. B., Udvardi, M. K., et al. (2011). The Medicago Genome Provides Insight into the Evolution of Rhizobial Symbioses. *Nature* 480, 520–524.
- Yuan, M., Zhu, J., Gong, L., He, L., Lee, C., Han, S., et al. (2019). Mutagenesis of *FAD2* Genes in Peanut with CRISPR/Cas9 Based Gene Editing. *BMC Biotechnol.* 19, 24.
- Zhang, H., Cao, Y., Zhang, H., Xu, Y., Zhou, C., Liu, W., et al. (2020). Efficient Generation of CRISPR/Cas9-Mediated Homozygous/Biallelic *Medicago Truncatula* Mutants Using a Hairy Root System. *Front. Plant Sci.* 11, 294.
- Zheng, N., Li, T., Dittman, J. D., Su, J., Li, R., Gassmann, W., et al. (2020). CRISPR/Cas9-Based Gene Editing Using Egg Cell-specific Promoters in Arabidopsis and Soybean. *Front. Plant Sci.* 11: 800.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/126488>