



## **Ascertainment of *In vivo* Antidiarrheal and *In vitro* Thrombolytic Effect of Ethanolic Extract of Leaves of *Amomum dealbatum***

**Md. Azimul Islam<sup>1</sup>, Mohammed Aktar Sayeed<sup>1</sup>, Md. Abdul Barek<sup>2\*</sup>,  
Enama Nabi Shetu<sup>1</sup> and Md. Nurul Faisal<sup>1</sup>**

<sup>1</sup>Department of Pharmacy, International Islamic University Chittagong, Kumira, Chittagong-4318, Bangladesh.

<sup>2</sup>Department of Pharmacy, Noakhali Science and Technology University, Sonapur, Noakhali-814, Bangladesh.

### **Authors' contributions**

*This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.*

### **Article Information**

DOI: 10.9734/JALSI/2019/v21i1130093

#### Editor(s):

(1) Dr. Ali Mohamed Elshafei Ali, Professor, Department of Microbial Chemistry, Genetic Engineering & Biotechnology Building, National Research Centre, Egypt.

#### Reviewers:

(1) Ochieng O. Anthony, Sumait University, Tanzania.

(2) Md. Shariful Islam, Southeast University, Bangladesh.

Complete Peer review History: <http://www.sdiarticle3.com/review-history/49046>

**Original Research Article**

**Received 11 March 2019**

**Accepted 16 May 2019**

**Published 25 May 2019**

### **ABSTRACT**

**Aims:** The present study aimed to investigate antidiarrheal and thrombolytic effect of ethanolic extract of leaves of *A. dealbatum* in mice.

**Study design:** Antidiarrheal effect was evaluated by castor oil-induced diarrhea method at two different concentrations in mice and *in vitro* thrombolytic activity was analyzed with clot lysis assay of human blood.

**Place and duration of study:** Department of Pharmacy, International Islamic University Chittagong, Kumira, Chittagong-4318, Bangladesh, between December 2018 and February 2019.

**Methodology:** The male Swiss mice's were divided into four groups (n = 5). First group was orally treated with 1% Tween-80 (10 ml/kg) and second group was orally treated with loperamide (5 mg/kg). Third and fourth group were orally treated with ethanolic extract of leaves of *A. dealbatum* at 200 and 400 mg/kg accordingly. Human RBCs were collected for conducting thrombolytic assay.

\*Corresponding author: E-mail: barekpharma@gmail.com;

During this study, 1.5 ml of venous blood was drawn from healthy volunteers (n = 10) and Streptokinase was employed as positive control and distilled water was employed as negative control.

**Results:** In castor oil induced diarrhea model, ethanolic extract of leaves of *A. dealbatum* at 200, 400 mg/kg and loperamide (5 mg/kg) significantly reduced the number of feces and increase percent of inhibition of defecations compared to negative control. The extract showed percent of inhibition of defecation of 16.67 and 37.50 for 200 and 400 mg/ml respectively where the positive control loperamide showed 66.67%. Percentage of clot disruptions were 4.51 ( $p < .001$ ), 75.69 ( $p < .001$ ) and 26.07 ( $p < .001$ ) for water, streptokinase and 10 mg/ml extract respectively.

**Conclusion:** Based on the results from in vivo and in vitro activities, the leaves of *A. dealbatum* were found to be a potential source of new antidiarrheal and thrombolytic agents.

**Keywords:** *Amomum dealbatum*; anti-diarrheal; castor oil; thrombolytic; clot disruptions.

## 1. INTRODUCTION

Plants are known to be the source of many chemical compounds was used by people of ancient cultures without knowledge of their active ingredients. World Health Organization (WHO) has provided a definition of medicinal plants, that is "A medicinal plant is any plant which, in one or more of its organs, contains substances that can be used for therapeutic purposes or which are precursors for synthesis of useful drug" [1]. In the Plant Kingdom, medicinal plants form the largest single grouping of plants. It is estimated that 30,000 species worldwide fall in this group, of which around 33% are trees [2]. In last few years, there has been great focus on the possible health benefits of natural substances with antidiarrheal, thrombolytic, antioxidant, antimicrobial, analgesic, antipyretic, sedative, antidepressant, antipsychotic, anticancer, anti-diabetic and others activities [3]. Therefore, it is necessary to establish scientific evidences for therapeutic use of such traditional medicinal plants. Zingiberaceae, the ginger family of flowering plants, the largest family of the order Zingiberales, containing 52 genera with a total of about 1600 known species [4]. The family is chiefly distributed throughout tropical and subtropical regions of Africa, Asia, China, Nepal, India, Thailand, Indonesia, Malaysia, Singapore, Brunei, Philippines, Papua New Guinea and the Americas [5]. *Amomum dealbatum* known locally as "Alachengay" which belongs to a family called Zingiberaceae. This plant is a robust perennial herb, growing up to 3 meters tall with a thick rhizome. Leaves oblong-lanceolate, pubescent beneath. Spikes oblong, peduncle as long as the spike. Corolla-tube cylindrical; segments obtuse, half as long as the tube; lip deflexed, ligulate, red-yellow. Fruit ovoid, strongly ribbed [6]. *A. dealbatum* is widely found in Bangladesh, Assam, China South-Central, East Himalaya,

Laos, Indonesia, Myanmar, Nepal, Thailand, Vietnam [6]. In Bangladesh they distributed in forests and shady places of Chittagong, Chittagong Hill Tracts and Sylhet [7]. Diarrhea is characterized by the passage of abnormally liquid or watery fecal matter associated with increased frequency of defecation (three or more times in a day) and abdominal pain [8,9]. It is the world's third highest killer disease and about 70% people are affected by diarrhea [10,11]. The conditions of diarrhea are particularly dangerous in infants and young children because of the rapidity with which serious dehydration occur [12]. This disease account for one in nine child deaths worldwide and around 760,000 children death every year [13]. So, many works have been carried out in order to discover new antidiarrheal compounds from natural sources for their diverse pharmacological and biological properties [14]. Thrombosis is a lethal disease which is characterized by the formation of blood clots (thrombus) in the circulatory system because of the imbalance of homeostatic system of physiological procedures [15]. This is connected with acute coronary disorders such as pulmonary emboli, deep vein thrombosis, strokes, heart attacks, and venous thromboembolic disorders that account for sudden morbidity and mortality [16]. Thrombosis leads to vascular blockade and while recovering it causes fatal consequences, such as cerebral or myocardial infarction and even death. Thrombolytic agents including tissue plasminogen activator (t-PA), alteplase, anistreplase, urokinase (UK), and streptokinase and recombinant t-PA therapies have been used as effective treatment for thrombolysis. UK and SK are widely used in India, Bangladesh and other developing countries due to lower cost [17] as compared to other thrombolytic drugs but the use is associated with high risk of anaphylactic reaction, systemic fibrinolysis, hemorrhage, slow

reperfusion rate and frequent early reclosures and lacks specificity [18]. Moreover, these drugs are not used in patients who have undergone surgery or those with a history of nervous lesions, gastrointestinal bleeding or hypertension [19]. For that reason, alternative options as traditional and herbal drugs are highly necessitated and numbers of plants have already been reported to show very emerging and potential thrombolytic agents. This study deals with the pharmacological actions namely antidiarrheal and thrombolytic effects of a newer source of indigenous medicinal plant *Amomum dealbatum*.

## 2. MATERIALS AND METHODS

### 2.1 Drugs and Chemicals

All chemicals and reagents used in this study were of analytical grade. Ethanol (Merck, Germany) was used as a solvent during extraction. Standard streptokinase was purchased from Popular Pharmaceuticals Limited, Bangladesh. Loperamide (Square Pharmaceuticals Limited), castor oil (WELL's Health Care, Spain) and Tween 80 (HiMedia Laboratories Pvt. Limited, Mumbai, India) were also used in this research.

### 2.2 Plant Materials

*Amomum dealbatum* was collected from kaptai shitapahar, Chittagong, Bangladesh on end of December 2017 and was identified by National Herbarium Institute, Mirpur, Dhaka, Bangladesh (Accession number: DACB-43725).

### 2.3 Extraction

After collection of whole plants of *A. dealbatum* was thoroughly washed with water. Then the selected plant part (leaves) was dried and powdered. About 520 g of the powdered materials of plant was taken separately in a clean, flat bottomed glass container and soaked in 2500 ml of ethanol at room temperature for two weeks accompanying occasional shaking and stirring. Then the solution was filtered using filter cloth and Whatman filter paper (Bibby RE200, Sterilin Ltd., UK) and concentrated with a rotary evaporator (RE-EV311-V, LabTeck S.R.L, Italy). It rendered a gummy concentrate of deep green color. The gummy concentrate was designated as crude ethanolic extract.

### 2.4 Experimental Animals

All animal procedures and experimental protocols were approved by the Research Ethics Committee of the institution and were carried out in accordance with the Guide for the Care and use of Laboratory Animals [20]. Swiss albino mice, weighing about 25–30 gram, were collected from Jahangirnagar University, Savar, Bangladesh. The animals were provided with standard laboratory food and distilled water ad libitum and maintained at natural day-night cycle having proper ventilation in the room. All the experiments were conducted in an isolated and noiseless condition. The study protocol was approved by the P&D Committee, Department of Pharmacy, International Islamic University Chittagong, Bangladesh (Pharm-P&D-37/07'12). The animals were acclimatized to laboratory condition for 10 days prior to experimentation.

### 2.5 Effect on Castor oil Induced Diarrhea

Castor oil induced diarrhea method described by Franca 2008 [21] was followed for this study. Four groups of five mice were selected for the final experiment. Group I received 1% Tween-80 (10 ml/kg), second group received loperamide (5 mg/kg) and other groups received ethanol extract 200 and 400 mg/kg accordingly. Castor oil (0.5 ml/animal) was administered after 60 minutes. Immediately after administering castor oil, each animal was kept in an individual cage with a floor lined with blotting paper. The characteristic diarrheal droppings (wet & dry feces) were noted and observed for 4 hours study for each mouse. 100% was considered as the total number of feces of control group [22]. At the beginning of each hour old papers were replaced with the new ones. Percentage of inhibition of defecation was calculated relative to the control using the following relationship-

$$\text{Inhibition of defecation (\%)} = \frac{A-B}{A} \times 100$$

Where, A is mean number of defecation feces of the control group and B is mean number of defecation caused by standard or plant extracts.

### 2.6 Thrombolytic Activity

The thrombolytic activity of plant extracts was evaluated by the method developed by Prasad et al. [23] with modification to use streptokinase as standard [18,24].

**Red blood cells (RBC) collection:** Human RBCs were collected for conducting thrombolytic assay. Male volunteers- weighing average 65 and free from diseases were selected to collect RBCs (using a protocol approved by Institutional Ethics Committee).

**Specimen:** 100 mg *A. dealbatum* ethanolic extract was suspended in 10 ml distilled water and the suspension was shaken vigorously on a vortex mixer. The suspension was kept overnight and decanted to remove the soluble supernatant, which was filtered through a 0.22-micron syringe filter. 100 µl of the plant extract was added to the eppendorf tube which contained the clots to check thrombolytic activity [23,24]. Streptokinase was employed as positive control and distilled water was employed as negative control.

**Thrombolytic assay:** During this study, 1.5 ml of venous blood was drawn from healthy volunteers (n = 10) and transferred to three different pre-weighed sterilized Eppendorf Tubes (0.5 ml/tube). The Eppendorf Tubes were incubated at 37 °C for 45 minutes. After formation of a clot, serum was completely discarded from the tubes (carried out without disturbing the clot formed). Each Eppendorf Tube was weighed to determine weight of the clot. Each Eppendorf Tube was appropriately labeled and 100 µl of the plant extract (10 mg/ml) was added to the tubes. 100 µl of streptokinase and 100 µl of water were distinctly added to the control tubes numbered. The tubes were incubated again at 37 °C for 90 minutes and observed for clot lysis. After the following incubation, the obtained fluid was discarded from the tubes. They were again weighed to observe the weight of released clot [23, 24]. Every test samples were examined in triplicate. Finally, the result was expressed as percentage of clot lysis which is calculated by the following equation:

$$\% \text{ of clot lysis} = \frac{\text{Weight of released clot}}{\text{Clot weight}} \times 100$$

## 2.7 Statistical Analysis

The data from antidiarrheal and thrombolytic assay were expressed as Mean ± Standard Error Mean (SEM) and analyzed by one-way analysis of variance (ANOVA) followed by Dunnett 't' test using SPSS software of 20 version.  $p < 0.05$  was considered statistically significant.

## 3. RESULTS

### 3.1 Effect on Castor Oil- induced Diarrhea

We evaluated the effect of ethanolic extract of *A. dealbatum* leaves on castor oil induced diarrhea. The trend in number of feces was also observed for control (14.40,  $p < .01$ ), standard (4.80,  $p < .01$ ), extracts 200 mg/kg (12.00,  $p = .05$ ) and 400 mg/kg (9.00,  $p < .01$ ) of plant sample (Table 1). When calculating percentage of inhibition of defecation, it was observed that the inhibition of defecation (%) in the dose of 200 mg/kg and 400 mg/kg are 16.67% and 37.50% respectively while standard loperamide (5 mg/kg) showed 66.67% (Fig. 1).


### 3.2 Thrombolytic Activity

The effects of ethanolic extract of leaves of *A. dealbatum* on in-vitro clot lysis are showed in Table 2. It is evident that percentage of clot lysis was 75.69% ( $p < .001$ ) when 100 µl of streptokinase (1,50,000 I.U.) was used as a positive control, while in the case of water (negative control) the percentage of clot lysis was negligible (4.51%,  $p < .001$ ) and the extract (10 mg/kg) showed moderate potentiality (26.07%,  $p < .001$ ) compared with streptokinase.

**Table 1. Effects of ethanolic extract of leaves of *A. dealbatum* on diarrhea induced by castor oil in mice**

Groups	Dose	No. of feces	% of inhibition of defecation
Control	10 ml/kg	14.40±0.87###	
Standard	5 mg/kg	4.80±0.20**	66.67
EE	200 mg/kg	12.00±1.38#	16.67
	400 mg/kg	9.00±0.55** ,##	37.50

Here, EE stands for ethanolic extract and Data are presented as mean ± S.E.M. ANOVA was employed, followed by Dunnett's test and significant differences were represented by \* $p < .05$ , \*\* $p < .01$ , \*\*\* $p < .001$  vs control group treated with . Tween 80 was employed as negative control and loperamide was employed as standard. # $p < .05$ , ## $p < .01$  and ### $p < .001$  in relation to the loperamide



**Fig. 1. Effect of ethanolic extract of leaves of *A. dealbatum* (200 mg/kg and 400 mg/kg) with positive and negative control on % inhibition of defecation**

**Table 2. Effects of ethanolic extracts of leaves of *A. dealbatum* leaves on *in-vitro* clot lysis**

Treatment		% of clot lysis for human blood
Control		4.51 ± 0.02 <sup>####</sup>
Streptokinase	100 µl	75.69 ± 0.54 <sup>***</sup>
EE	10 mg/ml	26.07 ± 0.28 <sup>***###</sup>

Here, EE stands for ethanolic extract and data was presented as mean ± SEM. ANOVA was employed, followed by Dunnett's test and significant differences were represented by \* $p < .05$ , \*\* $p < .01$ , \*\*\* $p < .001$  vs control group treated with vehicle. Distilled water was employed as negative control and streptokinase was employed as positive control. # $p < .05$ , ## $p < .01$  and #### $p < .001$  in relation to the Streptokinase

#### 4. DISCUSSION

Abnormally frequent defecation of feces of low consistency which may be due to a disturbance in the transport of water and electrolytes in the intestines are called diarrhea. Instead of the multiplicity of etiologies, (i) increased electrolytes secretion (secretory diarrhea), (ii) increased luminal osmolarity (osmotic diarrhea), (iii) deranged intestinal motility causing a decreased transit time, and (iv) decreased electrolytes absorption may be responsible for pathophysiology [25,26]. Nitric oxide and ricinoleic acid is the most active component of castor oil which is responsible for diarrhea [26,27]. Inhibition of intestinal Na<sup>+</sup> K<sup>+</sup> ATPase activity, consequently reducing normal fluid absorption, activation of adenylate cyclase or mucosal cAMP-mediated active secretion [28] and stimulation of prostaglandin formation and platelet activating factor [29] are several

proposed mechanisms to expound the castor oil induced diarrheal effect [30,31]. Inflammatory mediators (e.g., prostaglandins and histamine) are secreted due to irritation and inflammation in the intestinal mucosa in the presence of ricinoleic acid in the gut. The released prostaglandins commence vasodilatation, smooth muscle contraction, and mucus secretion in the small intestines. In experimental animals as well as in human beings, prostaglandins of the E series are envisaged to be strong diarrheagenic agents. Our study showed that the overall antidiarrheal study reveals the dose dependent activity. All mice from the control group (treated with vehicle) produced diarrhea after castor oil administration. The decrease in the severity of the diarrhea was measured by the percent of inhibition of defecation. In our study, ethanolic extracts of *A. dealbatum* leaves showed moderately reduced amount of feces in castor oil-induced mice and % inhibition of defecation was 16.67 and 37.50 at

200 and 400mg/kg respectively. The values were increased as the dose had been increased and showed significant antidiarrheal effect compared with positive control-loperamide. From these results, it can be enumerated that water and electrolytes secretion into the small intestine are reduced and may ameliorate electrolyte absorption from the intestinal lumen consistent with inhibition of hypersecretion [32]. Besides different pathophysiological conditions of diarrhea, hypermotility characterizes diarrhea where the secretory component is not the causative factor [33]. It was reported that some phytochemical compounds like flavonoids and polyphenols were responsible for the antidiarrheal activity properties [34]. However, studies also have shown that flavonoids have ability to inhibit intestinal motility, water and electrolytes secretion [35] and inhibit prostaglandins biosynthesis which are considered to delay castor oil-induced diarrhea [36]. So, the antidiarrheal activity of the ethanolic extract of the leaves of *A. dealbatum* could therefore be due to the presence of flavonoids and phenols. The result was in concord with other species of same family. The damaged regions of the endothelial cell surface or blood vessel are blocked by the deposition of platelets, tissue factor and fibrin is called thrombosis or blood clot formation [37]. In the formation process platelets played the major role and thrombosis is initiated when the activated platelets form platelets to platelets bonds and further bind to the leucocytes and bring them into a complex process of plaque formation and growth [38]. It is the thrombolytic agents which working by disrupting the fibrinogen and fibrin contained in a clot. Plasmin is one of the natural anti-thrombotic agents. After a long process of trial and error several thrombolytic drugs are discovered from various sources. Under this study, we tried to find whether the herbal preparation of *A. dealbatum* leaves possess clot lysis potentiality or not. The percent clot lytic activity was compared with water (negative control) and standard enzyme streptokinase (positive control). The mean % of clot lysis for water and streptokinase was found 4.51% ( $p < .001$ ) and 75.69% ( $p < .001$ ) separately. 10 mg/ml extracts of *A. dealbatum* leaves was give 26.07% ( $p < .001$ ) clot lytic activity which is moderate effect compare with the positive and negative control. The result agrees with previous reports where extract of *Amomum subulatum* exhibited fibrinolytic effect [39]. It was narrated that some phytochemicals like saponin, alkaloids and tannin are endangered for thrombolytic

activity [40]. Therefore the possibility of the presence of these phytochemicals in the leaves extract may be the probable reason of demonstrating the thrombolytic activity.

## 5. CONCLUSION

To the best of our knowledge, this is the first report about evaluation of *in vivo* antidiarrheal and *in vitro* thrombolytic activity of ethanolic extract of leaves of *A. dealbatum*. These findings suggest that the plant may be a potential source for the development of new antidiarrheal drug. Also the obtained results confirmed the presence of thrombolytic element in the leaves of *A. dealbatum*. However, further investigations are required to isolate the active constituents responsible for the observed effect and to elucidate the possible mechanisms of action responsible for the anti-diarrheal and thrombolytic activities of this plant.

## ETHICAL APPROVAL

The study protocol was approved by the ethical Committee, Department of Pharmacy, International Islamic University Chittagong, Bangladesh (Pharm-P&D-37/07'12).

## AVAILABILITY OF DATA AND MATERIALS

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Sofowora A. Medicinal plants and traditional medicine in Africa. Chichester, West Sussex New York: Wiley. 1982;xviii: 256.
2. Keshari P. Pradeep. A review of conservation and sustainable use of medicinal plant with special reference of *Tecomella undulata* (Sm.) Seem. Journal of Pharmacognosy and Phytochemistry. 2018;3:09-13.
3. Akinmoladun AC, et al. Phytochemical constituent and antioxidant activity of extract from the leaves of *Ocimum*

- gratissimum*. Scientific Research and Essay. 2007;2(5):163-166.
4. Phuaklee P, Sakpakdeejaroen I, Itharat A. Cytotoxic and antioxidant activities of two species of ginger extracts. Thai Journal of Pharmacology. 2010;32(1):82-85.
  5. Taylor L, Taylor TN, Krings M. Paleobotany: The Biology and evolution of fossil plants. 2<sup>nd</sup> ed. Academic Press; 2009.
  6. Fern K. Useful Tropical Plants; 2014.
  7. Uddin SB. Bangladesh ethnobotany online database. Department of Botany, Chittagong University: Chittagong 4331, Bangladesh; 2012.
  8. Ezekwesili CN, Obiora KA, Ugwu OP. Evaluation of anti-diarrhoeal property of crude aqueous extract of *Ocimum gratissimum* L. (Labiatae) in rats. Nigerian Society for Experimental Biology. 2004; 16(2):122-131.
  9. Birru EM, et al. Antidiarrheal activity of crude methanolic root extract of *Idigofera spicata* Forssk.(Fabaceae). BMC Complement Altern Med. 2016;16:272.
  10. Kabir MS, et al. Antioxidant, antidiarrheal, hypoglycemic and thrombolytic activities of organic and aqueous extracts of *Hopea odorata* leaves and in silico PASS prediction of its isolated compounds. BMC Complement Altern Med. 2016;16(1):474.
  11. Nigam V, Paarakh PM. Evaluation of anti-diarrhoeal activity of hydro alcoholic extract of *Chenopodium album* L. Indian Journal of Natural Products and Resources 4, 61-66., Evaluation of anti-diarrhoeal activity of hydro alcoholic extract of *Chenopodium album* L. Indian Journal of Natural Products and Resources. 2013;4(1):61-66.
  12. Shoba FG, Thomas M. Evaluation of anti-diarrhoeal effect of four medicinal plants on castor oil-induced gastrointestinal motility in mice. Advances in Applied Science Research. 2014;5(4):153-156.
  13. Degu A, Engidawork E, Shibeshi W. Evaluation of the anti-diarrheal activity of the leaf extract of *Croton macrostachyus* Hocsht. ex Del. (Euphorbiaceae) in mice model. BMC Complement Altern Med. 2016;16(1):379.
  14. Konaté K, et al. Antidiarrheal and antimicrobial profiles extracts of the leaves from *Trichilia emetica* Vahl. (Meliaceae). Asian Pacific Journal of Tropical Biomedicine. 2015;5(3):242-248.
  15. Abdel-Razik A, et al. De-novo portal vein thrombosis in liver cirrhosis: Risk factors and correlation with the model for end-stage liver disease scoring system. Eur J Gastroenterol Hepatol. 2015;27(5):585-92.
  16. Qin J, et al. A panel of microRNAs as a new biomarkers for the detection of deep vein thrombosis. J Thromb Thrombolysis. 2015;39(2):215-21.
  17. Mucklow JC. Thrombolytic treatment. Streptokinase is more economical than alteplase. BMJ. 1995;311(7018):1506.
  18. Naderi GA, et al. Fibrinolytic effects of Ginkgo biloba extract. Exp Clin Cardiol. 2005;10(2):85-7.
  19. Rahman MA, et al. Effects of organic extracts of six Bangladeshi plants on in vitro thrombolysis and cytotoxicity. BMC Complement Altern Med. 2013;13:25.
  20. Dragstedt LR. Ethical considerations in the use and care of laboratory animals. J Med Educ. 1960;35:1-3.
  21. Franca CS, et al. Analgesic and antidiarrheal properties of *Ocimum selloi* essential oil in mice. Fitoterapia. 2008; 79(7-8):569-73.
  22. Abdullahi M, et al. Medicinal and economic plants of Nupeland. Bida, Nigeria: Jube Evans Books and Publications; 2003.
  23. Prasad S, et al. Development of an *in vitro* model to study clot lysis activity of thrombolytic drugs. Thromb J. 2006;4: 14.
  24. Kawsar MH, et al. Studies of thrombolytic and cytotoxic properties of two asteraceous plants of Bangladesh. Bangladesh Pharmaceutical Journal. 2011; 14(2):103-106.
  25. Agbor GA, Leopold T, Jeanne NY. The antidiarrhoeal activity of *Alchornea cordifolia* leaf extract. Phytother Res. 2004; 18(11):873-6.
  26. Umer S, Tekewe A, Kebede N. Antidiarrhoeal and antimicrobial activity of *Calpurnia aurea* leaf extract. BMC Complement Altern Med. 2013;13:21.
  27. Racusen LC, Binder HJ. Ricinoleic acid stimulation of active anion secretion in colonic mucosa of the rat. J Clin Invest. 1979;63(4):743-9.
  28. Pinto A, et al. Time course of PAF formation by gastrointestinal tissue in rats after castor oil challenge. J Pharm Pharmacol. 1992;44(3):224-6.
  29. Mascolo N, et al. Nitric oxide and castor oil-induced diarrhea. J Pharmacol Exp Ther. 1994;268(1):291-5.
  30. Capasso F, et al. Dissociation of castor oil-induced diarrhoea and intestinal mucosal injury in rat: Effect of NG-nitro-L-arginine

- methyl ester. Br J Pharmacol. 1994;113(4): 1127-30.
31. Zafar Imam M, Sultana S, Akter S. Antinociceptive, antidiarrheal, and neuropharmacological activities of *Barringtonia acutangula*. Pharm Biol. 2012;50(9):1078-84.
  32. Shah S. Evaluation of diarrhea: The challenge continues! Part-I. Indian J Med Sci. 2004;58(2):75-8.
  33. Chitme HR, Chandra M, Kaushik S. Studies on anti-diarrhoeal activity of *Calotropis gigantea* R.Br. in experimental animals. J Pharm Pharm Sci. 2004;7(1): 70-5.
  34. Dosso K, et al. Antidiarrhoeal activity of an ethanol extract of the stem bark of *Piliostigma reticulatum* (Caesalpiniaceae) in rats. Afr J Tradit Complement Altern Med. 2012;9(2):242-9.
  35. Di Carlo G, et al. Inhibition of intestinal motility and secretion by flavonoids in mice and rats: Structure-activity relationships. J Pharm Pharmacol. 1993;45(12):1054-9.
  36. Brijesh S, et al. Studies on the antidiarrhoeal activity of *Aegle marmelos* unripe fruit: validating its traditional usage. BMC Complement Altern Med. 2009;9:47.
  37. Palta S, Saroa R, Palta A. Overview of the coagulation system. Indian J Anaesth. 2014;58(5):515-23.
  38. Das A, et al. Investigation of *in vitro* thrombolytic potential of ethanolic extract of *Momordica charantia* fruits: An anti-diabetic medicinal plant. Der Pharmacia Sinica. 2013;4:104-108.
  39. Kumar Verma S, Jain V, Singh DP. Effect of greater cardamom (*Amomum subulatum* Roxb.) on blood lipids, fibrinolysis and total antioxidant status in patients with ischemic heart disease. Asian Pacific Journal of Tropical Disease. 2012;2(2):739-743.
  40. Ali MR, et al. Preliminary phytochemical screening and *in vitro* thrombolytic potential of the methanolic extract of *enhydra fluctuans* Lour (Leaves). International Journal of Pharmamedix India. 2013;1(2):270-280.

© 2019 Islam et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

The peer review history for this paper can be accessed here:  
<http://www.sdiarticle3.com/review-history/49046>