



Enterovirus-A76 of South-East Asian Ancestry in a Captive Chimpanzee (*Pan troglodytes*) in Jos, Nigeria

A. O. Oragwa^{1,2}, U. E. George¹, T. O. C. Faleye^{1,3}, M. O. Adewumi^{1*}
and J. A. Adeniji^{1,4}

¹Department of Virology, College of Medicine, University of Ibadan, Ibadan, Oyo State, Nigeria.

²Department of Veterinary Microbiology and Pathology, Faculty of Veterinary Medicine, University of Jos, Jos, Plateau State, Nigeria.

³Department of Microbiology, Faculty of Science, Ekiti State University, Ado-Ekiti, Ekiti State, Nigeria.

⁴WHO, National Polio Laboratory, University of Ibadan, Ibadan, Oyo State, Nigeria.

Authors' contributions

This work was carried out in collaboration between all authors. All authors designed the study. Author AOO wrote the first draft of the manuscript. Authors MOA and JAA supervised the study. All authors managed the analyses of the study, read and approved the final manuscript.

Article Information

DOI: 10.9734/JAMB/2018/40008

Editor(s):

(1) Pongsak Rattanachaiakunsonon, Professor, Department of Biological Science, Faculty of Science, Ubon Ratchathani University, Thailand.

Reviewers:

(1) El-Yuguda Abdul-Dahiru, University of Maiduguri, Nigeria.

(2) Artemiy Sergeev, Russia.

(3) M. Zamri Saad, Universiti Putra Malaysia, Malaysia.

Complete Peer review History: <http://www.sciencedomain.org/review-history/23945>

Original Research Article

Received 11th January 2018

Accepted 20th March 2018

Published 2nd April 2018

ABSTRACT

Background: Contact points between humans and nonhuman primates (NHPs) during the past decades have increased significantly in Africa. This provides opportunities for largescale emergence of novel virus types and species with unpredictable pathogenicity and clinical impacts. This study was designed to investigate and catalogue Enteroviruses (EVs) present in captive NHPs in Nigeria.

Methods: Twenty-seven (27) fecal samples collected from captive NHPs in a Wild Life Park and Zoological garden at Jos, Nigeria in April 2016 were analyzed in this study. Samples were resuspended in a phosphate buffered saline (PBS)/chloroform mixture, and the clarified supernatant was subjected to RNA extraction, cDNA synthesis, a Panenterovirus 5'-UTR assay and three different enterovirus VP1 snPCR assays. All amplicons from the snPCR assays were sequenced, and enteroviruses identified using the enterovirus genotyping tool and phylogenetic analysis.

*Corresponding author: E-mail: adewumi1@hotmail.com;

Results: Eight (29.63%) (two each from Chimpanzees, Patas Monkey, Mona Monkey and Baboon) of the 27 samples were positive for the 5'-UTR assay. One (3.70%) of the 27 samples was positive for the enterovirus VP1 snPCR assays in addition to its positivity by 5'-UTR assay. The same sample happens to be one of the two samples from Chimpanzees that tested positive for the 5'-UTR assay, and it was subsequently identified as EV-A76 of South-East Asia ancestry.

Conclusions: This study documents the first recorded attempt to detect and identify enteroviruses in NHPs in Nigeria. It also reports the first detection and identification of EV-A76 in Nigeria and particularly in a NHP. It is of utmost importance that the enterovirus VP1 assays be improved to enable detection of EVs that have been detected in NHPs but yet to be described in humans.

Keywords: Nonhuman primates; Nigeria; enteroviruses; EV-A76; chimpanzee.

1. INTRODUCTION

Enteroviruses are non-enveloped viruses with icosahedral symmetry, a diameter of 27–30 nm and a single-stranded, positive sense, RNA genome of ~7,500nt. They are members of the genus *Enterovirus* in the family *Picornaviridae*, order *Picornvirales*. Within the genus are twelve species which include: EV-A to -H, EV-J and *Rhinovirus A* (RV-A) to -C (<http://ictvonline.org/virusTaxonomy.asp?vesion=2012>). Some species are mainly populated by EVs infecting humans e.g. EV-A to -D and RV-A to -C, while others contain predominantly EVs infecting nonhuman primates (NHPs) e.g. EV-H, and EV-J. Since the establishment of a correlation between the nucleotide sequence of one of the structural genes (VP1) and serotypes [1], it has been used for enterovirus serotype/genotype determination.

Though there has been some effort to document circulation and genetic diversity of EVs infecting African NHPs [2-4], no information exist about the situation in Nigeria. Nigeria is rich in rain and mangrove forest and home to a wide range of Old World primate species, including apes (chimpanzee and gorillas) and other species of Old World monkeys. Hence, it is important to document circulation and genetic diversity of EVs infecting NHPs in Nigeria since they may harbor enteroviruses capable of cross-species transmission [2-5], and even the polioviruses [6].

Recent findings have shown the presence of EVs which were long considered to be primarily human viruses in faecal samples collected from wild and captive NHPs [2,5,6]. This therefore suggests that NHPs may serve as reservoirs for EVs and mixing vessels for the emergence of recombinant strains. Furthermore, we recently detected EV-A119 [7] and EV-B111 [8], in Nigerian children diagnosed with acute flaccid paralysis (AFP), which have been previously shown to co-circulate between NHPs and

humans [2,3,5,6,9,10]. This therefore suggests that EVs may also co-circulate between NHPs and humans in Nigeria. This study was therefore designed to investigate and catalogue EVs present in captive NHPs in Nigeria as a first step towards describing the genetic diversity of EVs in NHPs in the country.

2. METHODOLOGY

2.1 Sample Collection and Storage

Twenty-seven fecal samples collected from NHPs in April 2016 were analyzed in this study. The samples were collected non-invasively by properly trained NHP handlers from the cages of NHPs in a Wild Life Park and Zoological garden in Jos, North-Central Nigeria. About 5g of stool was collected from freshly deposited stools on the floor of the animal cages (Table 1) and placed in a sterile container. Samples were properly labeled and transported in cold chain to the enterovirus research laboratory at the Department of Virology, College of Medicine, University College Hospital, Ibadan, Nigeria and preserved at -20°C until analysis.

2.2 Sample Processing

About one gram of each stool specimen was diluted in 3mL phosphate buffered saline (PBS), one milliliter chloroform and one gram of glass beads. The mixture was vortexed for two (2) minutes and subsequently centrifuged at 3000 rpm for 20 minutes. Afterwards, 2mL of the supernatant was aliquoted in one milliliter volumes into two cryovials per sample. One vial was stored at -20°C while the other was analyzed further.

2.3 RNA Extraction and cDNA Synthesis

RNA was extracted from 100 µL of each of the 27 samples using Jena Bioscience RNA extraction kit (Jena Bioscience, Jena, Germany) and was used for cDNA synthesis using Script

cDNA Synthesis Kit (Jena Bioscience, Jena, Germany) as previously described [11,12]. To be precise, two different cDNA (one with random hexamers [11], and the other with primers AN32-AN35 [12]) were made in 10 µL volumes, for all the samples. This mixture was incubated at 42°C for 10min followed by 50°C for 60 minutes in a Veriti thermocycler (Applied Biosystems, California, USA).

2.4 Polymerase Chain Reaction (PCR) Screens

2.4.1 Polymerase chain reaction (PCR) assay for the 5' UTR

The 5'-UTR assay was done in 30µL volume. The cDNA made with random hexamers was used for this assay. Every PCR tube contained 10µL of cDNA, 6 µL of Red Load Taq (Jena Bioscience, Jena, Germany), 13.4 µL of RNase free water and 0.3 µL each of the forward (PanEnt-5'-UTR-F) and reverse (PanEnt-5'-UTR-R) primers [11]. A Veriti thermal cycler (Applied Biosystems, California, USA) was used for thermal cycling as follows; 94°C for 3 minutes, then 45 cycles of 94°C for 30 seconds, 42°C for 30 seconds, and 60°C for 30 seconds, with ramp of 40% from 42°C to 60°C. This was then followed by 72°C for 7 minutes, and held at 4°C until the reaction was terminated. Subsequently, PCR products were resolved on 2% agarose gel stained with ethidium bromide and viewed using a UV transilluminator.

2.5 Seminested Polymerase Chain Reaction (snPCR) Assay for the VP1 Gene

As previously described [12,13], an snPCR assay was done in this case. The assay is a modification [12,13] of the WHO recommended RT-snPCR assay for direct detection of enteroviruses from clinical specimen [14]. A single first round PCR was done. Followed by three (PanEnterovirus [PE], Enterovirus Species A or C [EV-A/C] and Enterovirus Species B [EV-B]) different second round PCR assays. The cDNA made with primers AN32-AN35 was used for this assay. Each reaction of the first round assay was also done in 30 µL volume and contained 10 µL of cDNA, 6 µL of Red Load Taq (Jena Bioscience, Jena, Germany), 13.4 µL of RNase free water and 0.3 µL each of the forward (224) and reverse (222) primers. For the three second round PCR assays, the reaction mix was constituted as above except for the cDNA that

was substituted with the first-round PCR product. Also, instead of using 10 µL of cDNA, 3 µL of first round PCR product was used, and the RNase free water was increased to 20.4 µL. In addition, the primers were also changed. Primer AN88 was reverse primer used for all three second round assays. However, the forward primers were AN89, 189 and 187 for PE, EV-A/C, EV-B assays, respectively.

The same Veriti thermal cycler (Applied Biosystems, California, USA) used for the 5'-UTR assay was also used for the VP1 snPCR assay. Thermal cycling conditions were as described for the 5'-UTR assay but with few changes. The extension times were 60 and 30 seconds for the first and second round assays respectively. All PCR products were also resolved on 2% agarose gel stained with ethidium bromide and viewed using a UV transilluminator.

2.6 Amplicon Sequencing and Enterovirus Identification

Only amplicons with expected band size emanating from the second round PCR assays for the VP1 gene were shipped to MacroGen Inc, Seoul, South Korea, for purification and nucleotide sequencing using the respective primer pairs. The enterovirus genotyping tool (EGT) [15] was subsequently used for enterovirus type determination.

2.7 Phylogenetic Analysis

For multiple sequence alignment of the sequence(s) generated in this study and those of reference sequences downloaded from Genbank, the CLUSTAL W program in MEGA 5 software [16] was used with default settings. A neighbor-joining tree was then constructed using the Kimura-2 parameter model [17] and 1,000 bootstrap replicates in the same MEGA 5 software.

Nucleotide Sequence Accession Numbers: The sequence generated in this study has been deposited in GenBank under accession number KY798125.

3. RESULTS

3.1 Polymerase Chain Reaction (PCR) Assay for the 5' UTR

Eight (29.63%) of the 27 samples subjected to this assay had the expected band size (~114bp). The eight bands were detected in samples from four different NHP types namely, Chimpanzees,

Patas Monkey, Mona Monkey and Baboon (Table 1). It is however crucial to note that while five (5) of the bands had very high intensity, the remaining three (3) bands were very weak.

3.2 Seminested Polymerase Chain Reaction (snPCR) Assay for the VP1 Gene

Only one of the 27 (3.70%) samples subjected to these three assays had the expected band size (~350bp). The sample was positive for all three second round PCR assays (Table 1). It is one of the two samples from Chimpanzees that had been previously positive for the 5'-UTR PCR assay.

3.3 Amplicon Sequencing and Enterovirus Identification

The EGT independently identified all three sequences derived from the three amplicons generated from VP1 assay positive sample as Enterovirus A76 (EV-A76).

3.4 Phylogenetic Analysis

Phylogenetic analysis showed that the EV-A76 detected in this study is not related to all EV-A76 strains previously detected in sub-Saharan Africa from 2003 till date. Rather it is a member of one of the EV-A76 lineages circulating in South-East Asia and shares a common ancestor with a strain recovered in 2009 from the stool of a child in India (Fig. 1).

4. DISCUSSION

To the best of the authors' knowledge, this study documents the first recorded attempt to detect and identify enteroviruses in NHPs in Nigeria. Here we report the first detection and identification of EV-A76 in Nigeria and particularly in a NHP. Studies [18-20] have documented the recovery of EV-A76 in humans with different clinical conditions ranging from gastroenteritis [20], to central nervous system (CNS) manifestations like acute flaccid paralysis (AFP) [19,21] and encephalitis [18]. Hence, EV-A76s potential to have serious clinical manifestations in humans is obvious. Also crucial is the fact that EV-A76 has also been previously recovered in NHPs in Cameroon [2,3], and Bangladesh [5]. In fact, it was unambiguously shown [5] and confirmed in this study (Fig. 1) that

in 2008, the same strain of EV-A76 was simultaneously circulating in both humans and NHPs in Bangladesh. This confirms the zoonotic potential of EV-A76 and lays more credence to the need to better understand its epidemiology.

Prior this finding of EV-A76 in a Chimpanzee in Nigeria, it had been previously described in two other countries in sub-Saharan Africa; Cameroon [2,3,21] and Congo DRC [22]. It is however important to emphasize that the EV-A76 strain detected in this study is not related to those previously detected in the sub-region. Rather, it belonged to one of the lineages that had been circulating in South-East Asia since year 2000. In fact, the EV-A76 strain detected in this study is most closely related to (shares a common ancestor) a strain detected in India in 2009. How the infected Chimpanzee got exposed and infected by this imported strain of EV-A76 is currently not clear. However, considering the Chimpanzee in question was born in the park (Table 1), and had been in captivity in the same park since birth, it must have been exposed to this EV-A76 strain at the park. The possibility therefore exists that this strain of EV-A76 might be circulating either in the park or the encompassing community. In fact, the topology of the relationship between the India 2009 strain and that detected in this study suggests that several isolates of EV-A76 that are similar to both are yet to be described. Hence, there is need for further investigation to better describe and catalogue the diversity of EV-A76 in the region and Nigeria at large.

In this study, the 5'-UTR PCR assay detected enteroviruses in 29.63% (8/27) of the samples analyzed. However, the VP1 gene was amplified in only one of the eight samples positive for the 5'-UTR PCR assay. The gene was unamplifiable in the remaining seven samples. Why this is the case is currently not clear. However, as previously suggested [4], the possibility exists that the VP1 gene could not be amplified in the other seven samples because they are not members of EV-A – EV-D for which the VP1 amplification snPCR assay was originally developed [23]. With the increased significance of enteroviruses in NHPs, it is of utmost importance that the WHO recommended assay [14] be further improved to be able to consistently amplify the VP1 gene of non-Species A – D enteroviruses.

Table 1. Profile of NHPs sampled and results of enterovirus detection and identification assays

S/No	Name of Animal	Scientific Name	Site of Sample Collection	No of Animals in the cage	Sex	Age (Yrs)	Age in Captivity (Yrs)	Source/ Origin	Result		
									5'UTR	VP-1	Serotype
1	Olive Baboon	Papio anubis	JWLP	2	M/F	3		Plateau			
2	Patas Monkey	Erythrocebus patas	JZ	1	M	>10	10				
3	Patas Monkey	Erythrocebus patas	JWLP	4	M1/F3	1, 1.5, 2, 2.5	1, 1.5, 2, 2.5	Bitp	Positive		
4	Tantalus Monkey	Chlorocebus tantalus	JZ	1	F	~10	~10	Benin			
5	Baboon	Papio anubis	JZ	1	M	5	5	Plateau			
6	Chimpanzee	Pan troglodytes	JWLP	2	F	8	8	Bitp	Positive	Positive	EV-A76
7	Mona Monkey	Cercopithecus mona	JWLP	3	M1/F2			Plateau/ Taraba	Weak Positive		
8	Chimpanzee	Pan troglodytes	JZ	1	F	~54	~52	Savanna/ Benin			
9	Patas Monkey	Erythrocebus patas	JWLP	1	F	>3	~3	Plateau			
10	Patas Monkey	Erythrocebus patas	JZ	2	M	17	17	Central Africa	Weak Positive		
11	Baboon	Papio anubis	JZ	1	F	4	4	Plateau	Positive		
12	Baboon	Papio anubis	JZ	2	M/F	4&6	4&6				
13	Chimpanzee	Pan troglodytes	JZ	2	M/F	13	13				
14	Olive Baboon	Papio anubis	JWLP	1	M	4	4	Plateau	Positive		
15	Tantalus Monkey	Chlorocebus tantalus	JZ	1	F	20	20	Benin			
16	Mona Monkey	Cercopithecus mona	JZ	1	F	10	6_10	Abuja			
17	Baboon	Papio anubis	JZ	1	F	15	4	Benin			
18	Tantalus Monkey	Chlorocebus tantalus	JZ	1	F	>13	13				
19	Tantalus Monkey	Chlorocebus tantalus	JWLP	2	F	>1	1	Plateau			
20	Mona Monkey	Cercopithecus mona	JZ	1	M	20	20	Benin	Weak Positive		
21	Tantalus Monkey	Chlorocebus tantalus	JZ	2	F	>13	>13				
22	Chimpanzee	Pan troglodytes	JZ	1	M	>20	>20	Benin	Positive		
23	Western Baboon	Papio hamadryas	JWLP	1	M	7	7	Lagos			
24	Patas Monkey	Erythrocebus patas	JZ	2	M/F	>13	>13				
25	Patas Monkey	Erythrocebus patas	JWLP	1	M	1.5	1.5	Bitp			
26	Tantalus Monkey	Chlorocebus tantalus	JZ	2	M	13_9	13_9				
27	Olive Baboon	Papio anubis	JWLP	1	M	7	7	Plateau			

NB: JWLP= Jos Wild Life Park., JZ= Jos Zoo, Bitp= Born in the park




Fig. 1. Phylogram of EV-A76 identified in this study. The phylogram is based on an alignment of partial VP1 sequences. The newly sequenced strain is highlighted with Black Square. Strains recovered from children with AFP in Congo DRC and Cameroon is indicated with a black diamond and triangle, respectively. Strains recovered from NHPs in Bangladesh are highlighted with black circle. All strains recovered from NHPs in Cameroon form a cluster and are highlighted as ‘CPZ CAMEROON’ in the Phylogram. The GenBank accession number of the strains are indicated in the phylogram. Bootstrap values are indicated if > 50%

5. CONCLUSIONS

In this study, we report the first detection and identification of EV-A76 in Nigeria and particularly in a NHP. The EV-A76 strain detected in this study was not related to those previously detected in the sub-region but rather, belonged to one of the lineages that had been circulating in South-East Asia since year 2000. This result underscores the need to determine the genetic diversity of enteroviruses circulating in both captive and wild NHPs in Nigeria, as well as in humans living in close proximity to estimate the exposure and potential dissemination of new enteroviruses among humans in the forested regions in Nigeria.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Oberste MS, Maher K, Flemister MR, Marchetti G, Kilpatrick DR, Pallansch MA. Comparison of classic and molecular approaches for the identification of untypeable enteroviruses. *Journal of Clinical Microbiology*. 2000;38(3):1170–1174
2. Harvala H, Sharp CP, Ngole EM, Delaporte E, Peeters M, Simmonds P. Detection and genetic characterization of enteroviruses circulating among wild populations of chimpanzees in Cameroon: Relationship with human and simian enteroviruses. *Journal of Virology*. 2011; 85(9):4480-4486.
3. Sadeuh-Mba SA, Bessaud M, Joffret ML, Endegue Zanga MC, Balanant J, Mpoudi Ngole E, Njoum R, Reynes JM, Delpeyroux F, Rousset D. Characterization of enteroviruses from non-human primates in Cameroon revealed virus types widespread in humans along with candidate new types and species. *PLoS Negl Trop Dis*. 2014;8(7):E3052.
4. Mombo IM, Lukashev AN, Bleicker T, BruĖnink S, Berthet N, Maganga GD, et al. African non-human primates host diverse enteroviruses. *PLoS ONE*. 2017;12(1): e0169067.
5. Oberste MS, Feeroz MM, Maher K, Nix WA, Engel GA, Hasan KM, Begum S, Oh G, Chowdhury AH, Pallansch MA, Jones-Engel L. Characterizing the picornavirus landscape among synanthropic nonhuman primates in Bangladesh, 2007 to 2008. *Journal of Virology*. 2013a;87:558–571.
6. Oberste MS, Feeroz MM, Maher K, Nix WA, Engel GA, Begum S, Hasan KM, Oh G, Pallansch MA, Jones-Engel L. Naturally acquired Picornavirus infections in primates at the Dhaka Zoo. *Journal of Virology*. 2013b;87(1):572–580.
7. Adeniji JA, Oragwa AO, George UE, Ibok UI, Faleye TOC, Adewumi MO. Enterovirus A119 in a child with Acute Flaccid Paralysis, Nigeria. *J Hum Virol. Retrovirol*. 2016;4(2):00126.
8. Adeniji JA, Oragwa AO, George UE, Ibok UI, Faleye TOC, Adewumi MO. Preponderance of Enterovirus species C in RD-L20B cell culture negative stool samples from children diagnosed with Acute Flaccid Paralysis in Nigeria. *Archives of Virology*. 2017;162(10):3089-3101.
9. Ayukekbong J, Kabayiza JC, Lindh M, Nkuo-Akenji T, Tah F, Bergstrom T, Norder H. Shift of Enterovirus species among children in Cameroon-Identification of a new enterovirus, EV-A119. *J. Clin. Virol*. 2013;58(1):227-232.
10. Di Cristianzano V, Boettcher S, Diedrich S, Timmen-Wego M, Knops E, Luebke N, Kaiser R, Pfister H, Kabore Y, D'Alfonso R. Detection and characterization of enteroviruses and parechoviruses in healthy people living in the South of Cote d'Ivoire. *J. Clin. Virol*. 2015;71:40-43.
11. Adeniji JA, Faleye TOC. Impact of cell lines included in Enterovirus isolation protocol on perception of Nonpolio Enterovirus species C diversity. *Journal of Virological Methods*. 2014;207:238-247.
12. Faleye TOC, Adewumi MO, Coker BA, Nudamajo FY, Adeniji JA. Direct detection and identification of Enteroviruses from Faeces of healthy Nigerian children using a cell-culture independent RT-Seminested PCR assay. *Advances in Virology*; 2016a. Article ID: 1412838, 12 pages. Available:<http://dx.doi.org/10.1155/2016/1412838>
13. Faleye TOC, Adewumi MO, Kareem SA, Adesuyan YO, Fapohunda FA, Fasanya ST, Jimeto T, Lawrence OE, Obembe AA, Adeniji JA. The impact of a panenterovirus VP1 assay on our perception of the enterovirus diversity landscape of a sample. *J Hum Virol. Retrovirol*. 2016b; 4(3):00134.

14. World Health Organisation. Enterovirus surveillance guidelines: Guidelines for enterovirus surveillance in support of the Polio Eradication Initiative, Geneva; 2015.
15. Kroneman A, Vennema H, Deforche K, Van der Avoort H, Pe narandac S, Oberste MS, Vinjéc J, Koopmans M. An automated genotyping tool for enteroviruses and noroviruses. *Journal of Clinical Virology*. 2011;51:121–125.
16. Tamura K, Peterson D, Peterson N, Stecher G, Nei M, Kumar S. MEGA5: Molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Molecular Biology and Evolution*. 2011;28:2731-2739.
17. Kimura M. A simple method for estimating evolutionary rate of base substitutions through comparative studies of nucleotide sequences. *Journal of Molecular Evolution*. 1980;16(2):111-120.
18. Kumar A, Shukla D, Kumar R, Idris MZ, Misra UK, Dhole TN. Molecular epidemiological study of enteroviruses associated with encephalitis in children from India. *J. Clin. Microbiol*. 2012;50(11): 3509-3512.
19. Rao CD, Yergolkar P, and Shankarappa KS. Antigenic diversity of enteroviruses associated with nonpolio acute flaccid paralysis, India, 2007-2009. *Emerging Infect. Dis*. 2012;18(11):1833-1840.
20. Patil PR, Chitambar SD, Gopalkrishna V. Molecular surveillance of non-polio enterovirus infections in patients with acute gastroenteritis in Western India: 2004-2009. *J. Med. Virol*. 2015;87(1):154-61.
21. Sadeuh-Mba SA, Bessaud M, Massenet D, Joffret ML, Endegue MC, Njouom R, Reynes JM, Rousset D, Delpeyroux F. High frequency and diversity of species C enteroviruses in Cameroon and neighboring countries. *J. Clin. Microbiol*. 2013;51(3):759-770.
22. Junttila N, Leveque N, Kabue JP, Cartet G, Mushiya F, Muyembe-Tamfum JJ, Trompette A, Lina B, Magnius LO, Chomel JJ, Norder H. New enteroviruses, EV-93 and EV-94, associated with acute flaccid paralysis in the Democratic Republic of the Congo. *J. Med. Virol*. 2007;79(4):393-400.
23. Nix WA, Oberste MS, Pallansch MA. Sensitive, Seminested PCR Amplification of VP1 Sequences for Direct Identification of All Enterovirus Serotypes from Original Clinical Specimens. *Journal of Clinical Microbiology*. 2006;44(8):2698–2704.

© 2018 Oragwa et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<http://www.sciencedomain.org/review-history/23945>